



Syzygies of Prym and paracanonical curves of genus 8

Elisabetta Colombo, Gavril Farkas, Alessandro Verra, and Claire Voisin

Abstract. By analogy with Green's Conjecture on syzygies of canonical curves, the Prym-Green conjecture predicts that the resolution of a general level p paracanonical curve of genus g is natural. The Prym-Green Conjecture is known to hold in odd genus for almost all levels. Probabilistic arguments strongly suggested that the conjecture might fail for level 2 and genus 8 or 16. In this paper, we present three geometric proofs of the surprising failure of the Prym-Green Conjecture in genus 8, hoping that the methods introduced here will shed light on all the exceptions to the Prym-Green Conjecture for genera with high divisibility by 2.

Keywords. Paracanonical curve; syzygy; genus 8; moduli of Prym varieties

2010 Mathematics Subject Classification. 13D02; 14H10

[Français]

Titre. Syzygies de Prym et courbes paracanoniques de genre 8

Résumé. Par analogie avec la conjecture de Green sur les syzygies des courbes canoniques, la conjecture de Prym-Green prédit que la résolution d'une courbe générale, paracanonique, de genre g et de niveau p est naturelle. Cette conjecture est connue en genre impair pour presque tout niveau. Des arguments probabilistes ont fortement suggéré qu'elle pourrait s'avérer fausse pour le niveau 2 en genre 8 et 16. Dans cet article, nous présentons trois démonstrations géométriques de la surprenante non-validité de la conjecture de Prym-Green en genre 8, en espérant que les méthodes introduites apporteront un éclairage nouveau sur toutes les exceptions à la conjecture de Prym-Green pour des genres divisibles par une grande puissance de 2.

Received by the Editors on December 15, 2016, and in final form on June 23, 2017.

Accepted on June 27, 2017.

Elisabetta Colombo

Università di Milano, Dipartimento di Matematica, Via Cesare Saldini 50, 20133 Milano, Italy

e-mail: elisabetta.colombo@unimi.it

Gavril Farkas

Humboldt-Universität zu Berlin, Institut für Mathematik, Unter den Linden 6, 10099 Berlin, Germany

e-mail: farkas@math.hu-berlin.de

Alessandro Verra

Università Roma Tre, Dipartimento di Matematica, Largo San Leonardo Murialdo, 1-00146 Roma, Italy

e-mail: verra@mat.uniroma3.it

Claire Voisin

Collège de France 3, Rue d'Ulm, 75005 Paris, France

e-mail: claire.voisin@imj-prg.fr

Contents

| | |
|---|----|
| 1. Introduction | 2 |
| 2. Syzygies of paracanonical curves of genus 8 | 4 |
| 3. First proof: reducible spin curves | 10 |
| 4. Second proof: Divisor class calculations on $\overline{\mathcal{R}}_g$ | 12 |
| 5. Rank 2 vector bundles and singular quartics | 16 |
| 6. Miscellany | 19 |

1. Introduction

By analogy with Green's Conjecture on the syzygies of a general canonical curve [18], [19], the Prym-Green Conjecture, formulated in [10] and [3], predicts that the resolution of a *paracanonical* curve

$$\phi_{K_C \otimes \eta} : C \hookrightarrow \mathbf{P}^{g-2},$$

where C is a general curve of genus g and $\eta \in \text{Pic}^0(C)[\ell]$ is an ℓ -torsion point is natural. For even genus $g = 2i + 6$, the Prym-Green Conjecture amounts to the vanishing statement

$$K_{i,2}(C, K_C \otimes \eta) = K_{i+1,1}(C, K_C \otimes \eta) = 0, \quad (1.1)$$

in terms of Koszul cohomology groups. Equivalently, the genus g paracanonical level ℓ curve $C \subseteq \mathbf{P}^{g-2}$ satisfies the Green-Lazarsfeld property (N_i) . The Prym-Green Conjecture has been proved for all *odd* genera g when $\ell = 2$, see [8], or $\ell \geq \sqrt{\frac{g+2}{2}}$, see [9]. For even genus, the Prym-Green Conjecture has been established by degeneration and using computer algebra tools in [3] and [4], for all $\ell \leq 5$ and $g \leq 18$, with two possible mysterious exceptions in level 2 and genus $g = 8, 16$ respectively. The last section of [3] provides various pieces of evidence, including a probabilistic argument, strongly suggesting that for $g = 8$, one has $\dim K_{1,2}(C, K_C \otimes \eta) = 1$, and thus the vanishing (1.1) fails in this case. It is tempting to believe that the exceptions $g = 8, 16$ can be extrapolated to higher genus, and that for genera g with high divisibility by 2, there are genuinely novel ways of constructing syzygies of Prym-canonical curves waiting to be discovered. It would be very interesting to test experimentally the next relevant case $g = 24$. Unfortunately, due to memory and running time constraints, this is currently completely out of reach, see [3] and [7].

The aim of this paper is to confirm the expectation formulated in [3] and offer several geometric explanations for the surprising failure of the Prym-Green Conjecture in genus 8, hoping that the geometric methods described here for constructing syzygies of Prym-canonical curves will eventually shed light on all the exceptions to the Prym-Green Conjecture. We choose a general Prym-canonical curve of genus 8

$$\phi_{K_C \otimes \eta} : C \hookrightarrow \mathbf{P}^6,$$

with $\eta^{\otimes 2} = \mathcal{O}_C$. Set $L := K_C \otimes \eta$ and denote $I_{C,L}(k) := \text{Ker}\{\text{Sym}^k H^0(C, L) \rightarrow H^0(C, L^{\otimes k})\}$ for all $k \geq 2$. Observe that $\dim I_{C,L}(2) = \dim K_{1,1}(C, L) = 7$ and $\dim I_{C,L}(3) = 49$, therefore as $[C, \eta]$ varies in moduli, the multiplication map

$$\mu_{C,L} : I_{C,L}(2) \otimes H^0(C, L) \rightarrow I_{C,L}(3)$$

globalizes to a morphism of vector bundles of the same rank over the stack \mathcal{R}_8 classifying pairs $[C, \eta]$, where C is a smooth curve of genus 8 and $\eta \in \text{Pic}^0[2] \setminus \{\mathcal{O}_C\}$.

Theorem 1. *For a general Prym curve $[C, \eta] \in \mathcal{R}_8$, one has $K_{1,2}(C, L) \neq 0$. Equivalently the multiplication map $\mu_{C,L} : I_{C,L}(2) \otimes H^0(C, L) \rightarrow I_{C,L}(3)$ is not an isomorphism.*

We present three different proofs of Theorem 1. The first proof, presented in Section 3 uses the structure theorem already pointed out in [3] for degenerate syzygies of paracanonical curves in \mathbf{P}^6 . Precisely, if a paracanonical genus 8 curve $\phi_{K_C \otimes \eta} : C \hookrightarrow \mathbf{P}^6$, where $\eta \neq \mathcal{O}_C$, has a syzygy $0 \neq \gamma \in K_{1,2}(C, K_C \otimes \eta)$ of *sub-maximal rank* (see Section 2 for a precise definition), then the syzygy scheme of γ consists of an isolated point $p \in \mathbf{P}^6 \setminus C$ and a residual septic elliptic curve $E \subseteq \mathbf{P}^6$ meeting C transversally along a divisor e of degree 14, such that if e is viewed as a divisor on C and E respectively, then

$$e_C \in |K_C \otimes \eta^{\otimes 2}| \quad \text{and} \quad e_E \in |\mathcal{O}_E(2)|. \quad (1.2)$$

The union $D := C \cup E \hookrightarrow \mathbf{P}^6$, endowed with the line bundle $\mathcal{O}_D(1)$ is a degenerate spin curve of genus 22 in the sense of [5]. The locus of stable spin structures with at least 7 sections defines a subvariety of codimension $21 = \binom{7}{2}$ inside the moduli space $\overline{\mathcal{S}}_{22}$ of stable odd spin curves of genus 22. By restricting this condition to the locus of spin structures having $D := C \cup_e E$ as underlying curve, it turns out that one has enough parameters to realize this condition for a general $C \subseteq \mathbf{P}^6$ if and only if

$$\dim |K_C \otimes \eta^{\otimes 2}| = 7,$$

which happens precisely when $\eta^{\otimes 2} \cong \mathcal{O}_C$. Therefore for each Prym-canonical curve $C \subseteq \mathbf{P}^6$ of genus 8 there exists a corresponding elliptic curve $E \subseteq \mathbf{P}^6$ such that the intersection divisor $E \cdot C$ verifies (1.2), which forces $K_{1,2}(C, K_C \otimes \eta) \neq 0$.

The second and the third proofs involve the reformulation given in Section 2.B (see Proposition 5) of the condition that a paracanonical curve $\phi_L : C \hookrightarrow \mathbf{P}^6$ have a non-trivial syzygy. Precisely, if $\phi_L(C)$ is scheme-theoretically generated by quadrics, then $K_{1,2}(C, L) \neq 0$, if and only if there exists a quartic hypersurface in \mathbf{P}^6 singular along $C \subseteq \mathbf{P}^6$, which is not a quadratic polynomial in quadrics vanishing along C , that is, it does not belong to the image of the multiplication map

$$\mathrm{Sym}^2 I_{C,L}(2) \rightarrow I_{C,L}(4).$$

Equivalently, one has $H^1(\mathbf{P}^6, \mathcal{I}_{C/\mathbf{P}^6}^2(4)) \neq 0$.

The second proof presented in Section 4 uses intersection theory on the stack $\overline{\mathcal{R}}_8$. The virtual Koszul divisor of Prym curves $[C, \eta] \in \mathcal{R}_8$ having $K_{1,2}(C, K_C \otimes \eta) \neq 0$, splits into two divisors \mathfrak{D}_1 and \mathfrak{D}_2 respectively, corresponding to the case whether $C \subseteq \mathbf{P}^6$ is not scheme-theoretically cut out by quadrics, or $H^1(\mathbf{P}^6, \mathcal{I}_{C/\mathbf{P}^6}^2(4)) \neq 0$ respectively. We determine the virtual classes of both closures $\overline{\mathfrak{D}}_1$ and $\overline{\mathfrak{D}}_2$. Using an explicit uniruled parametrization of $\overline{\mathcal{R}}_8$ constructed in [11], we conclude that the class $[\overline{\mathfrak{D}}_2] \in CH^1(\overline{\mathcal{R}}_8)$ cannot possibly be effective (see Theorem 20). Therefore, again $K_{2,1}(C, K_C \otimes \eta) \neq 0$, for every Prym curve $[C, \eta] \in \mathcal{R}_8$.

The third proof given in Section 5 even though subject to a plausible, but still unproved transversality assumption, is constructive and potentially the most useful, for we feel it might offer hints to the case $g = 16$ and further. The idea is to consider rank 2 vector bundles E on C with canonical determinant and $h^0(C, E) = h^0(C, E(\eta)) = 4$. (Note that the condition that η is 2-torsion is equivalent to the fact that $E(\eta)$ also has canonical determinant, which is essential for the existence of such nonsplit vector bundles, cf. [15].) By pulling back to C the determinantal quartic hypersurface consisting of rank 3 tensors in

$$\mathbf{P} \left(H^0(C, E)^\vee \otimes H^0(C, E(\eta))^\vee \right) \cong \mathbf{P}^{15}$$

under the natural map $H^0(C, K_C \otimes \eta)^\vee \rightarrow H^0(C, E)^\vee \otimes H^0(C, E(\eta))^\vee$, we obtain explicit quartic hypersurfaces singular along the curve $C \subseteq \mathbf{P}^6$. Our proof that these are not quadratic polynomials

into quadrics vanishing along the curve, that is, they do not lie in the image of $\text{Sym}^2 I_{C,L}(2)$ remains incomplete, but there is a lot of evidence for this.

The methods of Section 5 suggests the following analogy in the next case $g = 16$. If $[C, \eta] \in \mathcal{R}_{16}$ is a Prym curve of genus 16, there exist vector bundles E on C with $\det E \cong K_C$ and satisfying $h^0(C, E) = h^0(C, E(\eta)) = 6$. Potentially they could be used to prove that $K_{5,2}(C, K_C \otimes \eta) \neq 0$ and thus confirm the next exception to the Prym-Green Conjecture.

2. Syzygies of paracanonical curves of genus 8

Let C be a general smooth projective curve of genus 8. For a non-trivial line bundle $\eta \in \text{Pic}^0(C)$, we shall study the *paracanonical* line bundle $L := K_C \otimes \eta$. When η is a 2-torsion point, we speak of the *Prym-canonical* line bundle L . For each paracanonical bundle L , we have $h^0(C, L) = 7$ and an induced embedding

$$\phi_L : C \hookrightarrow \mathbf{P}^6.$$

The goal is to understand the reasons for the non-vanishing of the Koszul group $K_{1,2}(C, L)$ of a Prym-canonical bundle L , as suggested experimentally by the results of [3], [4].

Let $I_C(2) = I_{C,L}(2) \subseteq H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(2))$, respectively $I_C(3) = I_{C,L}(3) \subseteq H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(3))$ be the ideal of quadrics, respectively cubics, vanishing on $\phi_L(C)$. It is well-known that whenever L is projectively normal, the non-vanishing of the Koszul cohomology group $K_{1,2}(C, L)$ is equivalent to the non-surjectivity of the multiplication map

$$\mu_{C,L} : H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \otimes I_C(2) \rightarrow I_C(3). \quad (2.3)$$

Note that

$$\dim I_C(2) = \binom{8}{2} - 21 = 7, \quad \text{and} \quad \dim I_C(3) = \binom{9}{3} - 3 \cdot 14 + 7 = 49,$$

respectively, so that the two spaces appearing in the map (2.3) have the same dimension. Denote by P_8^{14} the universal degree 14 Picard variety over \mathcal{M}_8 consisting of pairs $[C, L]$, where $[C] \in \mathcal{M}_8$ and $L \neq K_C$. The jumping locus

$$\mathfrak{R}053 := \left\{ [C, L] \in P_8^{14} : K_{1,2}(C, L) \neq 0 \right\}$$

is a divisor. It turns out, cf. Theorem 5.3 of [3] and Proposition 8, that $\mathfrak{R}053$ splits into two components depending on the *rank* of the corresponding non-zero syzygy from $K_{1,2}(C, L)$.

Definition 2. The rank of a non-zero syzygy $\gamma = \sum_{i=0}^6 \ell_i \otimes q_i \in \text{Ker}(\mu_{C,L})$ is the dimension of the subspace $\langle \ell_0, \dots, \ell_6 \rangle \subseteq H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1))$. The syzygy scheme $\text{Syz}(\gamma)$ of γ is the largest subscheme $Y \subseteq \mathbf{P}^6$ such that $\gamma \in H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \otimes I_Y(2)$.

It is shown in [3], that $\mathfrak{R}053$ splits into divisors $\mathfrak{R}053_6$ and $\mathfrak{R}053_7$, depending on whether the syzygy $0 \neq \gamma \in \text{Ker}(\mu_{C,L})$ has rank 6 or 7 respectively. By a specialization argument to irreducible nodal curves, it follows from [3] that $\mathcal{R}_8 \not\subseteq \mathfrak{R}053_7$. A direct, more transparent proof of this fact will be given in Proposition 13.

2.A. Paracanonical curves of genus 8 with special syzygies and elliptic curves

We summarize a few facts already stated or recalled in Section 5 of [3] concerning rank 6 syzygies of paracanonical curves in \mathbf{P}^6 . Very generally, let

$$\gamma = \sum_{i=1}^6 \ell_i \otimes q_i \in H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \otimes H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(2))$$

be a rank 6 linear syzygy among quadrics in \mathbf{P}^6 . The linear forms ℓ_1, \dots, ℓ_6 define a point $p \in \mathbf{P}^6$. Following Lemma 6.3 of [16], there exists a skew-symmetric matrix of linear forms $A := (a_{ij})_{i,j=1,\dots,6}$, such that

$$q_i = \sum_{j=1}^6 \ell_j a_{ij}.$$

In the space \mathbf{P}^{20} with coordinates ℓ_1, \dots, ℓ_6 and a_{ij} for $1 \leq i < j \leq 6$, one considers the 15-dimensional variety X_6 defined by the 6 quadratic equations $\sum_{j=1}^6 \ell_j a_{ij} = 0$, where $i = 1, \dots, 6$ and by the cubic equation $\text{Pfaff}(A) = 0$ in the variables a_{ij} . The original space \mathbf{P}^6 embeds in \mathbf{P}^{20} via evaluation. The syzygy scheme $\text{Syz}(\gamma)$ is the union of the point p and of the intersection D of \mathbf{P}^6 with the variety X_6 . It follows from Theorem 4.4 of [6], that for a general rank 6 syzygy γ as above, $D \subseteq \mathbf{P}^6$ is a smooth curve of genus 22 and degree 21 such that $\mathcal{O}_D(1)$ is a theta characteristic.

In the case at hand, that is, when $[C, L] \in \mathfrak{Kos}_{\mathfrak{S}_6}$, the curve D must be reducible, for it has C as a component. More precisely:

Lemma 3. *For a general paracanonical curve $C \subseteq \mathbf{P}^6$ having a rank 6 syzygy, the curve D is nodal and consists of two components $C \cup E$, where $E \subseteq \mathbf{P}^6$ is an elliptic septic curve. Furthermore, $\mathcal{O}_D(2) = \omega_D$. The intersection $e := C \cdot E$, viewed as a divisor on C satisfies $e_C \in |\mathcal{O}_C(2) \otimes K_C^\vee|$, and as a divisor on E , satisfies $e_E \in |\mathcal{O}_E(2)|$.*

Remark 4. Note that C is Prym-canonical or canonical if and only if $e_C \in |K_C|$.

The construction above is reversible. Firstly, general element $[C, L] \in \mathfrak{Kos}_{\mathfrak{S}_6}$ can be reconstructed as the residual curve of a reducible spin curve $D \subseteq \mathbf{P}^6$ of genus 22 containing an elliptic curve $E \subseteq \mathbf{P}^6$ with $\deg(E) = 7$ as a component such that the union of D and some point $p \in \mathbf{P}^6 \setminus E$ is the syzygy scheme of a rank 6 linear syzygy among quadrics in \mathbf{P}^6 .

Furthermore, given a reducible spin curve $D = C \cup_e E \subseteq \mathbf{P}^6$ of genus 22 as above, that is, with $\omega_D \cong \mathcal{O}_D(2)$, the genus 8 component C has a nontrivial syzygy of rank 6 involving the quadrics in the 6-dimensional subspace $I_D(2) \subseteq I_C(2)$, see Lemma 29 for a proof of this fact.

2.B. Syzygies and quartics singular along paracanonical curves

We first discuss an alternative characterization of the non-surjectivity of the map $\mu_{C,L}$:

Proposition 5. *Assume the paracanonical curve $\phi_L(C)$ is projectively normal and scheme-theoretically cut out by quadrics. Then $K_{1,2}(C, L) \neq 0$ if and only if there exists a degree 4 homogeneous polynomial on \mathbf{P}^6 , which vanishes to order at least 2 along C but does not belong to the image of the multiplication map $\text{Sym}^2 I_{C,L}(2) \rightarrow I_{C,L}(4)$.*

Proof. We work on the variety $X \xrightarrow{\tau} \mathbf{P}^6$ defined as the blow-up of \mathbf{P}^6 along $\phi_L(C)$. Let E be the exceptional divisor of the blow-up, and consider the line bundle $H := \tau^* \mathcal{O}_{\mathbf{P}^6}(2)(-E)$ on X . Its space of sections identifies to $I_C(2)$, and our assumption that C is scheme-theoretically cut out by quadrics says equivalently that H is a globally generated line bundle on X . The nonvanishing of $K_{1,2}(C, L)$ is equivalent to the non-surjectivity of the multiplication map

$$H^0(X, H) \otimes H^0(X, \tau^* \mathcal{O}(1)) \rightarrow H^0(X, H \otimes \tau^* \mathcal{O}(1)), \quad (2.4)$$

where we use the identification

$$H^0(X, H \otimes \tau^* \mathcal{O}(1)) = H^0(X, \tau^* \mathcal{O}(3)(-E)) = I_C(3).$$

As H is globally generated by its space $W := I_C(2)$ of global sections, the Koszul complex

$$0 \rightarrow \bigwedge^7 W \otimes \mathcal{O}_X(-7H) \rightarrow \dots \rightarrow \bigwedge^2 W \otimes \mathcal{O}_X(-2H) \rightarrow W \otimes \mathcal{O}_X(-H) \rightarrow \mathcal{O}_X \rightarrow 0 \quad (2.5)$$

is exact. We now twist this complex by $\tau^* \mathcal{O}_{\mathbf{P}^6}(1)(H)$ and take global sections. The last map is then the multiplication map (2.4). The successive terms of this twisted complex are

$$\bigwedge^i W \otimes \mathcal{O}_X(\tau^* \mathcal{O}(1))((-i+1)H),$$

for $0 \leq i \leq 7$. The spectral sequence abutting to the hypercohomology of this complex, that is 0, has

$$E_2^{0,0} = \text{coker} \left\{ W \otimes H^0(X, \tau^* \mathcal{O}(1)) \rightarrow H^0(X, H \otimes \tau^* \mathcal{O}(1)) \right\} \quad (2.6)$$

and the terms $E_1^{i,-i-1}$ for $i < -1$ are equal to $\bigwedge^{-i} W \otimes H^{-i-1}(X, \tau^* \mathcal{O}(1)((i+1)H))$. Similarly, we have

$$E_1^{i,-i} = \bigwedge^{-i} W \otimes H^{-i}(X, \tau^* \mathcal{O}(1)((i+1)H)).$$

Lemma 6. (i) *We have*

$$E_1^{i,-i-1} = \bigwedge^{-i} W \otimes H^{-i-1}(X, \tau^* \mathcal{O}(1)((i+1)H)) = 0, \quad (2.7)$$

for $-i-1 = 5, \dots, 1$.

(ii) *For $-i-1 = 6$, that is, $i = -7$, we have*

$$E_1^{-7,6} = \bigwedge^7 W \otimes H^6(X, \tau^* \mathcal{O}(1)(-6H)) = \bigwedge^7 W \otimes I_C(4)_2^\vee, \quad (2.8)$$

where $I_C(4)_2 \subseteq I_C(4)$ is the set of quartic polynomials vanishing at order at least 2 along C , and

$$E_1^{-6,6} = \bigwedge^6 W \otimes H^6(X, \tau^* \mathcal{O}(1)(-5H)) = \bigwedge^6 W \otimes I_C(2)^\vee. \quad (2.9)$$

(iii) *We have $E_1^{i,-i} = 0$, for $-6 < i < 0$.*

Proof of Lemma 6. (i) We want equivalently to show that

$$H^\ell(X, \tau^* \mathcal{O}(1)(-\ell H)) = 0, \quad \text{when } \ell = 5, \dots, 1.$$

Recall that $H = \tau^* \mathcal{O}(2)(-E)$. Furthermore,

$$K_X = \tau^* \mathcal{O}_{\mathbf{P}^6}(-7)(4E). \quad (2.10)$$

So we have to prove that

$$H^\ell(X, \tau^* \mathcal{O}(-2\ell+1)(\ell E)) = 0, \quad \text{for } \ell = 5, \dots, 1. \quad (2.11)$$

Examining the spectral sequence induced by τ , and using the fact that

$$R^s \tau_*(\mathcal{O}_X(tE)) = 0$$

for $s \neq 0, 4$ and also for $s = 4, t \leq 4$, we see that for $1 \leq \ell \leq 4$,

$$H^\ell(X, \tau^* \mathcal{O}(-2\ell + 1)(\ell E)) = H^\ell(\mathbf{P}^6, \mathcal{O}(-2\ell + 1) \otimes R^0 \tau_* \mathcal{O}_X(\ell E)).$$

For $1 \leq \ell \leq 4$, the right hand side is zero, because it is equal to $H^\ell(\mathbf{P}^6, \mathcal{O}(-2\ell + 1))$.

For $\ell = 5$, we have to compute the space $H^5(X, \tau^* \mathcal{O}(-9)(5E))$, which by Serre duality and by (2.10), is dual to the space

$$H^1(X, \tau^* \mathcal{O}(2)(-E)) = H^1(\mathbf{P}^6, \mathcal{O}(2) \otimes \mathcal{I}_C) = 0.$$

(ii) We have to compute the spaces $H^6(X, \tau^* \mathcal{O}(1)(-6H))$ and $H^6(X, \tau^* \mathcal{O}(1)(-5H))$. As $H := \tau^* \mathcal{O}(2)(-E)$, this is rewritten as $H^6(X, \tau^* \mathcal{O}(-11)(6E))$ and $H^6(X, \tau^* \mathcal{O}(-9)(5E))$ respectively. If we dualize using (2.10), we get

$$H^6(X, \tau^* \mathcal{O}(-11)(6E))^\vee = H^0(X, \tau^* \mathcal{O}(4)(-2E)) = I_C(4)_2,$$

$$H^6(X, \tau^* \mathcal{O}(-9)(5E))^\vee = H^0(X, \tau^* \mathcal{O}(2)(-E)) = I_C(2).$$

(iii) We have

$$E_1^{i, -i} = E_1^{-6, 6} = \bigwedge^{-i} W \otimes H^{-i}(X, \tau^* \mathcal{O}(1)((i+1)H)) = \bigwedge^{-i} W \otimes H^{-i}(X, \tau^* \mathcal{O}(2i+3)((-i-1)E)).$$

For $1 \leq -i \leq 5$, we have $R^s \tau_* \mathcal{O}_X((-i-1)E) = 0$ unless $s = 0$. Furthermore, we have

$$R^0 \tau_* \mathcal{O}_X((-i-1)E) = \mathcal{O}_{\mathbf{P}^6},$$

so that

$$H^{-i}(X, \tau^* \mathcal{O}(2i+3)((-i-1)E)) = H^{-i}(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(2i+3)) = 0.$$

□

Corollary 7. *Only one $E_2^{p,q}$ -terms of this spectral sequence is possibly nonzero in degree -1 , namely*

$$E_2^{-7, 6} = \text{Ker} \left\{ \bigwedge^7 W \otimes I_C(4)_2^\vee \rightarrow \bigwedge^6 W \otimes I_C(2)^\vee \right\}. \quad (2.12)$$

Furthermore, all the differentials d_r starting from $E_2^{-7, 6}$ vanish for $2 \leq r < 7$.

Note that the map

$$\bigwedge^7 W \otimes I_C(4)_2^\vee \rightarrow \bigwedge^6 W \otimes I_C(2)^\vee$$

is nothing but the transpose of the multiplication map

$$W \otimes I_C(2) \rightarrow I_C(4)_2,$$

up to trivialization of $\bigwedge^7 W$. It follows that

$$(E_2^{-7, 6})^\vee = \text{Coker} \left\{ W \otimes I_C(2) \rightarrow I_C(4)_2 \right\}. \quad (2.13)$$

Corollary 7 concludes the proof of the proposition since it implies that we have an isomorphism given by d_7 between (2.12) and (2.6), or a perfect duality between (2.12) and the cokernel (2.13). □

Proposition 5 has the following consequence. Recall that P_8^{14} is the moduli space of pairs $[C, L]$, with C being a smooth curve of genus 8 and $L \neq K_C$ a paracanonical line bundle.

Proposition 8. *The Koszul divisor \mathfrak{Kos}_3 of P_8^{14} is the union of two divisors, one of them being the set of pairs $[C, L]$ such that $\phi_L(C)$ is not scheme-theoretically cut out by quadrics, the other being the set of pairs $[C, L]$ such that $H^1(\mathbf{P}^6, \mathcal{I}_C^2(4)) \neq 0$, or equivalently, such that there exists a quartic which is singular along $\phi_L(C)$ but does not lie in $\text{Sym}^2 I_C(2)$.*

Proof. We first have to prove that the locus of pairs $[C, L]$ such that $\phi_L(C)$ is not scheme-theoretically cut-out by quadrics is contained in the divisor \mathfrak{Kos}_3 . This is a consequence of the following lemmas:

Lemma 9. *If $L \neq K_C$ is a projectively normal paracanonical line bundle on a curve of genus 8, then $\phi_L(C)$ is scheme-theoretically cut out by cubics.*

Proof of Lemma 9. We observe that the twisted ideal sheaf $\mathcal{I}_C(3)$ is regular in Castelnuovo-Mumford sense. Indeed, we have

$$H^i(\mathbf{P}^6, \mathcal{I}_C(3-i)) = H^{i-1}(C, L^{\otimes(3-i)})$$

for $i \geq 2$, and the right hand side is obviously 0 for $i-1 \geq 2$, and also 0 for $i-1 = 1$ since $H^1(C, L) = 0$ because $L \neq K_C$ and $\deg L = 2g - 2$. For $i = 1$, we have

$$H^1(\mathbf{P}^6, \mathcal{I}_C(2)) = 0$$

by projective normality. Being regular, the sheaf $\mathcal{I}_C(3)$ is generated by global sections. \square

Corollary 10. *If C, L are as above, and C is not scheme-theoretically cut out by quadrics, then the multiplication map*

$$I_C(2) \otimes H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \rightarrow I_C(3)$$

is not surjective.

To conclude the proof of the proposition, we just have to show that the sublocus of P_8^{14} where L is not projectively normal is not a divisor, since the statement of the proposition will be then an immediate consequence of Proposition 5. We argue along the lines of [12]. First of all, a line bundle L of degree 14 is not generated by sections if and only if $L = K_C(-x + y)$ for some points $x, y \in C$. This determines a codimension 6 locus of P_8^{14} . Similarly L is not very ample if and only if $L = K_C(-x - y + z + t)$, for some points x, y, z, t of C , which is satisfied in a codimension 4 locus of P_8^{14} . Finally, assume L is very ample but $\phi_L(C)$ is not projectively normal. Equivalently

$$\text{Sym}^2 H^0(C, L) \rightarrow H^0(C, L^{\otimes 2})$$

is not surjective, which means that there exists a rank 2 vector bundle F on C which is a nontrivial extension

$$0 \longrightarrow K_C \otimes L^\vee \longrightarrow F \longrightarrow L \longrightarrow 0,$$

such that $h^0(C, F) = 7$. If $x, y, z \in C$, there exists a nonzero section $\sigma \in H^0(C, F)$ vanishing on x, y and z , and thus F is also an extension

$$0 \longrightarrow D \longrightarrow F \longrightarrow K_C \otimes D^\vee \longrightarrow 0, \tag{2.14}$$

where D is a line bundle such that $h^0(C, D(-x - y - z)) \neq 0$, and $h^0(C, L \otimes D^\vee) \neq 0$. We thus have $h^0(C, D) + h^0(C, K_C \otimes D^\vee) \geq 7$ and $\text{Cliff}(D) \leq 2$. As D is effective of degree at least 3, one has the following possibilities:

a) $h^0(C, K_C \otimes D^\vee) = 0$, and then $D = L$, which contradicts the fact that the extension (2.14) is not split;

b) $h^0(C, K_C \otimes D^\vee) = 1$ and $h^0(C, D) \geq 6$, and then $D = L(-x)$ and $h^0(K_C \otimes L^\vee(x)) \neq 0$, so $L = K_C(x - y)$, which happens in a locus of codimension at least 6 in P_8^{14} ;

c) D contributes to the Clifford index of C . As the locus of curves $[C] \in \mathcal{M}_8$ with $\text{Cliff}(C) \leq 2$ is of codimension 2 in \mathcal{M}_8 , this situation does not occur in codimension 1. \square

We shall need later on the following result:

Lemma 11. *Let $\phi_L : C \hookrightarrow \mathbf{P}^6$ be a projectively normal paracanonical curve of genus 8. If C is scheme-theoretically cut out by quadrics, the multiplication map*

$$\mathrm{Sym}^2 I_{C,L}(2) \rightarrow I_{C,L}(4) \quad (2.15)$$

is injective.

Proof. As the restriction map $\phi_L^* : H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(2)) \rightarrow H^0(C, L^{\otimes 2})$ is surjective, its kernel $I_{C,L}(2)$ is of dimension 7. Let as before $\tau : X \rightarrow \mathbf{P}^6$ be the blow-up of \mathbf{P}^6 along $\phi_L(C)$, and let E be its exceptional divisor. We view $I_{C,L}(2)$ as $H^0(X, \tau^*\mathcal{O}(2)(-E))$ and our assumption is that $I_{C,L}(2)$ generates the line bundle $H := \tau^*\mathcal{O}(2)(-E)$ everywhere on X . Thus $I_{C,L}(2)$ provides a morphism

$$\psi : X \rightarrow \mathbf{P}(I_{C,L}(2)). \quad (2.16)$$

Now we have $\deg c_1(H)^6 \neq 0$ by Sublemma 12 below, and thus the morphism ψ has to be generically finite, hence dominant since both spaces have dimension 6. It is then clear that the pull-back map

$$\psi^* : H^0(\mathbf{P}(I_{C,L}(2)), \mathcal{O}(2)) \rightarrow H^0(X, H^{\otimes 2})$$

is injective. On the other hand, this morphism is nothing but the map (2.15). \square

Sublemma 12. *With the same notation as above, we have*

$$\deg c_1(H)^6 = 8. \quad (2.17)$$

Proof. We have

$$c_1(H)^6 = \sum_i \binom{6}{i} (-2)^i h^i \cdot E^{6-i},$$

where $h := \tau^*c_1(\mathcal{O}_{\mathbf{P}^6}(1))$, and

$$h^i \cdot E^{6-i} = 0$$

for $i \neq 6, 1, 0$. Furthermore

$$h^6 = 1, \text{ and } h \cdot E^5 = \deg \phi_L(C) = 14$$

and $E^6 = c_1(N_C)$. By adjunction formula

$$\deg c_1(N_C) = 7\deg \phi_L(C) + \deg K_C = 8 \cdot 14.$$

It follows that

$$\deg c_1(H)^6 = 64 - 6 \cdot 28 + 8 \cdot 14 = 8,$$

which proves (2.17). \square

Proposition 5 and Lemma 11 describe precisely the splitting of the Koszul divisor \mathfrak{Kos}_3 into the divisors \mathfrak{Kos}_{36} and \mathfrak{Kos}_{37} corresponding to paracanonical curves $[C, L] \in P_8^{14}$ having a non-zero syzygy $\gamma \in K_{1,2}(C, L)$ of rank 6 or respectively 7. Precisely, \mathfrak{Kos}_{36} is a unirational divisor (cf. [3] Theorem 5.3) consisting of those paracanonical curves $C \subseteq \mathbf{P}^6$ for which $H^1(\mathbf{P}^6, \mathcal{I}_C^2(4)) \neq 0$. The divisor \mathfrak{Kos}_{37} consists of paracanonical curves $C \subseteq \mathbf{P}^6$ which are not scheme-theoretically cut out by quadrics.

3. First proof: reducible spin curves

3.A. The syzygy is degenerate

The first observation is the following result (already observed experimentally in [3]), which turns out to be useful for the description given below of the general paracanonical curve of genus 8 with nontrivial syzygies.

Proposition 13. *Let $C \subseteq \mathbf{P}^6$ be a smooth paracanonical curve of genus 8 and degree 14, scheme-theoretically generated by quadrics. Then a nontrivial syzygy*

$$\gamma \in \text{Ker} \{I_C(2) \otimes H^0(\mathcal{O}_{\mathbf{P}^6}(1)) \rightarrow I_C(3)\}$$

must be degenerate, that is of rank at most 6.

Proof. We use the morphism

$$\psi : X \rightarrow \mathbf{P}(I_C(2))$$

introduced in (2.16), where $\tau : X \rightarrow \mathbf{P}^6$ is the blow-up of C with exceptional divisor E , and $H := \tau^*\mathcal{O}_{\mathbf{P}^6}(-2E)$. This gives us a morphism

$$(\tau, \psi) : X \rightarrow \mathbf{P}^6 \times \mathbf{P}^6$$

which is of degree 1 on its image, and the syzygy γ induces a hypersurface Y of bidegree $(1, 1)$ in $\mathbf{P}^6 \times \mathbf{P}^6$ containing the 6-dimensional variety $(\tau, \psi)(X)$. Assume to the contrary that γ has maximal rank 7, or equivalently that Y is smooth. Then by the Lefschetz Hyperplane Restriction Theorem, the restriction map $H^{10}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z}) \rightarrow H^{10}(Y, \mathbb{Z})$ is surjective, so that $[(\tau, \psi)(X)]_Y \in H^{10}(Y, \mathbb{Z})$ is the restriction of a class $\beta \in H^{10}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$, which implies that

$$[(\tau, \psi)(X)] = \beta \cdot [Y] \text{ in } H^{12}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z}), \quad (3.18)$$

where $[Y] \in H^2(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$ is the class of Y , that is $h_1 + h_2$, with h_i for $i = 1, 2$ being the pull-backs of the hyperplane classes on each factor. Note that $H^{12}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$ is the set of degree 6 homogeneous polynomials with integral coefficients in h_1 and h_2 . We now have:

Lemma 14. *An element $\alpha \in H^{12}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$ is of the form $(h_1 + h_2) \cdot \beta$ if and only if it satisfies the condition*

$$\sum_{i=0}^6 (-1)^i h_1^i \cdot h_2^{6-i} \cdot \alpha = 0 \text{ in } H^{24}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z}) = \mathbb{Z}. \quad (3.19)$$

Proof of Lemma 14. We have $(h_1 + h_2) \cdot (\sum_i (-1)^i h_1^i \cdot h_2^{6-i}) = 0$ in $H^{14}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$, so one implication is obvious. That the two conditions are equivalent then follows from the fact that both conditions determine a saturated corank 1 sublattice of $H^{12}(\mathbf{P}^6 \times \mathbf{P}^6, \mathbb{Z})$. \square

To conclude that γ has to be degenerate, in view of Lemma 14, it suffices to prove that the class $[(\tau, \psi)(X)]$ does not satisfy (3.19). Since $(\tau, \psi)^*h_1 = c_1(H)$ and $(\tau, \psi)^*h_2 = 2c_1(H) - E$, it is enough to prove that

$$\sum_{i=0}^6 (-1)^i c_1(H)^i \cdot (2c_1(H) - E)^{6-i} \neq 0,$$

which follows from the computations made in the proof of Sublemma 12. \square

3.B. Syzygies and spin curves of genus 22 in \mathbf{P}^6

Recall that $\overline{\mathcal{S}}_g^-$ denotes the moduli stack of odd stable spin curves of genus g , see [5] for details. We start with a nodal genus 22 spin curve of the form $[D := C \cup E, \vartheta] \in \overline{\mathcal{S}}_{22}^-$, where C is a smooth genus 8 curve, E is a smooth elliptic curve and $e := C \cap E$ consists of 14 distinct points, thus $p_a(D) = 22$. Assume $\vartheta \in \text{Pic}^{21}(D)$ verifies $\vartheta^{\otimes 2} \cong \omega_D$, hence the restricted line bundles ϑ_E and ϑ_C have degrees 7 and 14 respectively. Furthermore, $h^0(E, \vartheta_E) = 7$, whereas $h^0(C, \vartheta_C) = 7$ if and only if $\vartheta_C \cong K_C$. The intersection divisor e on the two components of D is characterized by

$$e_C \in |\vartheta_C^{\otimes 2} \otimes K_C^\vee| \quad \text{and} \quad e_E \in |\vartheta_E^{\otimes 2}|.$$

Note in particular that $e_C \in |K_C|$ if and only if $\vartheta_C^{\otimes 2} = K_C^{\otimes 2}$, that is (C, ϑ_C) is canonical or Prym canonical.

The line bundle ϑ on D fits into the Mayer-Vietoris exact sequence:

$$0 \longrightarrow \vartheta \longrightarrow \vartheta_C \oplus \vartheta_E \xrightarrow{r} \mathcal{O}_e(\vartheta) \longrightarrow 0,$$

where r is defined by the isomorphisms on the fibers of ϑ_C and ϑ_E over the points in e . Given $\vartheta_C \in \text{Pic}^{14}(C)$ with $\vartheta_C^{\otimes 2} = K_C(e)$ and $\vartheta_E \in \text{Pic}^7(E)$ with $\vartheta_E^{\otimes 2} = \mathcal{O}_E(e)$, there is a finite number of stable spin curves $[D, \vartheta] \in \overline{\mathcal{S}}_{22}^-$ such that the restrictions of ϑ to C and E are isomorphic to ϑ_C and ϑ_E respectively. Passing to global sections in the Mayer-Vietoris sequence, we obtain the exact sequence:

$$0 \longrightarrow H^0(D, \vartheta) \longrightarrow H^0(C, \vartheta_C) \oplus H^0(E, \vartheta_E) \xrightarrow{r} H^0(\mathcal{O}_e(\vartheta)) \longrightarrow \dots \quad (3.20)$$

Note that r is represented by a 14×14 matrix and $h^0(D, \vartheta) = 14 - \text{rk}(r)$. In the case of a reducible spin curve coming from the syzygy of a paracanonical genus 8 curve in $\mathfrak{R}053_6$, one has $h^0(D, \vartheta) = \text{rk}(r) = 7$.

3.C. Proof of Theorem 1 via reducible spin curves

Theorem 1 states that every Prym canonical curve of genus 8 has a syzygy of rank 6. First we observe the existence of such a curve having the generic behavior described in Lemma 3.

Lemma 15. *There exists a curve $[C, \eta] \in \mathcal{R}_8$, whose Prym canonical model is scheme theoretically cut out by quadrics, and $K_{2,1}(C, K_C \otimes \eta)$ is 1-dimensional, generated by a syzygy γ of rank 6. The syzygy scheme of γ is the union of a point p and a nodal curve $D = C \cup E$, such that E is a smooth elliptic curve of degree 7 and $e := C \cdot E \in |K_C|$ consists of 14 mutually distinct points. Moreover, no cubic polynomial on \mathbf{P}^6 vanishes with multiplicity 2 along C .*

Proof. Examples of singular Prym canonical curves having all these properties have been produced in [3] Proposition 4.4 or [4]. A generic deformation in $\overline{\mathcal{R}}_8$ of these singular examples will provide the required smooth Prym canonical curve. \square

(First) proof of Theorem 1. We denote by X the moduli space of elements $[C, \eta, x_1, \dots, x_{14}]$, where $[C, \eta] \in \mathcal{R}_8$ is a smooth Prym curve of genus 8 and $x_i \in C$ are pairwise distinct points such that $x_1 + \dots + x_{14} \in |K_C| \cong \mathbf{P}^7$. Since the fibres of the forgetful map $X \rightarrow \mathcal{R}_8$ are 7-dimensional, it follows that X is an irreducible variety of dimension 28.

Let T be the locally closed parameter space of odd genus 22 spin curves having the form

$$\left([D := C \cup_{\{x_1, \dots, x_{14}\}} E, \vartheta] : [C] \in \mathcal{M}_8, \sum_{i=1}^{14} x_i \in |K_C|, [E, x_1, \dots, x_{14}] \in \mathcal{M}_{1,14}, \vartheta^{\otimes 2} = \omega_D \right).$$

Observe that points in T , apart from the spin structure $[D, \vartheta] \in \overline{\mathcal{S}}_{22}$ also carry an underlying Prym structure $[C, \eta := K_C \otimes \vartheta_C^\vee] \in \mathcal{R}_8$, for $\vartheta_C^{\otimes 2} \cong K_C(x_1 + \cdots + x_{14}) \cong K_C^{\otimes 2}$. One has an induced finite morphism $T \rightarrow X \times \mathcal{M}_{1,14}$, as well as a map $\mu : T \rightarrow \mathcal{R}_8$ forgetting the 14-pointed elliptic curve. It follows that $\dim T = \dim X + \dim \mathcal{M}_{1,14} = 42$. The locus

$$T_7 := \{[D, \vartheta] \in T : h^0(D, \vartheta) \geq 7\}$$

has the structure of a skew-symmetric degeneracy locus. Applying [13] Theorem 1.10, each component of T_7 has codimension at most $\binom{7}{2} = 21$ inside T , that is, $\dim(T_7) \geq \dim(\mathcal{R}_8)$.

By passing to a general 8-nodal Prym canonical curve $[C, \eta]$, following [3] Proposition 4.5, as well as Lemma 15, we have that $\dim K_{1,2}(C, K_C \otimes \eta) = 1$. In particular, the fibre $\mu^{-1}([C, \eta])$ contains an isolated point, which shows that T_7 is non-empty and has a component which maps dominantly under μ onto \mathcal{R}_8 . Theorem 1 now follows. \square

Remark 16. The same construction can be carried out at the level of general paracanonical curves $[C, L] \in P_8^{14}$, where $L \in \text{Pic}^{14}(C) \setminus \{K_C\}$. The key difference is that we replace T by a variety T' parametrizing objects

$$\left([D := C \cup_{\{x_1, \dots, x_{14}\}} E, \vartheta, L] : [C, x_1, \dots, x_{14}] \in \mathcal{M}_{14,8}, L \in \text{Pic}^{14}(C) \setminus \{K_C\}, \right. \\ \left. \sum_{i=1}^{14} x_i \in |L^{\otimes 2} \otimes K_C^\vee|, [E, x_1, \dots, x_{14}] \in \mathcal{M}_{1,14}, \vartheta^{\otimes 2} = \omega_D \right).$$

Similarly, we have a morphism $\mu' : T' \rightarrow P_8^{14}$ retaining the pair $[C, L]$ alone. The main difference compared to the Prym canonical case is that now

$$\dim |L^{\otimes 2} \otimes K_C^\vee| = 6,$$

therefore $\dim(T') = \dim(P_8^{14}) + \dim(\mathcal{M}_{1,14}) + 6 = 49$. The degeneracy locus $T'_7 \subseteq T'$ defined by the condition $h^0(D, \vartheta) \geq 7$ has codimension 21 inside T' , that is,

$$\dim(T'_7) = 28 = \dim(P_8^{14}) - 1.$$

It follows that the image $\mu'(T'_7) \subseteq P_8^{14}$ has codimension 1, which is in accordance with $\mathfrak{K}053_6$ being a divisor in P_8^{14} .

4. Second proof: Divisor class calculations on $\overline{\mathcal{R}}_g$

Recall [10] that $\overline{\mathcal{R}}_g$ is the Deligne-Mumford moduli space of Prym curves of genus g , whose geometric points are triples $[X, \eta, \beta]$, where X is a quasi-stable curve of genus g , $\eta \in \text{Pic}(X)$ is a line bundle of total degree 0 such that $\eta_E = \mathcal{O}_E(1)$ for each smooth rational component $E \subseteq X$ with $|E \cap \overline{X} - E| = 2$ (such a component is said to be *exceptional*), and $\beta : \eta^{\otimes 2} \rightarrow \mathcal{O}_X$ is a sheaf homomorphism whose restriction to any non-exceptional component is an isomorphism. If $\pi : \overline{\mathcal{R}}_g \rightarrow \overline{\mathcal{M}}_g$ is the map dropping the Prym structure, one has the formula

$$\pi^*(\delta_0) = \delta'_0 + \delta''_0 + 2\delta_0^{\text{ram}} \in CH^1(\overline{\mathcal{R}}_g), \quad (4.21)$$

where $\delta'_0 := [\Delta'_0]$, $\delta''_0 := [\Delta''_0]$, and $\delta_0^{\text{ram}} := [\Delta_0^{\text{ram}}]$ are irreducible boundary divisor classes on $\overline{\mathcal{R}}_g$, which we describe by specifying their respective general points.

We choose a general point $[C_{xy}] \in \Delta_0 \subset \overline{\mathcal{M}}_g$ corresponding to a smooth 2-pointed curve (C, x, y) of genus $g - 1$ and consider the normalization map $\nu : C \rightarrow C_{xy}$, where $\nu(x) = \nu(y)$. A general point

of Δ'_0 (respectively of Δ''_0) corresponds to a pair $[C_{xy}, \eta]$, where $\eta \in \text{Pic}^0(C_{xy})[2]$ and $\nu^*(\eta) \in \text{Pic}^0(C)$ is non-trivial (respectively, $\nu^*(\eta) = \mathcal{O}_C$). A general point of Δ_0^{ram} is a Prym curve of the form (X, η) , where $X := C \cup_{\{x,y\}} \mathbf{P}^1$ is a quasi-stable curve with $p_a(X) = g$ and $\eta \in \text{Pic}^0(X)$ is a line bundle such that $\eta_{\mathbf{P}^1} = \mathcal{O}_{\mathbf{P}^1}(1)$ and $\eta_C^2 = \mathcal{O}_C(-x-y)$. In this case, the choice of the homomorphism β is uniquely determined by X and η . In what follows, we work on the partial compactification $\tilde{\mathcal{R}}_g \subseteq \overline{\mathcal{R}}_g$ of \mathcal{R}_g obtained by removing the boundary components $\pi^{-1}(\Delta_j)$ for $j = 1, \dots, \lfloor \frac{g}{2} \rfloor$, as well as Δ_0'' . In particular, $CH^1(\tilde{\mathcal{R}}_g) = \mathbb{Q}\langle \lambda, \delta'_0, \delta_0^{\text{ram}} \rangle$.

For a stable Prym curve $[X, \eta] \in \tilde{\mathcal{R}}_g$, set $L := \omega_X \otimes \eta \in \text{Pic}^{2g-2}(X)$ to be the paracanonical bundle. For $i \geq 1$, we introduce the vector bundle \mathcal{N}_k over $\tilde{\mathcal{R}}_g$, having fibres

$$\mathcal{N}_k[X, \eta] = H^0(X, L^{\otimes k}).$$

The first Chern class of \mathcal{N}_k is computed in [10] Proposition 1.7:

$$c_1(\mathcal{N}_k) = \binom{k}{2} (12\lambda - \delta'_0 - 2\delta_0^{\text{ram}}) + \lambda - \frac{k^2}{4} \delta_0^{\text{ram}}. \quad (4.22)$$

Then we define the locally free sheaves \mathcal{G}_k on $\tilde{\mathcal{R}}_g$ via the exact sequences

$$0 \longrightarrow \mathcal{G}_k \longrightarrow \text{Sym}^k \mathcal{N}_1 \longrightarrow \mathcal{N}_k \longrightarrow 0,$$

that is, satisfying $\mathcal{G}_k[X, \eta] := I_{X,L}(k) \subseteq \text{Sym}^k H^0(X, L)$. Using (4.22) one computes $c_1(\mathcal{G}_k)$.

We also need the class of the vector bundle \mathcal{G} with fibres

$$\mathcal{G}[X, \eta] = H^0(X, \omega_X^{\otimes 5} \otimes \eta^{\otimes 4}) = H^0(X, \omega_X \otimes L^{\otimes 4}).$$

Lemma 17. *One has $c_1(\mathcal{G}) = 121\lambda - 10\delta'_0 - 24\delta_0^{\text{ram}} \in CH^1(\tilde{\mathcal{R}}_g)$.*

Proof. We apply Grothendieck-Riemann-Roch to the universal Prym curve $f : \mathcal{C} \rightarrow \tilde{\mathcal{R}}_g$. Denote by $\mathcal{L} \in \text{Pic}(\mathcal{C})$ the universal Prym bundle, whose restriction to each Prym curve is the corresponding 2-torsion point, that is, $\mathcal{L}|_{f^{-1}([X, \eta])} = \eta$, for each point $[X, \eta] \in \tilde{\mathcal{R}}_g$. Since $R^1 f_*(\omega_f^{\otimes 5} \otimes \mathcal{L}^{\otimes 4}) = 0$, we write

$$c_1(\mathcal{G}) = f_* \left[\left(1 + 5c_1(\omega_f) + 4c_1(\mathcal{L}) + \frac{(5c_1(\omega_f) + 4c_1(\mathcal{L}))^2}{2} \right) \cdot \left(1 - \frac{c_1(\omega_f)}{2} + \frac{c_1^2(\Omega_f^1) + [\text{Sing}(f)]}{12} \right) \right]_2.$$

We use then the formulas $f_*(c_1^2(\mathcal{L})) = -\delta_0^{\text{ram}}/2$ and $f_*(c_1(\mathcal{L}) \cdot c_1(\omega_f)) = 0$ (see [10], Proposition 1.6) coupled with Mumford's formula $f_*(c_1^2(\Omega_f^1) + [\text{Sing}(f)]) = 12\lambda$ as well with the identity

$$\kappa_1 := f_*(c_1^2(\omega_f)) = 12\lambda - \delta'_0 - 2\delta_0^{\text{ram}},$$

in order to conclude. □

The Koszul locus

$$\mathcal{Z}_8 := \mathfrak{Kos}_3 \cap \mathcal{R}_8 = \left\{ [C, \eta] \in \mathcal{R}_8 : K_{1,2}(C, K_C \otimes \eta) \neq 0 \right\}$$

is a virtual divisor on \mathcal{R}_8 , that is, the degeneracy locus of a map between vector bundles of the same rank over $\tilde{\mathcal{R}}_8$. If it is a genuine divisor (which we aim to rule out), the class of its closure in $\tilde{\mathcal{R}}_8$ is given by [3] Theorem F:

$$[\overline{\mathcal{Z}}_8] = 27\lambda - 4\delta'_0 - 6\delta_0^{\text{ram}} \in CH^1(\tilde{\mathcal{R}}_8).$$

Remark 18. Some of the considerations above can be extended to higher order torsion points. We recall that $\mathcal{R}_{g,\ell}$ is the moduli space of pairs $[C, \eta]$, where C is a smooth curve of genus g and $\eta \in \text{Pic}^0(C)$ is a non-trivial ℓ -torsion point. It is then shown in [3] that the locus $\mathcal{Z}_{8,\ell} := \mathfrak{Kos}_{\mathfrak{J}} \cap \mathcal{R}_{8,\ell} \subseteq P_8^{14}$ is a divisor on $\mathcal{R}_{8,\ell}$ for each other level $\ell \geq 3$. The class of the compactification of $\mathcal{Z}_{8,\ell}$ is given by the following formula, see [3] Theorem F:

$$[\overline{\mathcal{Z}}_{8,\ell}] = 27\lambda - 4\delta'_0 - \sum_{a=1}^{\lfloor \frac{\ell}{2} \rfloor} \frac{4(a^2 - a\ell + \ell^2)}{\ell} \delta_0^{(a)} \in CH^1(\widetilde{\mathcal{R}}_{8,\ell}).$$

We refer to [3] Section 1.4, for the definition of the boundary divisor classes $\delta_0^{(a)}$, where $a = 1, \dots, \lfloor \frac{\ell}{2} \rfloor$. If $\pi : \overline{\mathcal{R}}_{g,\ell} \rightarrow \overline{\mathcal{M}}_g$ is the map forgetting the level ℓ structure, then

$$\pi^*(\delta_0) = \delta'_0 + \delta''_0 + \ell \sum_{\ell=1}^{\lfloor \frac{\ell}{2} \rfloor} \delta_0^{(a)}.$$

We fix now a genus 8 Prym-canonically embedded curve $\phi_L : C \hookrightarrow \mathbf{P}^6$. As usual, we denote the kernel bundle by $M_L := \Omega_{\mathbf{P}^6|C}^1(1)$, hence we have the exact sequence

$$0 \longrightarrow N_C^\vee \otimes L^{\otimes 4} \longrightarrow M_L \otimes L^{\otimes 3} \longrightarrow K_C \otimes L^{\otimes 4} \longrightarrow 0. \quad (4.23)$$

This can be interpreted as an exact sequence of vector bundles over $\widetilde{\mathcal{R}}_8$. Denoting by \mathcal{H} the vector bundle over $\widetilde{\mathcal{R}}_8$ with fibres $H^0(C, N_C^\vee \otimes L^{\otimes 4})$, we compute using the previous formulas and the fact that $\text{rk}(\mathcal{N}_1) = h^0(C, L) = 7$ and $\text{rk}(\mathcal{N}_3) = h^0(C, L^{\otimes 3}) = 35$:

$$c_1(\mathcal{H}) = 35c_1(\mathcal{N}_1) + 7c_1(\mathcal{N}_3) - c_1(\mathcal{N}_4) - c_1(\mathcal{G}) = 100\lambda - 5\delta'_0 - \frac{53}{2}\delta_0^{\text{ram}}. \quad (4.24)$$

Thus $\mathfrak{D}_1 = \mathfrak{Kos}_{\mathfrak{J}_7} \cap \mathcal{R}_8$ and $\mathfrak{D}_2 = \mathfrak{Kos}_{\mathfrak{J}_6} \cap \mathcal{R}_8$. We have already seen in Proposition 5 that $K_{1,2}(C, L) \neq 0$ if and only if either $\phi_L(C) \subseteq \mathbf{P}^6$ is not scheme-theoretically cut out by quadrics, or else, $H^1(\mathbf{P}^6, \mathcal{I}_C^2(4)) \neq 0$. We write

$$\mathcal{Z}_8 = \mathfrak{D}_1 + \mathfrak{D}_2, \quad \text{where}$$

$$\mathfrak{D}_1 := \left\{ [C, \eta] \in \mathcal{R}_8 : \phi_L(C) \subseteq \mathbf{P}^6 \text{ is scheme-theoretically not cut out by quadrics} \right\}$$

and

$$\mathfrak{D}_2 := \left\{ [C, \eta] \in \mathcal{R}_8 : H^1(\mathbf{P}^6, \mathcal{I}_C^2(4)) \neq 0 \right\}.$$

We have already observed that $\dim I_{C,L}(2) = 7$ and $\chi(\mathbf{P}^6, \mathcal{I}_C^2(4)) = 28$. If \mathcal{Z}_8 is a divisor, then \mathfrak{D}_2 is a divisor as well and for $[C, \eta] \in \mathcal{R}_8 \setminus \mathfrak{D}_2$, we have that

$$\dim \text{Sym}^2 I_{C,L}(2) = \dim I_{C,L}(4)_2 = 28.$$

Paying some attention to its definition, the divisor \mathfrak{D}_1 can be thought as the degeneracy locus

$$\left\{ [C, \eta] \in \mathcal{R}_8 : \text{Sym}^2 I_{C,L}(2) \xrightarrow{\neq} I_{C,L}(4)_2 \right\},$$

which is an effective divisor on $\widetilde{\mathcal{R}}_8$. We compute the class of this divisor:

Theorem 19. *We have the following formulas:*

$$[\overline{\mathfrak{D}}_1] = 7\lambda - \frac{1}{2}\delta'_0 - \frac{3}{4}\delta_0^{\text{ram}} \in CH^1(\widetilde{\mathcal{R}}_8)$$

and

$$[\overline{\mathfrak{D}}_2] = 20\lambda - \frac{7}{2}\delta'_0 - \frac{21}{4}\delta_0^{\text{ram}} \in CH^1(\widetilde{\mathcal{R}}_8).$$

Proof. We first globalize over $\widetilde{\mathcal{R}}_8$ the following exact sequence:

$$0 \longrightarrow I_{C,L}(4)_2 \longrightarrow I_{C,L}(4) \longrightarrow H^0(C, N_C^\vee \otimes L^{\otimes 4}) \longrightarrow H^1(\mathbf{P}^6, \mathcal{I}_C^2(4)) \longrightarrow 0.$$

Denote by \mathcal{A} the sheaf on $\widetilde{\mathcal{R}}_8$ supported along the divisor \mathcal{D}_2 , whose fibre over a general point of that divisor is equal to $H^1(\mathbf{P}^6, \mathcal{I}_C^2(4))$. There is a surjective morphism of sheaves

$$\mathcal{H} \rightarrow \mathcal{A}$$

and denote by \mathcal{G}'_4 its kernel. Since \mathcal{A} is locally free along \mathcal{D}_2 and $\widetilde{\mathcal{R}}_8$ is a smooth stack, using the Auslander-Buchsbaum formula we find that \mathcal{G}'_4 is a locally free sheaf of rank equal to $\text{rk}(\mathcal{H}) = \chi(C, N_C^\vee(4L)) = 19 \cdot 7$. Precisely, \mathcal{G}'_4 is an elementary transformation of \mathcal{H} along the divisor \mathcal{D}_2 . Furthermore, $c_1(\mathcal{G}'_4) = c_1(\mathcal{H}) - [\overline{\mathcal{D}}_2]$.

The morphism $\mathcal{G}_4 \rightarrow \mathcal{H}$ globalizing the maps $I_{C,L}(4) \rightarrow H^0(C, N_C^\vee \otimes L^{\otimes 4})$ factors through the subsheaf \mathcal{G}'_4 and we form the exact sequence:

$$0 \longrightarrow \mathcal{G}_4^2 \longrightarrow \mathcal{G}_4 \longrightarrow \mathcal{G}'_4 \longrightarrow 0.$$

The multiplication maps $\text{Sym}^2 I_{C,L}(2) \rightarrow I_{C,L}(4)_2$ globalize to a sheaf morphism

$$\nu : \text{Sym}^2(\mathcal{G}_2) \rightarrow \mathcal{G}_4^2$$

between locally free sheaves of the same rank 28 over the stack $\widetilde{\mathcal{R}}_8$. The degeneration locus of ν is precisely the divisor $\overline{\mathcal{D}}_1$. We compute:

$$c_1(\text{Sym}^2(\mathcal{G}_2)) = 8c_1(\mathcal{G}_2) = 8(8c_1(\mathcal{N}_1) - c_1(\mathcal{N}_2)) = -40\lambda + 8(\delta'_0 + \delta_0^{\text{ram}}),$$

and

$$c_1(\mathcal{G}_4^2) = 120c_1(\mathcal{N}_1) - c_1(\mathcal{N}_4) - c_1(\mathcal{H}) + [\overline{\mathcal{D}}_2] = -53\lambda + 11\delta'_0 + \frac{25}{2}\delta_0^{\text{ram}} + [\overline{\mathcal{D}}_2].$$

We obtain the relation $[\overline{\mathcal{D}}_1] - [\overline{\mathcal{D}}_2] = -13\lambda + 3\delta'_0 + \frac{9}{2}\delta_0^{\text{ram}}$. Since at the same time

$$[\overline{\mathcal{D}}_1] + [\overline{\mathcal{D}}_2] = [\mathcal{Z}_8] = 27\lambda - 4\delta'_0 - 6\delta_0^{\text{ram}},$$

we solve the system and conclude. \square

We are now in a position to give a second proof of Theorem 1:

Theorem 20. *The class $[\overline{\mathcal{D}}_2]$ cannot be effective. It follows that $\mathcal{Z}_8 = \mathcal{R}_8$ and $K_{1,2}(C, K_C \otimes \eta) \neq 0$, for every Prym curve $[C, \eta] \in \mathcal{R}_8$.*

Proof. We use the sweeping curve of the boundary divisor Δ'_0 of $\widetilde{\mathcal{R}}_8$ constructed via Nikulin surfaces in [11] Lemma 3.2: Precisely, through the general point of Δ'_0 there passes a rational curve $\Gamma \subseteq \Delta'_0$, entirely contained in $\widetilde{\mathcal{R}}_8$, having the following numerical characters:

$$\Gamma \cdot \lambda = 8, \quad \Gamma \cdot \delta'_0 = 42, \quad \text{and} \quad \Gamma \cdot \delta_0^{\text{ram}} = 8.$$

We note that $\Gamma \cdot \overline{\mathcal{D}}_2 < 0$. Writing $\overline{\mathcal{D}}_2 \equiv \alpha \cdot \delta'_0 + E$, where $\alpha \geq 0$ and E is an effective divisor whose support is disjoint from Δ'_0 , we immediately obtain a contradiction. \square

The divisors \mathfrak{D}_1 and \mathfrak{D}_2 can be defined in an identical manner at the level of each moduli space $\overline{\mathcal{R}}_{8,\ell}$ of twisted level ℓ curves of genus g . As already pointed out, in the case $\ell \geq 3$ it follows from [3] Proposition 4.4 that both \mathfrak{D}_1 and \mathfrak{D}_2 are actual divisors. Repeating the same calculations as for $\ell = 2$, we obtain the following formula on the partial compactification $\widetilde{\mathcal{R}}_{8,\ell}$ of $\mathcal{R}_{8,\ell}$:

$$[\overline{\mathfrak{D}}_2] = 20\lambda - \frac{7}{2}\delta'_0 - \sum_{a=1}^{\lfloor \frac{\ell}{2} \rfloor} \frac{1}{2\ell} (7a^2 - 7a\ell + 17\ell^2 - 20\ell)\delta_0^{(a)} \in CH^1(\widetilde{\mathcal{R}}_{8,\ell}). \quad (4.25)$$

As an application, we mention a different proof of one of the main results from [1]:

Theorem 21. *The canonical class of $\overline{\mathcal{R}}_{8,\ell}$ is big for $\ell \geq 3$. It follows that $\overline{\mathcal{R}}_{8,\ell}$ is a variety of general type for $\ell = 3, 4, 6$.*

Proof. Using formula (4.25), it is a routine exercise to check that for $\ell \geq 3$ the canonical class computed in [3] Proposition 1.5

$$K_{\widetilde{\mathcal{R}}_{8,\ell}} = 13\lambda - 2\delta'_0 - (\ell + 1) \sum_{a=1}^{\lfloor \frac{\ell}{2} \rfloor} \delta_0^{(a)}$$

can be written as a *positive* combination of the big class λ and the effective class $[\overline{\mathfrak{D}}_2]$, hence it is big. Arguing along the lines of [3] Remark 3.5, it is easy to extend this result to the full compactification $\overline{\mathcal{R}}_{8,\ell}$ and deduce that $K_{\overline{\mathcal{R}}_{8,\ell}}$ is big.

To conclude that $\overline{\mathcal{R}}_{8,\ell}$ is of general type, one needs, apart from the bigness of the canonical class $K_{\widetilde{\mathcal{R}}_{8,\ell}}$ of the moduli stack, a result that the singularities of the coarse moduli space $\overline{\mathcal{R}}_{8,\ell}$ impose no adjunction conditions. This is only known for $2 \leq \ell \leq 6, \ell \neq 5$, see [2]. \square

5. Rank 2 vector bundles and singular quartics

Our goal in this section is to propose a construction of syzygies of Prym canonical curves of genus 8. We also sketch the proof of the fact that these syzygies are nontrivial. We fix again a general element $[C, \eta] \in \mathcal{R}_8$ and set $L := K_C \otimes \eta$. According to Proposition 5, in order to prove that $K_{2,1}(C, L) \neq 0$, we have to produce quartic hypersurfaces in \mathbf{P}^6 which vanish at order at least 2 along $\phi_L(C)$, but do not lie in the image of the map $\text{Sym}^2 I_{C,L}(2) \rightarrow I_{C,L}(4)$. The goal of this section is to produce such quartics from rank 2 vector bundles on C . The (incomplete) proof that the quartics we construct are not in the image of $\text{Sym}^2 I_{C,L}(2)$ depends on an unproved general position statement (*), but there might be other approaches exploiting the fact that the hypersurfaces in question are determinantal.

The following construction produces quartics vanishing at order 2 along C . Let E be a rank 2 vector bundle on C , with determinant K_C . Assume

$$h^0(C, E) = 4, \quad h^0(C, E(\eta)) = 4. \quad (5.26)$$

Setting $V_0 := H^0(C, E)$ and $V_1 := H^0(C, E(\eta))$, we have a natural map

$$V_0 \otimes V_1 \rightarrow H^0(C, L),$$

defined using evaluation and the following composite map:

$$H^0(E) \otimes H^0(E(\eta)) \rightarrow H^0(E \otimes E(\eta)) \cong H^0(\mathcal{E}nd E \otimes L) \xrightarrow{\text{Tr}} H^0(C, L). \quad (5.27)$$

This map gives dually a morphism

$$H^0(C, L)^\vee \rightarrow V_0^\vee \otimes V_1^\vee,$$

(which will be proved below to be injective for a general choice of E). We consider the quartic hypersurface D_4 on $\mathbf{P}(V_0^\vee \otimes V_1^\vee)$ parametrizing tensors of rank at most 3.

Lemma 22. *The restriction $D_{4,E}$ of this quartic to $\mathbf{P}(H^0(C, L)^\vee) \subseteq \mathbf{P}(V_0^\vee \otimes V_1^\vee)$ is singular along the curve C .*

Proof. The quartic D_4 is singular along the set $T_2 \subseteq \mathbf{P}(V_0^\vee \otimes V_1^\vee)$ of tensors of rank at most 2. The quartic $D_{4,E}$ in $\mathbf{P}(H^0(C, L)^\vee)$ is thus singular along $T_2 \cap \mathbf{P}(H^0(C, L)^\vee)$, which obviously contains $C \subseteq \mathbf{P}(H^0(C, L)^\vee)$, since at a point $p \in C$, the map $V_0 \otimes V_1 \rightarrow H^0(C, L)$ composed with the evaluation at p factors through $E|_p \otimes E(\eta)|_p$. \square

By Brill-Noether theory, the variety $W_7^1(C)$ of degree 7 pencils on C is 4-dimensional. There should thus exist finitely many elements $D \in W_7^1(C)$ with the property that

$$h^0(C, D) \geq 2, h^0(C, D \otimes \eta) \geq 2. \quad (5.28)$$

We now have the following lemma:

Lemma 23. *Let $[C, \eta] \in \mathcal{R}_8$ be as above and $D \in W_7^1(C)$ satisfying (5.28). Then*

(i) $h^0(C, D) = 2$ and $h^0(C, D \otimes \eta) = 2$. *The multiplication map*

$$\left(H^0(C, D) \otimes H^0(C, K_C \otimes D^\vee) \right) \oplus \left(H^0(C, D \otimes \eta) \otimes H^0(C, K_C \otimes D^\vee \otimes \eta) \right) \rightarrow H^0(C, K_C)$$

is surjective (in fact, an isomorphism).

(ii) *The multiplication map*

$$\left(H^0(C, D) \otimes H^0(C, K_C \otimes D^\vee \otimes \eta) \right) \oplus \left(H^0(C, D \otimes \eta) \otimes H^0(C, K_C \otimes D^\vee) \right) \rightarrow H^0(C, K_C(\eta))$$

is surjective.

Proof. This can be proved by a degeneration argument, for example by degenerating C to the union of two curves of genus 4 meeting at one point. \square

By Brill-Noether theory, the following corollary follows from (i) above:

Corollary 24. *For $[C, \eta]$ as above, the set of pencils $D \in W_7^1(C)$ satisfying (5.28) is finite.*

Given such a D , we form the rank 2 vector bundle

$$E = D \oplus (K_C \otimes D^\vee)$$

on C which satisfies the conditions (5.26). The associated quartic is however not interesting for our purpose, due to the following fact:

Lemma 25. *The quartic on $\mathbf{P}(H^0(C, L)^\vee)$ associated to the vector bundle $D \oplus (K_C \otimes D^\vee)$ is the union of the two quadrics Q_0 and Q_1 associated respectively with the multiplication maps*

$$H^0(D) \otimes H^0((K_C \otimes D^\vee)(\eta)) \rightarrow H^0(K_C(\eta)) \text{ and } H^0(D(\eta)) \otimes H^0(K_C \otimes D^\vee) \rightarrow H^0(K_C(\eta)).$$

Both these quadrics contain C .

Proof. Indeed we have in this case

$$V_0 = H^0(C, E) = H^0(C, D) \oplus H^0(C, K_C \otimes D^\vee), \text{ respectively}$$

$$V_1 = H^0(C, E(\eta)) = H^0(C, D \otimes \eta) \oplus H^0(C, K_C \otimes D^\vee \otimes \eta).$$

Furthermore, it is clear that the map of (5.27) factors through the projection

$$V_0 \otimes V_1 \rightarrow \left(H^0(C, D) \otimes H^0(C, K_C \otimes D^\vee \otimes \eta) \right) \oplus \left(H^0(C, K_C \otimes D^\vee) \otimes H^0(C, D \otimes \eta) \right)$$

and induces on each summand the multiplication map. The quadric Q_0 is by definition associated with the the multiplication map

$$\mu_0 : H^0(C, D) \otimes H^0(C, K_C \otimes D^\vee \otimes \eta) \rightarrow H^0(C, K_C \otimes \eta),$$

and is the set of elements f in $\mathbf{P}(H^0(K_C \otimes \eta))^\vee$ such that $\mu_0^*(f)$ is a tensor of rank ≤ 1 . Similarly for Q_1 , with D being replaced with $D(\eta)$. Finally we use the fact that a tensor

$$(\mu_0^*f, \mu_1^*f) \in \left(H^0(C, D) \otimes H^0(C, K_C \otimes D^\vee \otimes \eta) \right) \oplus \left(H^0(C, K_C \otimes D^\vee) \otimes H^0(C, D \otimes \eta) \right)$$

has rank at most 3 if and only if one of μ_0^*f and μ_1^*f has rank at most 1. \square

We recall from [15] or [14] that the Brill-Noether condition $h^0(C, E) \geq 4$ imposes only $10 = \binom{5}{2}$ equations on the parameter space of rank 2 vector bundles E with determinant K_C . As $\det E(\eta) \cong K_C$ as well, we conclude that the equations (5.26) impose only 20 conditions. As the moduli space $SU_C(2, K_C)$ of semistable rank 2 vector bundles on C having determinant K_C has dimension $3g - 3 = 21$, in our case we conclude that there is a positive dimensional family of such vector bundles on C satisfying (5.26).

We now sketch the proof of the fact that for C general of genus 8 and $D \in W_7^1(C)$ satisfying (5.28), for a general deformation E of the vector bundle $D \oplus (K_C \otimes D^\vee)$ satisfying $\det E \cong K_C$ and $h^0(C, E) = 4$, the associated quartic $D_{4,E}$ singular along C is not defined by an element of $\text{Sym}^2 I_C(2)$. Combined with Proposition 5, this provides a third approach to Theorem 1. The proof of this fact rests on an unproven general position statement (*), so it is incomplete.

Sketch of proof of the nontriviality of the syzygy. The vector bundle E is generated by sections, as it is a general section-preserving deformation of the vector bundle

$$D \oplus (K_C \otimes D^\vee)$$

which is generated by global sections, and similarly for $E(\eta)$. Along $C \subseteq \mathbf{P}(H^0(C, L)^\vee)$, then the rational map

$$\mathbf{P}(H^0(C, L)^\vee) \dashrightarrow \mathbf{P}(H^0(E)^\vee \otimes H^0(E(\eta))^\vee)$$

is well-defined and the image of C is contained in the locus $T_{2,E}$ of tensors of rank exactly 2. In fact, the case of $D \oplus (K_C \otimes D^\vee)$ shows that this map is a morphism for general E (one just needs to know that $H^0(C, K_C \otimes \eta)$ is generated by the two vector spaces $H^0(D) \otimes H^0(K_C \otimes D^\vee \otimes \eta)$ and $H^0(D \otimes \eta) \otimes H^0(K_C \otimes D^\vee)$ respectively, or rather their images under the multiplication map. Note that on $T_{2,E}$, there is a rank 2 vector bundle M which restricts to E on C .

In the case of the split vector bundle $E_{\text{sp}} = D \oplus (K_C \otimes D^\vee)$, Lemma 25 shows that the Zariski closure $\overline{T_{2,E_{\text{sp}}}}$ parameterizing tensors of rank ≤ 2 in $\mathbf{P}(H^0(C, L)^\vee) \subseteq \mathbf{P}(V_0^\vee \otimes V_1^\vee)$ is equal to the singular locus of $D_{4,E_{\text{sp}}}$ and consists of the union of the two planes P_0, P_1 defined as the singular loci of the quadrics Q_0, Q_1 respectively, and the intersection $Q_0 \cap Q_1$. The locus $\overline{T_{2,E_{\text{sp}}}} \setminus T_{2,E_{\text{sp}}}$ is the locus where the tensor has rank 1, and this happens exactly along the two conics $P_0 \cap Q_1$ and $P_1 \cap Q_0$. The curve C is contained in $Q_0 \cap Q_1$ and does not intersect $P_0 \cup P_1$. In particular, the rational map $\phi : \widetilde{\mathbf{P}^6} \dashrightarrow \mathbf{P}^6$ given by the linear system $I_C(2)$ is well defined along $P_0 \cup P_1$. We believe that the following general position statement concerning the two planes P_i is true for general C and D, η as above.

(*) The surfaces $\phi(P_i)$ are projectively normal Veronese surfaces, generating a hyperplane $\langle \phi(P_i) \rangle \subseteq \mathbf{P}^6$. Furthermore, the surface $\phi(P_0) \cup \phi(P_1) \subseteq \mathbf{P}^6$ is contained in a unique quadric in \mathbf{P}^6 , namely the union of the two hyperplanes $\langle \phi(P_0) \rangle$ and $\langle \phi(P_1) \rangle$.

We now prove that, assuming (*), for a general vector bundle E as above, the associated quartic $D_{4,E}$ singular along C is not defined by an element of $\text{Sym}^2 I_C(2)$. As P_0, P_1 are 2-dimensional reduced components of $\overline{T_{2,E_{\text{sp}}}}$, hence of the right dimension, the theory of determinantal hypersurfaces shows that for general E as above, there is a reduced surface $\Sigma_E \subseteq \overline{T_{2,E}}$ whose specialization when $E = E_{\text{sp}}$ contains $P_0 \cup P_1$. Let $\mathcal{E} \rightarrow C \times B$ be a family of vector bundles on C parameterized by a smooth curve B , with general fiber E and special fiber E_{sp} . Denote by \mathcal{E}_b the restriction of \mathcal{E} to $C \times \{b\}$. Property (*) then implies that $\phi(\Sigma_{\mathcal{E}_b})$ for general $b \in B$ is contained in at most one quadric $Q_{\mathcal{E}_b}$ in \mathbf{P}^6 . We argue by contradiction and assume that the quartic D_{4,\mathcal{E}_b} is a pull-back $\phi^{-1}(Q)$ for general b . One thus must have $Q = Q_{\mathcal{E}_b}$. Next, the determinantal quartic D_{4,\mathcal{E}_b} is singular along T_{2,\mathcal{E}_b} , hence along $\Sigma_{\mathcal{E}_b}$. Let $b \mapsto q_{\mathcal{E}_b} \in \text{Sym}^2 I_C(2)$, where $q_{\mathcal{E}_b}$ is a defining equation for the quadric $Q_{\mathcal{E}_b}$. Then we find that the first order derivative of the family $\phi^* q_{\mathcal{E}_b}$ at b_0 also vanishes along $\Sigma_{\mathcal{E}_{b_0}}$, hence it must be proportional to $\phi^* q_{\mathcal{E}_{b_0}}$. We then conclude that the quadric $Q_{\mathcal{E}_b}$ is in fact constant, and thus must be equal to the quadric $Q_{E_{\text{sp}}}$. We now reach a contradiction by proving the following lemma. \square

Lemma 26. *If the determinantal quartic D_{4,\mathcal{E}_b} is constant, equal to $D_{\text{sp}} = Q_0 \cup Q_1$, then the vector bundle \mathcal{E}_b on C does not deform with $b \in B$.*

Proof. Denoting $V_{0,b} := H^0(C, \mathcal{E}_b)$, $V_{1,b} := H^0(C, \mathcal{E}_b(\eta))$, we have the multiplication map

$$V_{0,b} \otimes V_{1,b} \rightarrow H^0(C, K_C \otimes \eta)$$

which is surjective for generic b since it is surjective for $\mathcal{E}_0 = D \oplus (K_C \otimes D^\vee)$ (see Lemma 23). The determinantal quartic D_{4,\mathcal{E}_b} is the vanishing locus of the determinant of the corresponding bundle map

$$\sigma_b : V_{0,b} \otimes \mathcal{O}_{\mathbf{P}(H^0(C, K_C(\eta))^\vee)} \rightarrow V_{1,b}^\vee \otimes \mathcal{O}_{\mathbf{P}(H^0(C, K_C(\eta))^\vee)}(1) \quad (5.29)$$

on $\mathbf{P}(H^0(C, K_C \otimes \eta)^\vee)$. We know that $D_{4,\mathcal{E}_b} = Q_0 \cup Q_1$ for any $b \in B$, where the quadrics Q_i are singular (of rank 4), but with singular locus P_i not intersecting $C \subseteq Q_0 \cap Q_1$. The morphism σ_b has rank exactly 1 generically along each Q_i and the kernel of $\sigma|_{D_{4,b}}$ determines a line bundle $\mathcal{K}_{i,b}$ on its smooth locus $Q_i \setminus P_i$. This line bundle is independent of b since $\text{Pic}(Q_i \setminus P_i)$ has no continuous part. The restriction of $\mathcal{K}_{i,b}$ to C is thus constant. Finally, on the smooth part of $(Q_0 \cap Q_1)_{\text{reg}}$, the kernel $\text{Ker}(\sigma)$ contains the two line bundles $\mathcal{K}_{i,b}|_{Q_0 \cap Q_1}$. Restricting to $C \subseteq (Q_0 \cap Q_1)_{\text{reg}}$, we conclude that $\text{Ker} \sigma_b|_C$ contains $\mathcal{K}_{i,0}|_C$ for $i = 0, 1$. For $b = 0$, one has

$$\text{Ker} \sigma_0|_C = \mathcal{K}_{0,0}|_C \oplus \mathcal{K}_{1,0}|_C$$

and this thus remains true for general b . Finally, it follows from the construction and the fact that \mathcal{E}_b is generated by its sections that $\text{Ker} \sigma_b|_C = \mathcal{E}_b^\vee$, which finishes the proof. \square

6. Miscellany

6.A. Extra remarks on the geometry of paracanonical curves of genus 8 with a nontrivial syzygy

We now comment on an interesting rank 2 vector bundle appearing in our situation. Again, let $\phi_L : C \hookrightarrow \mathbf{P}^6$ be a paracanonical curve of genus 8. We assume L is scheme-theoretically cut out by quadrics. Denoting by N_C the normal bundle of C in the embedding in \mathbf{P}^6 , we consider the natural

map $I_C(2) \otimes \mathcal{O}_C \rightarrow N_C^\vee \otimes L^{\otimes 2}$ (which is surjective by our assumption) given by differentiation along $\phi_L(C)$, and let F denote its kernel. We thus have the short exact sequence:

$$0 \longrightarrow F \longrightarrow I_C(2) \otimes \mathcal{O}_C \longrightarrow N_C^\vee \otimes L^{\otimes 2} \longrightarrow 0. \quad (6.30)$$

If $K_{1,2}(C, L) \neq 0$, the map $\mu : I_C(2) \otimes H^0(\mathbf{P}^6, \mathcal{O}(1)) \rightarrow I_C(3)$ is not surjective, hence not injective. A fortiori, the map

$$\bar{\mu} : I_C(2) \otimes H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \rightarrow H^0(C, N_C^\vee \otimes L^{\otimes 3})$$

induced by (6.30) is not injective, so that $h^0(C, F(L)) \neq 0$. In fact, the equivalence between the statements $h^0(C, F(L)) \neq 0$ and $K_{1,2}(C, L) \neq 0$ follows from the same argument once we know that there is no cubic polynomial on \mathbf{P}^6 vanishing with multiplicity 2 along C .

We observe now that F is a vector bundle of rank 2 on the curve C , with determinant equal to $\det N_C \otimes L^{\otimes (-2)} \cong K_C \otimes L^{\otimes (-3)}$. Hence if $F(L)$ has a nonzero section, assuming this section vanishes nowhere along C , then $F(L)$ is an extension of $K_C \otimes L^\vee$ by \mathcal{O}_C . This provides an extension class

$$e \in H^1(C, L \otimes K_C^\vee) = H^0(C, K_C^{\otimes 2} \otimes L^\vee)^\vee. \quad (6.31)$$

Assume now $L \otimes K_C^\vee =: \eta$ is a nonzero 2-torsion element of $\text{Pic}^0(C)$. Then

$$e \in H^0(C, L)^\vee.$$

On the other hand, according to Theorem 20, there exists a nontrivial syzygy

$$\gamma = \sum_{i=1}^6 \ell_i \otimes q_i \in K_{1,2}(C, L) = \text{Ker}\{H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1)) \otimes I_C(2) \rightarrow I_C(3)\},$$

which is degenerate by Proposition 13. As we saw already, it has in fact rank 6 for generic $[C, \eta]$, hence determines a nonzero element

$$f \in H^0(\mathbf{P}^6, \mathcal{O}_{\mathbf{P}^6}(1))^\vee = H^0(C, L)^\vee = H^1(C, K_C \otimes L^\vee) = H^1(C, L \otimes K_C^\vee), \quad (6.32)$$

which is well-defined up to a coefficient.

Proposition 27. *The two elements e and f are proportional.*

Proof. Equivalently, we show that the kernels of the two linear forms $e, f \in H^0(C, L)^\vee$ are equal. Viewing γ as an element of $\text{Hom}(I_C(2)^\vee, H^0(C, L))$, we have $\text{Ker}(f) = \text{Im}(\gamma)$. On the other hand, the kernel of e identifies with

$$\text{Im}\left\{j : H^0(C, F \otimes L^{\otimes 3} \otimes K_C^\vee) \rightarrow H^0(C, L)\right\},$$

where the map j is obtained by twisting the exact sequence $0 \rightarrow \mathcal{O}_C \rightarrow F(L) \rightarrow K_C \otimes L^\vee \rightarrow 0$ by K_C . We have $F \otimes L^{\otimes 3} \otimes K_C^\vee \cong F^\vee$ since $\det F \cong K_C \otimes L^{\otimes (-3)}$, hence there is a natural morphism

$$i^* : I_C(2)^\vee \otimes \mathcal{O}_C \rightarrow F^\vee \cong F(L^{\otimes 3} \otimes K_C^\vee)$$

dual to the inclusion $F \hookrightarrow I_C(2) \otimes \mathcal{O}_C$ of (6.30). The proposition follows from the following claim:

Claim. *The morphism $\alpha : I_C(2)^\vee \rightarrow H^0(C, L)$ is equal to $j \circ i^*$.*

Forgetting about the last identification $F^\vee \cong F \otimes L^{\otimes 3} \otimes K_C^\vee$, the claim amounts to the following general fact: For an evaluation exact sequence on a variety X

$$0 \longrightarrow G \longrightarrow W \otimes \mathcal{O}_X \longrightarrow M \longrightarrow 0$$

and for a section $s \in H^0(X, G(L)) = H^0(X, \text{Hom}(G^\vee, L))$ giving an element

$$s' \in \text{Ker}\left\{W \otimes H^0(X, L) \rightarrow H^0(X, M \otimes L)\right\} \subseteq \text{Hom}(W^\vee, H^0(X, L)),$$

the induced map $s : H^0(X, G^\vee) \rightarrow H^0(X, L)$ composed with the map $W^\vee \rightarrow H^0(X, G^\vee)$ equals the map $s' : W^\vee \rightarrow H^0(X, L)$. \square

6.B. Further properties

Using the exact sequence (6.30) in the general case of a genus 8 paracanonical curve $[C, L] \in P_8^{14}$, we obtain:

Lemma 28. *A section $s \in H^0(C, F(L)) \subseteq I_{C,L}(2) \otimes H^0(C, L) = \text{Hom}(I_{C,L}(2)^\vee, H^0(C, L))$ of rank 6, determines an element $e \in |2L - K_C|$.*

Proof. The multiplication by $s \in H^0(F(L)) \subseteq I_{C,L}(2) \otimes H^0(C, L) = H^0(I_{C,L}(2)^\vee \otimes L)$ determines the natural maps $F^\vee \rightarrow L$ and $g_s : I_C(2)^\vee \otimes \mathcal{O}_C \rightarrow L$ sitting in the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(g_s) & \longrightarrow & I_C(2)^\vee \otimes \mathcal{O}_C & \longrightarrow & L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & 2L - K_C & \longrightarrow & F^\vee & \longrightarrow & L \longrightarrow 0 \end{array},$$

where $I_C(2)^\vee \otimes \mathcal{O}_C \rightarrow F^\vee$ is the dual of the natural inclusion of (6.30). Passing to global sections we get the inclusion $H^0(\text{Ker}(g_s)) = \text{Ker}\{I_{C,L}(2)^\vee \rightarrow H^0(C, L)\} \hookrightarrow H^0(2L - K_C)$, which by hypothesis is 1-dimensional hence it defines an element $e \in |2L - K_C|$. \square

Via the exact sequence (6.30) we can also show directly the following result that has been used in Section 3:

Lemma 29. *If there is a spin curve $D = C \cup E \hookrightarrow \mathbf{P}^6$ of genus 22 and degree 21 containing the genus 8 paracanonical curve $[C, L]$ as in Lemma 3, then $H^0(C, F(L)) \neq 0$. If there is no cubic polynomial on \mathbf{P}^6 vanishing with multiplicity 2 along C , then $K_{1,2}(C, L) \neq 0$.*

Proof. Let $e = C \cap E$ and recall $c_1(F) = -3L + K_C$ and $\mathcal{O}_C(e) = 2L - K_C$. Note that $I_D(2) \subseteq I_C(2)$ is 6-dimensional. Tensor then the first vertical exact sequence of the following diagram by L and pass to global sections.

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & L^\vee & \longrightarrow & I_D(2) \otimes \mathcal{O}_C & \longrightarrow & \mathcal{I}_D/(\mathcal{I}_D \cap \mathcal{I}_C^2)(2) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F & \longrightarrow & I_C(2) \otimes \mathcal{O}_C & \longrightarrow & N_C^\vee(2) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{O}_C(-e) & \longrightarrow & \mathcal{O}_C & \longrightarrow & \mathcal{O}_{C|e} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

\square

6.C. Nontrivial syzygies of paracanonical curves via vector bundles

We return to the proof of Theorem 20 given in Section 5. Consider now a general paracanonical curve $[C, K_C \otimes \eta] \in P_8^{14}$. For a rank 2 vector bundle on C of degree 14, with noncanonical determinant, the equation $h^0(C, E) \geq 4$ imposes 16 conditions. Similarly, if $\epsilon \in \text{Pic}^0(C)$, the equation $h^0(C, E \otimes \epsilon) \geq 4$ imposes 16 conditions on the parameter space of E . Given C , there are $29 = 4g - 3$ parameters for E , and $8 = g$ parameters for ϵ . It follows that we have at least a 5-dimensional family of pairs (E, ϵ) , such that

$$h^0(C, E) \geq 4 \quad \text{and} \quad h^0(C, E \otimes \epsilon) \geq 4. \quad (6.33)$$

Furthermore, the construction of Section 5 (together with Proposition 5) shows that for a general triple (C, E, ϵ) as above, one has $K_{2,1}(C, L) \neq 0$, where $L := \det E \otimes \epsilon$. Assuming the map $(E, \epsilon) \mapsto L$

is generically finite on its image, we constructed in this way a five dimensional family of paracanonical line bundles $L \in \text{Pic}^{14}(C)$ with a nontrivial syzygy: $K_{1,2}(C, L) \neq 0$. This family has the following property:

Lemma 30. *If $L = \det E \otimes \epsilon$, where E satisfies (6.33), the line bundle $K_C^{\otimes 2} \otimes L^\vee$ satisfies the same property. The family above, which has dimension at least five, is thus invariant under the involution $L \mapsto K_C^{\otimes 2} \otimes L^\vee$ on P_8^{14} , whose fixed locus is the Prym moduli space \mathcal{R}_8 .*

Proof. This follows from Serre duality, replacing E with $E^\vee \otimes K_C$ and $E \otimes \epsilon$ by $E^\vee \otimes \epsilon^\vee \otimes K_C$ plus the fact that $\det(E^\vee \otimes K_C) \otimes \epsilon^\vee \cong K_C^{\otimes 2} \otimes \det E^\vee \otimes \epsilon^\vee$. \square

One can ask in general the following question:

Question 31. *Is the divisor $\mathfrak{K}053$ on P_8^{14} invariant under the involution $L \mapsto K_C^{\otimes 2} \otimes L^\vee$?*

References

- [1] G. Bruns, *Twists of Mukai bundles and the geometry of the level 3 modular variety over $\overline{\mathcal{M}}_8$* , Trans. Amer. Math. Soc, to appear. [arXiv:1606.07038](#)
- [2] A. Chiodo and G. Farkas, *Singularities of the moduli space of level curves*, J. Eur. Math. Soc. **19** (2017), no. 3, 603–658. [MR-3612865](#)
- [3] A. Chiodo, D. Eisenbud, G. Farkas and F.-O Schreyer, *Syzygies of torsion bundles and the geometry of the level ℓ modular variety over \mathcal{M}_g* , Invent. Math. **194** (2013), no. 1, 73–118. [MR-3103256](#)
- [4] E. Colombo and P. Frediani, *On the Koszul cohomology of canonical and Prym-canonical binary curves*, Bull. Lond. Math. Soc. **45** (2013), no. 5, 1031–1040. [MR-3104994](#)
- [5] M. Cornalba, *Moduli of curves and theta-characteristics*. In: Lectures on Riemann surfaces (Trieste, 1987), pp. 560–589, World Sci. Publ., Teaneck, NJ, 1989. [MR-1082361](#)
- [6] D. Eisenbud, H. Lange, G. Martens and F.-O. Schreyer, *The Clifford dimension of a projective curve*, Compositio Math. **72**(1989), no. 2, 173–204. [MR-1030141](#)
- [7] B. Eröcal, O. Motsak, F.-O. Schreyer and A. Steenpass, *Refined algorithms to compute syzygies*, J. Symbolic Comput. **74** (2016), 308–327. [MR-3424044](#)
- [8] G. Farkas and M. Kemeny, *The Generic Green-Lazarsfeld secant conjecture*, Invent. Math. **203** (2016), no. 1, 265–301. [MR-3437872](#)
- [9] G. Farkas and M. Kemeny, *The Prym-Green Conjecture for torsion bundles of high order*, Duke Math. J. **166** (2017), no. 6, 1103–1124. [MR-3635900](#)
- [10] G. Farkas and K. Ludwig, *The Kodaira dimension of the moduli space of Prym varieties*, J. Eur. Math. Soc. **12** (2010), no. 3, 755–795. [MR-2639318](#)
- [11] G. Farkas and A. Verra, *Prym varieties and moduli of polarized Nikulin surfaces*, Adv. Math. **290** (2016), 314–328. [MR-3451926](#)
- [12] M. Green and R. Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Invent. Math. **83** (1986), no. 1, 73–90. [MR-0813583](#)

- [13] J. Harris, *Theta-characteristics on algebraic curves*, Trans. Amer. Math. Soc. **271** (1982), no. 2, 611–638. [MR-0654853](#)
- [14] S. Mukai, *Vector bundles and Brill-Noether theory*. In: Current topics in complex algebraic geometry (Berkeley, CA, 1992/93), pp. 145–158, Math. Sci. Res. Inst. Publ., 28, Cambridge Univ. Press, Cambridge, 1995. [MR-1397062](#)
- [15] D. Mumford, *Theta characteristics of an algebraic curve*, Ann. Sci. École Norm. Sup. (4) **4** (1971), 181–192. [MR-0292836](#)
- [16] F.-O. Schreyer, *A standard basis approach to syzygies of canonical curves*, J. Reine Angew. Math. **421** (1991), 83–123. [MR-1129577](#)
- [17] C. Voisin, *Sur l'application de Wahl des courbes satisfaisant la condition de Brill-Noether-Petri*, Acta Math. **168** (1992), no. 3–4, 249–272. [MR-1161267](#)
- [18] C. Voisin, *Green's generic syzygy conjecture for curves of even genus lying on a K3 surface*, J. Eur. Math. Soc. **4** (2002), no. 4, 363–404. [MR-1941089](#)
- [19] C. Voisin, *Green's canonical syzygy conjecture for generic curves of odd genus*, Compos. Math. **141** (2005), no. 5, 1163–1190. [MR-2157134](#)