

---

## An atlas of K3 surfaces with finite automorphism group

Xavier Roulleau

**Abstract.** We study the geometry of K3 surfaces with finite automorphism group and Picard number at least 3. We describe these surfaces classified by Nikulin and Vinberg as double covers of simpler surfaces or as embedded in a projective space. We study, moreover, the configurations of their finite set of  $(-2)$ -curves.

**Keywords.** K3 surfaces, finite automorphism groups, finite number of  $(-2)$ -curves, Vinberg algorithms

**2020 Mathematics Subject Classification.** 14J28

arXiv:2003.08985v6 [math.AG] 24 Nov 2022

---

Received by the Editors on April 17, 2020, and in final form on April 26, 2022.

Accepted on June 21, 2022.

Xavier Roulleau

Aix-Marseille Université, CNRS, Centrale Marseille, I2M UMR 7373, 13453 Marseille, France

*e-mail:* [Xavier.Roulleau@univ-amu.fr](mailto:Xavier.Roulleau@univ-amu.fr)

The author is thankful to the Max Planck Institute for Mathematics in Bonn for financial support.

© by the author(s)

This work is licensed under <http://creativecommons.org/licenses/by-sa/4.0/>

## Contents

1. Introduction . . . . .	2
2. Notation, Preliminaries . . . . .	6
3. Rank 3 lattices . . . . .	13
4. Rank 4 lattices . . . . .	30
5. Rank 5 lattices . . . . .	31
6. Rank 6 lattices . . . . .	38
7. Rank 7 lattices . . . . .	46
8. Rank 8 lattices . . . . .	51
9. Rank 9 lattices . . . . .	58
10. Rank 10 lattices . . . . .	64
11. Rank 11 lattices . . . . .	71
12. Rank 12 lattices . . . . .	75
13. Rank 13 lattices . . . . .	80
14. Rank 14 lattices . . . . .	83
15. Rank at least 15 lattices . . . . .	87
16. Table: Number of $(-2)$ -curves . . . . .	91
References . . . . .	93

## 1. Introduction

Algebraic K3 surfaces  $X$  over  $\mathbb{C}$  with finite automorphism group were classified by their Picard lattices  $\text{NS}(X)$  and Picard number  $\rho = \text{rk}(\text{NS}(X)) \geq 3$  by Nikulin for  $\rho = 5, 6, \dots$  in 1981, *cf.* [Nik83], and for  $\rho = 3$  in 1985, *cf.* [Nik85], and by Vinberg for  $\rho = 4$  (in 1981, published in 2007 in [Vin07]). The result of their classification is a list of 118 Néron–Severi lattices.

In the present paper, we exhibit for each of these surfaces a geometric construction, as a double plane, a double cover of the Hirzebruch surface  $F_n$ ,  $n \in \{2, 3, 4\}$  (which part has also been more or less explicitly done in [AN06] and [Zha98]) or a complete intersection surface of degree  $k \in \{4, 6, 8\}$  in  $\mathbb{P}^{\frac{1}{2}k+1}$ . Let us recall that a K3 surface with Picard number  $\rho > 2$  has finite automorphism group if and only if it contains only a non-zero finite number of  $(-2)$ -curves. The  $(-2)$ -curves play a key role for K3 surfaces; for example, one can describe their ample cone using these curves. We are thus especially interested in the configuration of these curves on the K3 surfaces. In [Nik83] and especially in [Nik85] for  $\rho = 3$ , Nikulin described the number and some configurations of these curves. Our contribution to the subject comes for higher-rank Picard lattices,

when there are more  $(-2)$ -curves than expected. Among the K3 surfaces we describe, there are two series which we find most remarkable.

The first series are the surfaces with Néron–Severi lattice of type  $U(2) \oplus \mathbf{A}_1^{\oplus n}$ , for  $n \in \{2, \dots, 7\}$ . Such a surface  $X$  is the desingularization of the double cover of the plane branched over a sextic curve with  $n + 1$  nodes in general position. The surface  $X$  is also naturally the double cover  $X \rightarrow Z$  of a degree  $8 - n$  del Pezzo surface  $Z$  branched over a curve in the linear system  $|-2K_Z|$ . The pull-back on  $X$  of the  $(-1)$ -curves on  $Z$  are  $(-2)$ -curves, and one has a description of the configuration of the  $(-2)$ -curves on  $X$  from the well-known configuration of  $(-1)$ -curves on the del Pezzo surface. In particular, for  $n = 2, \dots, 7$ , the number of  $(-2)$ -curves on the K3 surface is 6, 10, 16, 27, 56, 240, respectively. Let us describe the case with 240  $(-2)$ -curves.

**Theorem 1.1.** *Let  $X$  be a general K3 surface with Néron–Severi lattice isometric to  $U(2) \oplus \mathbf{A}_1^{\oplus 7}$ . There exists a double cover  $f_1: X \rightarrow \mathbb{P}^2$  branched over a sextic curve with 8 nodes  $p_1, \dots, p_8$  such that the images of the 240  $(-2)$ -curves on  $X$  are*

- the 8 points  $p_1, \dots, p_8$ ,
- the 28 lines through  $p_i, p_j$  with  $i \neq j$ ,
- the 56 conics that go through 5 points in  $\{p_1, \dots, p_8\}$ ,
- the 56 cubics that go through 7 points  $p_j$  with a double point at 1 of these points,
- the 56 quartics through the 8 points  $p_j$  with double points at 3 of them,
- the 28 quintics through the 8 points  $p_j$  with double points at 2 of them,
- the 8 sextics through the 8 points with double points at all except a single point with multiplicity 3.

Let  $Z \rightarrow \mathbb{P}^2$  be the blow-up at the points  $p_k, k \in \{1, \dots, 8\}$ ; the surface  $X$  is also a double cover of  $Z$ . The pull-back on  $X$  of the pencil  $|-K_Z|$  gives another double cover  $f_2: X \rightarrow \mathbb{P}^2$ . Its branch locus is a smooth sextic curve to which 120 conics are tangent at every intersection point. These 120 conics are the images of the 240  $(-2)$ -curves on  $X$ .

The two involutions corresponding to the covers  $f_1, f_2$  generate the automorphism group  $(\mathbb{Z}/2\mathbb{Z})^2$  of  $X$ .

The second series are the surfaces with Néron–Severi lattice of type  $U \oplus \mathbf{E}_8 \oplus \mathbf{A}_1^{\oplus 4}$ ,  $U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1^{\oplus 3}$ ,  $U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$ ,  $U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 5}$  or  $U \oplus \mathbf{A}_1^{\oplus 8}$ . These surfaces contain, respectively, 27, 39, 59, 90, 145  $(-2)$ -curves. Let us give an example of the results obtained for the case of  $U \oplus \mathbf{A}_1^{\oplus 8}$ . Let  $F_4$  be the Hirzebruch surface with a section  $s$  such that  $s^2 = -4$ . Let  $f$  be a fiber of the unique fibration, and let  $L = 4f + s$ ,  $L' = 5f + s$ .

**Theorem 1.2.** *Let  $X$  be a general K3 surface with Néron–Severi lattice isometric to  $U \oplus \mathbf{A}_1^{\oplus 8}$ . There exists a double cover  $f_1: X \rightarrow F_4$  branched over a curve  $B = s + b$ , where  $b$  is a curve in  $|3L|$  with 8 nodes  $p_1, \dots, p_8$  and  $s \cap b = \emptyset$ ; that double cover is such that the images of the 145  $(-2)$ -curves on  $X$  are*

- the 8 points  $p_1, \dots, p_8$ ,
- the section  $s$ ,
- the 8 fibers through the 8 points  $p_1, \dots, p_8$ ,
- the 8 curves in  $|L'|$  going through 7 of the 8 points  $p_1, \dots, p_8$ ,
- the 56 sections in the linear system  $|L|$  that pass through 5 of the 8 points  $p_1, \dots, p_8$ ,
- the 56 curves in the linear system  $|2L|$  that pass through 5 of the 8 points with multiplicity 1 and through the 3 remaining with multiplicity 2,
- the 8 curves in the linear system  $|3L|$  that pass through the 8 points with multiplicity 2, except at 1 point where the multiplicity is 3.

There exists, moreover, a double cover of  $\mathbb{P}^2$  branched over a sextic curve with one node. Through that node, there are 8 lines that are tangent to the sextic at another intersection point. There are 64 conics that are 6-tangent to the sextic. The  $72 = 8 + 64$  curves and the node are the images of the 145  $(-2)$ -curves on  $X$ .

The two involutions corresponding to the two covers we described generate the automorphism group  $(\mathbb{Z}/2\mathbb{Z})^2$  of  $X$ .

The other members of the family have a similar description as double covers of  $F_2$  or  $F_4$ , and one can describe their  $(-2)$ -curves similarly. As above, these surfaces have another model as a double cover of  $\mathbb{P}^2$  branched over a sextic curve. In most cases, the set of  $(-2)$ -curves can be naturally decomposed as the union of two sets of  $(-2)$ -curves having a different behavior: one set generates the Néron–Severi lattice; it has a dual graph which is directly related to the name of the lattice and can be easily represented. The second set is a set of extra  $(-2)$ -curves which, when they exist, form an interesting configuration, sometimes very symmetric. Often the curves of the second set occur as pull-backs under a double cover of conics or lines in  $\mathbb{P}^2$  that are tangent to the sextic branch curve at an intersection point.


In this paper, we also describe the configurations of the  $(-2)$ -curves contained in the K3 surfaces. The perhaps nicest configuration is as follows.

**Theorem 1.3.** *Let  $X$  be a general K3 surface with Néron–Severi lattice  $U(4) \oplus A_1^{\oplus 3}$ . The surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$  such that there exist 12 conics that are tangent to the sextic at an intersection point; the 24  $(-2)$ -curves on  $X$  are mapped in pairs to the 12 conics.*

*There exists a partition of the 24  $(-2)$ -curves into three sets  $S_1, S_2, S_3$  of 8 curves each, such that for curves  $B, B'$  in two different sets  $S, S'$ , one has  $BB' \in \{0, 4\}$  and for any  $B \in S$ , there are exactly 4 curves  $B'$  in  $S'$  such that  $BB' = 4$ , and symmetrically for  $B'$ . The sets  $S$  and  $S'$  form an  $8_4$  configuration called the Möbius configuration.*

*The moduli space of K3 surfaces polarized by  $U(4) \oplus A_1^{\oplus 3}$  is unirational.*

The following graph is the Levi graph of the Möbius configuration, where vertices in red are curves in  $S$ , vertices in blue are curves in  $S'$ , and an edge links a red curve to a blue curve if and only if their intersection number is 4:



From that graph, we can moreover read off the intersection numbers of the curves in  $S$  (and  $S'$ ) as follows:

For any red curve  $B$ , there are four blue curves linked to it by an edge. Consider the complementary set of blue curves; this is another set of four blue curves, all linked through an edge to the same red curve  $B'$ . Then we have  $BB' = 6$ , and for any other red curve  $B'' \neq B'$ , we have  $BB'' = 2$ . Symmetrically, the intersection numbers between the blue curves follow the same rule.

Nikulin also studied that configuration of 24 curves, which he obtained by lattice considerations (see [Nik83, Section 8.3]), showing a very nice relation with the 24 roots of  $D_4$ .

Another result is about the “famous” 95 families of K3 hypersurfaces in weighted projective threefolds. In Section 2.6, we remark that among these K3 surfaces, many are surfaces with finite automorphism group, and their moduli space is unirational. We also study the unirationality of many other moduli spaces of K3 surfaces with finite automorphism group.

At the end of the paper, in Section 16, we give the list of the K3 surfaces with finite automorphism group and their number of  $(-2)$ -curves, for Picard number at least 3. In this paper is missing the classification for Picard number 4: that is the subject of another paper [ACR20] with Artebani and Correa Diesler.

In [Kon89], Kondo classifies the automorphism group of general K3 surfaces in each of the 118 families. For 105 of these families, Kondo proves that the automorphism group of a general K3 surface is either trivial

or the group  $\mathbb{Z}/2\mathbb{Z}$ , leaving indeterminate which cases actually happen. Our work clarifies the situation for each of these families; the automorphism group of a general K3 surface of each family can be found in Section 16.

It is not possible for the author to cite every contribution on the subject of K3 surfaces with finite automorphism group. However, let us give some details on [AN06], where Alexeev and Nikulin classify log del Pezzo surfaces of index at most 2 (*i.e.*, normal surfaces  $X$  with quotient singularities such that the anti-canonical divisor  $-K_X$  is ample and  $2K_X$  is a Cartier divisor). They also describe possible configurations of exceptional curves on a minimal resolution of  $X$ , *i.e.*, smooth rational curves with negative self-intersection.

The paper of Alexeev and Nikulin gives the description of the  $(-2)$ -curves in Theorems 1.1 and 1.2. Table 3 in [AN06] gives these configurations for the “special” resolutions of the most degenerate del Pezzo surfaces, those with the largest configuration of  $(-2)$ -curves. But the procedure for the general case is also provided in [AN06]: if a certain ADE configuration disappears on a deformation, then one should add the corresponding Weyl group orbit of the  $(-1)$ -curves. The  $(-2)$ -curves on the double cover K3 surface are the preimages of the  $(-1)$ -,  $(-2)$ - and  $(-4)$ -curves on these resolutions.

For example, for the double covers of the degree 1 del Pezzo surfaces: On the most degenerate “almost” del Pezzo surface, there is an  $E_8$  configuration of  $(-2)$ -curves and a single  $(-1)$ -curve. The corresponding K3 surface has  $16 + 1 = 17$   $(-2)$ -curves. On the most general del Pezzo surfaces, there are no  $(-2)$ -curves, and there is a single  $W(E_8)$ -orbit of  $(-1)$ -curves, which has cardinality  $|W(E_8)/W(E_7)| = 240$ . Indeed, this description of the  $(-1)$ -curves on a del Pezzo surface is well known. The corresponding K3 surface has 240  $(-2)$ -curves.

Table 3 in [AN06] contains this information for all 2-elementary hyperbolic lattices where the K3 surfaces have finite automorphism group, excluding the  $(19, 1, 1)$  case which has  $g = 1$  (see the notation in [AN06]). The lattices in Theorems 1.1 and 1.2 are 2-elementary.

To be more precise, the paper [AN06] computes the fundamental chambers  $P^{2,4}$  for all the 2-elementary lattices in question, for the full reflection group. That group is generated by the reflections in the  $(-2)$ -vectors and in the 2-divisible  $(-4)$ -vectors. And by [AN06, Proposition 2.4.1], the walls of the fundamental chamber  $P^2$  for the Weyl group  $W^2$  are the orbits of the  $(-2)$ -walls of  $P^{2,4}$  under the group generated by the  $(-4)$ -walls of  $P^{2,4}$ . In terms of graphs, the set of walls of  $P^2$  (that is, the  $(-2)$ -curves on  $X$ ) is the union of the orbits of the white vertices in the Coxeter graph of  $P^{2,4}$  (see [AN06, Table 3, p. 93]) under the Weyl group corresponding to the black subgraph.

On the subject of K3 surfaces with finite automorphism group, let us also mention the paper [Zha98], where Zhang classifies the quotients of K3 surfaces by an involution.

At last, let us also cite [AHL10] by Artebani, Hausen and Laface, and [McK10] by McKernan, where the Cox ring of some K3 surfaces is studied. It turns out that a K3 surface has a finitely generated Cox ring if and only if its cone of effective divisor classes is polyhedral or, equivalently, if its automorphism group is finite. K3 surfaces with finite automorphism group and Picard number  $\rho = 2$  have been described by Piatetskii-Shapiro and Shafarevich [PS71] and also studied in [GLP10] (see also [AHL10, Ott13] for their Cox rings). The Cox rings of K3 surfaces with a non-symplectic involution and  $2 \leq \rho \leq 5$  have been described in [AHL10]. Also in [AHL10] is studied the Cox ring for all K3 surfaces which are general double covers of del Pezzo surfaces. Finally, let us mention (see [PS71]) that the general K3 surface with Picard number 1 has trivial automorphism group, unless its Néron–Severi lattice is generated by  $D$  with  $D^2 = 2$ , in which case the automorphism group is generated by the non-symplectic involution.

## Acknowledgements

The author is very grateful to the referees for many suggestions, comments, ideas improving the paper; one of the referees explained to us how to finish the computation of the number of  $(-2)$ -curves for certain difficult cases (see Section 2.3). The author also thanks Alessandra Sarti and Alan Thompson for very useful

email exchanges about moduli spaces of K3 surfaces. Part of this paper was written during the author's stay at Max Planck Institute for Mathematics in Bonn, to which the author is grateful for its hospitality. The computations have been done using Magma software; *cf.* [BCP97].

## 2. Notation, Preliminaries

### 2.1. Notation and conventions

We work over the complex numbers. Linear equivalence between divisors is denoted by  $\equiv$ . On a K3 surface, linear and numerical equivalences coincide. A  $(-2)$ -curve is unique in its numerical equivalence class, so we will often not distinguish between a  $(-2)$ -curve and its equivalence class.

The configuration of a set of  $(-2)$ -curves  $C, C', \dots$  is given by its dual graph, where a vertex represents a  $(-2)$ -curve and two vertices are linked by an edge if the intersection  $C \cdot C'$  (sometimes also denoted by  $CC'$ ) of the corresponding curves  $C, C'$  satisfies  $C \cdot C' > 0$ . Unless explicitly stated, when  $n = C \cdot C'$ , we label the edge  $n$ :



Moreover, a thick edge



between the two vertices  $C, C'$  means that  $C \cdot C' = 2$ , and no label means that  $C \cdot C' = 1$ .

We denote by  $\mathbf{a}_n, \mathbf{d}_n, \mathbf{e}_n$  the Du Val curve singularities, also called simple curve singularities (see [BHP<sup>+</sup>04, Chapter II, Section 8]). Let  $C_6 \hookrightarrow \mathbb{P}^2$  be a reduced plane sextic curve with at most **ade** singularities. Let  $X$  be the K3 surface which is the minimal desingularization of the double cover branched over  $C_6$ . We denote by  $\eta: X \rightarrow \mathbb{P}^2$  the natural map. We say that a line is tritangent to  $C_6$  if the multiplicities at the intersection points of the line and  $C_6$  are even. Similarly, we say that a conic is 6-tangent to  $C_6$  if the multiplicities of the intersection points at the conic and  $C_6$  are even. The following result is well known.

**Lemma 2.1.** *Let  $R_d$  be a tritangent line ( $d = 1$ ) or a 6-tangent conic ( $d = 2$ ). The pull-back on  $X$  of  $R_d$  splits:  $\eta^*R_d = A + B$ , where  $A, B$  are two smooth rational curves with intersection number  $d^2 + 2$ . One has  $D_2 \cdot A = D_2 \cdot B = d$  and  $A + B \equiv dD_2$ , where  $D_2$  is the pull-back by  $\eta$  of a line.*

*Conversely, if two  $(-2)$ -curves  $A, B$  are such that there exist a nef, base-point free divisor  $D_2$  of square 2 and a  $d \in \{1, 2\}$  such that  $dD_2 = A + B$ , then the image of  $A$  and  $B$  by the natural map obtained from  $|D_2|$  is a rational curve of degree  $d$ .*

For a symmetric matrix  $Q$  with integral entries, we denote by  $[Q]$  the lattice with Gram matrix  $Q$ . We denote by  $U$  the lattice  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ . If  $L$  is a lattice and  $m \in \mathbb{Z}$ , then  $L(m)$  is the lattice with the same group as  $L$  but with Gram matrix multiplied by  $m$ .

We denote by  $\mathbf{A}_n, \mathbf{D}_n, \mathbf{E}_n$  the negative-definite lattices that correspond to the root systems denoted with the same letters. A double cover branched over a curve with a singularity  $\mathbf{b}_n$  ( $\mathbf{b}$  in  $\mathbf{a}, \mathbf{d}, \mathbf{e}$ ) is a singular surface with  $B_n$  singularity ( $B$  in  $A, D, E$ ), see [BHP<sup>+</sup>04, Chapter III, Section 6]. Its minimal resolution is by a union of  $(-2)$ -curves, which curves generate a lattice  $\mathbf{B}_n$ .

In this paper, by an elliptic fibration of a K3 surface  $X$ , we mean a morphism  $X \rightarrow \mathbb{P}^1$  with connected fibers. We will frequently use implicitly the linear equivalence relations obtained from (singular) fibers of an elliptic fibration; such information can be read off the dual graph of the  $(-2)$ -curves. The Kodaira classification of singular fibers of elliptic fibrations and their dual graph with their weight can be found, *e.g.*, in [BHP<sup>+</sup>04, Section V.7]. We denote by  $\tilde{\mathbf{A}}_n, \tilde{\mathbf{D}}_n, \tilde{\mathbf{E}}_n$  the types of singular fibers of an elliptic fibration in Kodaira's classification.

Let  $N$  be a lattice of signature  $(1, \rho - 1)$ . We denote by  $\mathcal{M}_N$  the moduli of K3 surfaces  $X$  polarized by a primitive embedding  $N \hookrightarrow \text{NS}(X)$ .

## 2.2. Some results of Saint-Donat on linear systems of K3 surfaces

To be self-contained, let us recall the following results of Saint-Donat [Sai74].

**Theorem 2.2.** *Let  $D$  be a divisor on a K3 surface  $X$ .*

- a) ([Sai74, Proposition 2.6]). *If  $D$  is effective and non-zero and  $D^2 = 0$ , then  $D = aE$ , where  $|E|$  is a free pencil and  $a \in \mathbb{N}$ .*
- b) ([Sai74, (2.7.3)]). *If  $D$  is big and nef, then  $h^0(D) = 2 + \frac{1}{2}D^2$  and either  $|D|$  has no fixed part, or  $D = aE + \Gamma$ , where  $|E|$  is a free pencil and  $\Gamma$  is an effective  $(-2)$ -class such that  $\Gamma \cdot E = 1$ .*
- c) ([Sai74, Section 4.1]). *If  $D$  is big and nef and  $|D|$  has no fixed part, then  $|D|$  is base-point free and either  $\varphi_{|D|}$  is 2-to-1 onto its image (hyperelliptic case), or it maps  $X$  birationally onto its image, contracting the  $(-2)$ -curves  $\Gamma$  such that  $D \cdot \Gamma = 0$  to singularities of type ADE.*

For a divisor  $D$  such that the linear system  $|D|$  has no base points, we denote by  $\varphi_D$  the morphism associated to  $|D|$ . In case the linear system  $|D|$  is hyperelliptic, we have the following.

**Theorem 2.3** (See [Sai74, Proposition 5.7] and its proof). *Let  $|D|$  be a complete linear system on the K3 surface  $X$ . Suppose that  $|D|$  has no fixed components. Then  $|D|$  is hyperelliptic if and only if either*

- a)  $D^2 \geq 4$ , and there is a fiber of an elliptic fibration  $F$  such that  $F \cdot D = 2$ ; or
- b)  $D^2 \geq 4$ , and there is an irreducible curve  $D'$  with  $D'^2 = 2$  and  $D \equiv 2D'$  (thus in that case  $D^2 = 8$ );  
then the image of the map  $\varphi_D$  is the Veronese surface; or
- c)  $D^2 = 2$ ;  
then  $\varphi_D$  is a double cover of  $\mathbb{P}^2$ .

In case a), the image of the associated map  $\varphi_D: X \rightarrow \mathbb{P}^{\frac{1}{2}D^2+1}$  is a rational normal scroll of degree  $\frac{1}{2}D^2$ , except in the following three cases:

- i)  $D \equiv 4F + 2\Gamma$ , where  $F$  is a fiber and  $\Gamma$  is a  $(-2)$ -curve such that  $F\Gamma = 1$ .  
In that case,  $D^2 = 8$ , and  $\varphi_D(X)$  is a cone in  $\mathbb{P}^5$  over a rational normal quartic curve in a hyperplane  $\mathbb{P}^4 \subset \mathbb{P}^5$ . The map  $\varphi_D$  factors through  $X \xrightarrow{\varphi} \mathbb{F}_4 \rightarrow \mathbb{P}^5$ , where  $\varphi$  is a morphism onto the Hirzebruch surface  $\mathbb{F}_4$  and  $\mathbb{F}_4 \rightarrow \mathbb{P}^5$  is the contraction map of the unique section  $s$  such that  $s^2 = -4$ . The branch locus of  $\varphi$  is the union of  $s$  and a reduced curve  $B'$  in the linear system  $|3s + 12f|$  such that  $s \cap B' = \emptyset$  (here  $f$  is a fiber of the unique fibration  $\mathbb{F}_4 \rightarrow \mathbb{P}^1$ ).
- ii)  $D \equiv 3F + 2\Gamma_0 + \Gamma_1$ , where  $\Gamma_0, \Gamma_1$  are  $(-2)$ -curves such that  $\Gamma_0 \cdot F = 1$ ,  $\Gamma_1 \cdot F = 0$  and  $\Gamma_0 \cdot \Gamma_1 = 1$ .  
In that case,  $D^2 = 6$ , and  $\varphi_D(X)$  is a cone in  $\mathbb{P}^4$  over a rational normal cubic curve in a hyperplane  $\mathbb{P}^3 \subset \mathbb{P}^4$ . There is a factorization of  $\varphi_D$  through  $X \xrightarrow{\varphi} \mathbb{F}_3 \rightarrow \mathbb{P}^4$ . The branch locus of  $\varphi$  is the union of the unique section  $s$  such that  $s^2 = -3$  and a reduced curve  $B'$  in the linear system  $|3s + 10f|$  such that  $sB' = 1$ .
- iii) u)  $D \equiv 2F + \Gamma_0 + \Gamma_1$ , where  $\Gamma_0, \Gamma_1$  are  $(-2)$ -curves such that  $\Gamma_0 \cdot F = \Gamma_1 \cdot F = 1$  and  $\Gamma_0 \cdot \Gamma_1 = 0$  (then  $D^2 = 4$ ).  
v)  $D \equiv 2F + 2\Gamma_0 + 2\Gamma_1 + \dots + 2\Gamma_n + \Gamma_{n+1} + \Gamma_{n+2}$ , where  $n \geq 0$  and the  $\Gamma_i$  are  $(-2)$ -curves, such that  $D^2 = 4$ .

In cases iii) u) and iii) v),  $\varphi_D(X)$  is a quadric cone in  $\mathbb{P}^3$ , and there is a factorization  $X \xrightarrow{\varphi} \mathbb{F}_2 \rightarrow \varphi_D(X)$ . In these two cases, the branch locus  $B$  of  $\varphi$  is in  $|4s + 8f|$ . In case u),  $B$  does not contain the section  $s$  such that  $s^2 = -2$ ; in case v), one has  $B = s + B'$ , with  $B'$  reduced and  $sB' = 2$ .

*Remark 2.4.* For brevity, we will say that a divisor  $D$  is base-point free or is hyperelliptic if the associated linear system  $|D|$  is; we hope this will not induce any confusion to the reader.

## 2.3. About the computations

Let  $X$  be a K3 surface with finite automorphism group and Néron–Severi lattice  $L$  of rank  $\rho > 2$ . We did our search of  $(-2)$ -curves on  $X$  using an algorithm of Shimada [Shil4, Section 3] as described in [Rou19].

The inputs of that algorithm are the Gram matrix of a basis of the Néron–Severi lattice and an ample class. For each Néron–Severi lattice  $L$  involved, we computed such an ample class  $D$  as follows. The lattices  $L$  in this paper are mainly of the form

$$L = U(k) \oplus K \quad \text{or} \quad L = [m] \oplus K \quad (k, m \in \mathbb{N}^*),$$

where  $K$  is a direct sum of ADE lattices. To obtain an ample divisor  $D \in L$ , we construct a divisor  $D' \in L$  such that on a set of roots of  $K$  which is also a basis of  $K$ , the divisor  $D'$  has positive intersection. Then we add a suitable multiple  $a$  of an element  $u$  in the  $U[k]$ - or  $[m]$ -part with positive square so that one gets a divisor  $D'' = au + D' \in L$  with  $(D'')^2 > 0$ . We then check that the divisor  $D''$  is ample by verifying that the negative-definite orthogonal complement  $D''^\perp$  does not contain roots, *i.e.*, elements  $v$  such that  $v^2 = -2$  (if this is not the case, then we increase the parameter  $a$ ). Then, the transitivity of the action of the Weyl group

$$W = \langle s_\delta : x \rightarrow x + (x \cdot \delta)\delta \mid \delta \in \Delta \rangle$$

(with  $\Delta = \{\delta \in L \mid \delta^2 = -2\}$ ) on the chambers of the positive cone (see [Huy16, Proposition 8.2.6]) allows us to choose  $D = D''$  as an ample class for the K3 surfaces with lattice  $L$ .

In many cases, the first-found ample divisor  $D$  is such that  $D^2$  is large. By computing the  $(-2)$ -curves on the K3 surface and by using Shimada’s algorithm in [Roul9], we are able to find ample classes with smaller self-intersection. For each lattice, we give the ample class  $D$  with the smallest  $D^2$  we found.

With that knowledge of an ample class  $D$ , one can run Shimada’s algorithm for the computation of the (classes of) the  $(-2)$ -curves  $C$  on  $X$  which have degree  $C \cdot D$  less than or equal to a fixed bound. We then use the test function in [ACL21] to check that we obtain the complete list of  $(-2)$ -curves, which worked well (and confirmed in another way the already known cases by Nikulin), except for the cases

$$U(2) \oplus \mathbf{A}_1^{\oplus 7}, \quad U \oplus \mathbf{A}_1^{\oplus 8}, \quad U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 5}, \quad U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}, \quad U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1^{\oplus 3},$$

where the computation were too heavy to finish (there are too many facets on the effective cone), and thus we have only a lower bound. For these cases, one can obtain the exact number of  $(-2)$ -curves using the following approach communicated to us by one of the referees.

One starts by embedding the lattice  $\text{NS}(X)$  into another Néron–Severi lattice  $\text{NS}'$  of larger rank for which one has already determined the set of classes of  $(-2)$ -curves. For example, let us take

$$II_{1,17} = U \oplus \mathbf{E}_8 \oplus \mathbf{E}_8$$

for  $\text{NS}'$ , as in Section 15.4. The set of  $(-2)$ -curves  $A_1, \dots, A_{19}$  and their configuration had been determined in the classical work of Vinberg [Vin75]. Let  $\mathcal{P}_{18}$  be the positive cone of  $II_{1,17} \otimes \mathbb{R}$  containing an ample class and  $\mathcal{N}_{18}$  the set of the closures of connected components of the complement of the union of all hyperplanes  $(r)^\perp$  of  $\mathcal{P}_{18}$  defined by  $(-2)$ -vectors  $r$ . Suppose that we have a primitive embedding

$$\text{NS}(X) \hookrightarrow \text{NS}' = II_{1,17},$$

and let  $\mathcal{P}_X$  be the positive cone of  $\text{NS}(X) \otimes S$  containing an ample class. We assume that  $\mathcal{P}_X$  is embedded into  $\mathcal{P}_{18}$  and regard  $\mathcal{P}_X$  as a subspace of  $\mathcal{P}_{18}$ . We consider the closed subsets

$$\mathcal{P}_X \cap N' \quad (N' \in \mathcal{N}_{18})$$

of  $\mathcal{P}_X$  that contain a non-empty open subset of  $\mathcal{P}_X$ . In [Shil5], these closed subsets  $\mathcal{P}_X \cap N'$  are called induced chambers. Let  $N_X$  be the closure of the ample cone of  $\text{NS}(X)$ . Since  $N_X$  has only finitely many walls, the cone  $N_X$  is tessellated by a finite number of induced chambers. By the method in [Shil5], we can determine the set of induced chambers contained in  $N_X$ , and hence the set of walls of  $N_X$ , that is, the classes of  $(-2)$ -curves on  $X$ . More precisely, let  $G$  be the subgroup of  $O(\text{NS}(X))$  consisting of the isometries  $g \in O(\text{NS}(X))$  that preserve  $N_X$  and extend to an isometry of  $\text{NS}' = II_{1,17}$ . We can calculate a complete set of representatives of orbits of the action of  $G$  on the set of induced chambers contained in  $N_X$ .



For each  $\text{NS}(X)$  in the above five lattices, we embed  $\text{NS}(X)$  into  $II_{1,17}$  primitively as follows. Let  $\rho$  be the rank of  $\text{NS}(X)$ , and let  $A_{i_1}, \dots, A_{i_{18-\rho}}$  be the first  $18 - \rho$  elements of the sequence

$$A_3, A_5, A_7, A_9, A_1, A_{17}, A_{15}, A_{13}, A_{11}$$

of  $(-2)$ -curves  $A_1, \dots, A_{19}$  in Section 15.4. Then  $\text{NS}(X)$  is isometric to the orthogonal complement of these classes  $A_{i_1}, \dots, A_{i_{18-\rho}}$  in  $II_{1,17}$ . The order of  $G$  and the orbit decomposition of induced chambers in  $N_X$  under the action of  $G$  are given in the following table, where, for example, 42456960 and  $[1, 55], [2, 7]$  in the first line mean that there exist 55 orbits of size  $|G|$  and 7 orbits of size  $|G|/2$  (with stabilizer group of order 2), and hence there exist

$$\left(55 + \frac{7}{2}\right)|G| = 42456960$$

induced chambers contained in  $N_X$ .

$\rho$	$\text{NS}(X)$	$\#(-2)$	$ G $	Chambers	Orbits
9	$U(2) \oplus \mathbf{A}_1^{\oplus 7}$	240	725760	42456960	$[1, 55], [2, 7]$
10	$U \oplus \mathbf{A}_1^{\oplus 8}$	145	80640	30683520	$[1, 372], [2, 17]$
11	$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 5}$	90	10080	10800720	$[1, 1061], [2, 21]$
12	$U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$	59	1440	2400480	$[1, 1649], [2, 36]$
13	$U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1^{\oplus 3}$	39	240	376200	$[1, 1557], [2, 21]$

We give at the end of this atlas a table with the number of  $(-2)$ -curves in the 118 different cases.

One can also compute the finite set of all fibrations on  $X$  as follows. By [Kov94, Proposition 2.4], if the class of a fiber  $F$  in  $\text{NS}(X)$  is indecomposable (*i.e.*, is not the sum of two effective divisors), then it is an extremal class in the closure of the effective cone. But by [Kov94, Theorem 6.1], since there are only a finite number of  $(-2)$ -curves in our surface  $X$ , the  $(-2)$ -curves are the extremal classes. Therefore, there is a singular fiber of the elliptic fibration which is the sum of  $(-2)$ -curves on  $X$ .

Now Kodaira's classification gives a finite number of possibilities for the types  $\tilde{\mathbf{A}}_n, \tilde{\mathbf{D}}_n, \tilde{\mathbf{E}}_n$  of the reducible fibers  $F$  on  $X$ . Since a fiber  $F$  in  $\text{NS}(X)$  is a sum of  $(-2)$ -curves (which are finitely many), there is a finite number of possible degrees  $FD$  of fibers  $F$  with respect to  $D$ . Thus there is an upper bound for  $FD$ , so that one can compute all fibrations on  $X$  using Shimada's algorithm (see [Roul9]).

For the computations, we used Magma [BCP97]. Our algorithms are available as ancillary files of the submission of this paper on arXiv.

## 2.4. Algorithms for double planes

In this paper, many double plane models of K3 surfaces are studied. Algorithms to obtain geometric properties of the double covering (singularities of the branch curve  $B$ , configuration of irreducible components of  $B$ , enumeration of splitting curves, ...) from the lattice data are developed in the paper [Shil0]. They can be modified to the case where the linear system has a fixed component. Many arguments in the present paper (expressing a class  $D_2$  such that  $(D_2)^2 = 2$  as various sums of classes of  $(-2)$ -curves with non-negative coefficients) can be automated by these algorithms.

## 2.5. Nikulin star-shaped dual graphs and lattices of type $U \oplus K$

In [Nik83], Nikulin obtained the following result.

**Proposition 2.5** ([Nik83, Corollary 1.6.5]; see also [Kon89, Lemma 3.1]). *Let  $X$  be a K3 surface. Assume that  $\text{NS}(X) \simeq U \oplus K$ .*

- i) There is an elliptic pencil  $\pi: X \rightarrow \mathbb{P}^1$  with a section. A fiber and the section generate the lattice isometric to  $U$ .*

ii) If  $K = \mathbf{G} \oplus K'$  with the lattice  $\mathbf{G}$  generated by irreducible elements of square  $-2$ , then  $\pi$  has a singular fiber of type  $\tilde{\mathbf{G}}$ , where  $\mathbf{G}$  is among the lattices

$$\mathbf{A}_n, n \geq 1, \quad \mathbf{D}_n, n \geq 4, \quad \mathbf{E}_6, \mathbf{E}_7, \mathbf{E}_8.$$

Assume  $\mathrm{NS}(X) \simeq U \oplus \bigoplus_{i \in I} \mathbf{G}^{(i)}$ , where the  $\mathbf{G}^{(i)}$  are lattices generated by irreducible elements of square  $-2$ . The dual graph of the  $(-2)$ -curves contained in the singular fibers of type  $\tilde{\mathbf{G}}^{(i)}$  and the section form a (so-called) star-shaped graph (see [Nik83]), which we denote by  $\mathrm{St}(\mathrm{NS}(X))$ .

**Theorem 2.6** (See [Nik83, Theorem 0.2.2 and Corollary 3.9.1]). *Let  $X$  be a K3 surface with Néron–Severi lattice isometric to  $U \oplus K$ , where  $K$  is in the following list:*

$$\begin{aligned} & \mathbf{A}_2; \quad \mathbf{A}_1 \oplus \mathbf{A}_2, \mathbf{A}_3; \quad \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_2, \mathbf{A}_2^{\oplus 2}, \mathbf{A}_1 \oplus \mathbf{A}_3, \mathbf{A}_4; \\ & \quad \mathbf{A}_1 \oplus \mathbf{A}_2^{\oplus 2}, \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_3, \mathbf{A}_2 \oplus \mathbf{A}_3, \mathbf{A}_1 \oplus \mathbf{A}_4, \mathbf{A}_5, \mathbf{D}_5; \\ & \quad \mathbf{A}_2^{\oplus 3}, \mathbf{A}_3^{\oplus 2}, \mathbf{A}_2 \oplus \mathbf{A}_4, \mathbf{A}_1 \oplus \mathbf{A}_5, \mathbf{A}_6, \mathbf{A}_2 \oplus \mathbf{D}_4, \mathbf{A}_1 \oplus \mathbf{D}_5, \mathbf{E}_6; \quad \mathbf{A}_7, \\ & \quad \mathbf{A}_3 \oplus \mathbf{D}_4, \mathbf{A}_2 \oplus \mathbf{D}_5, \mathbf{D}_7, \mathbf{A}_1 \oplus \mathbf{E}_6; \quad \mathbf{A}_2 \oplus \mathbf{E}_6; \quad \mathbf{A}_2 \oplus \mathbf{E}_8; \quad \mathbf{A}_3 \oplus \mathbf{E}_8. \end{aligned}$$

*The K3 surface has finite automorphism group, and  $\mathrm{St}(\mathrm{NS}(X))$  is the dual graph of all  $(-2)$ -curves on  $X$ .*

Let  $X$  be a K3 surface such that  $\mathrm{NS}(X) \simeq U \oplus \bigoplus_{i \in I} \mathbf{G}^{(i)}$ , where the  $\mathbf{G}^{(i)}$  are lattices generated by irreducible elements of square  $-2$ . Let  $F$  be a fiber of the natural fibration  $\pi: X \rightarrow \mathbb{P}^1$  and  $E$  be the section as in Proposition 2.5. The divisor  $D_2 = 2F + E$  is nef of square 2, with base points since  $D_2F = 1$ . By Theorem 2.3, the divisor  $D_8 = 2D_2$  is base-point free and hyperelliptic, and it defines a morphism  $X \rightarrow \mathbb{P}^5$  which factors through the Hirzebruch surface  $\mathbf{F}_4$  so that the branch locus of  $\eta: X \rightarrow \mathbf{F}_4$  is the disjoint union of the unique section  $s$  such that  $s^2 = -4$  and  $B$ , a curve in the linear system  $|3s + 12f|$ , where  $f$  is a fiber of the unique fibration of  $\mathbf{F}_4$ . We immediately have the following.

**Proposition 2.7.** *The image by  $\eta$  of the section  $E$  is the section  $s$ ; the pull-back on  $X$  of the pencil  $|f|$  is the pencil of elliptic curves in the elliptic fibration  $\pi: X \rightarrow \mathbb{P}^1$ . A singular fiber of  $\pi$  of type  $\tilde{\mathbf{A}}_n, \tilde{\mathbf{D}}_n, \tilde{\mathbf{E}}_n$  is mapped onto a fiber of  $\mathbf{F}_4 \rightarrow \mathbb{P}^1$  that cuts  $B$  at, respectively, an  $\mathbf{a}_n, \mathbf{d}_n, \mathbf{e}_n$  singularity of  $B$ .*

## 2.6. About the famous 95, their moduli spaces and K3 surfaces with finite automorphism group

In [Bel02] are studied the Néron–Severi lattices of the so-called “famous 95” families of K3 surfaces. These “famous 95” have been constructed by Reid (unpublished); the list of these families appeared in Yomemura [Yon90] from the point of view of singularity theory. The K3 surfaces involved are (singular) anti-canonical divisors in weighted projective threefolds  $\mathbb{W}\mathbb{P}^3 = \mathbb{W}\mathbb{P}^3(\bar{a})$  (here  $\bar{a}$  is the weight  $\bar{a} = (a_1, \dots, a_4)$ ). As we will see, many of these K3 surfaces have finite automorphism group.

Let  $d_{\bar{a}}$  be the degree of an anti-canonical divisor in  $\mathbb{W}\mathbb{P}^3$ . It turns out that each general degree  $d_{\bar{a}}$  surface  $\bar{X}$  in  $\mathbb{W}\mathbb{P}^3$  has the same singularities, and its minimal desingularization is a K3 surface  $X$ . The main result of [Bel02] is the computation of the Néron–Severi lattice  $\mathrm{NS}(X)$  for a general member of each of the 95 families.

For  $\bar{a}$  among the 95 possible weights, let  $L_{\bar{a}} \simeq \mathrm{NS}(X)$  be the lattice of the Néron–Severi lattice of a general K3 surface  $X$  with singular model  $\bar{X} \subset \mathbb{W}\mathbb{P}^3(\bar{a})$ , and let  $\mathcal{M}_{\bar{a}}$  be the moduli space of  $L_{\bar{a}}$ -polarized K3 surfaces.

**Proposition 2.8.** *There is a birational map between  $\mathcal{M}_{\bar{a}}$  and the moduli space of degree  $d_{\bar{a}}$  surfaces in  $\mathbb{W}\mathbb{P}^3(\bar{a})$  modulo automorphisms.*

*Proof.* For each of the 95 cases of  $\bar{a}$ , we compute the dimension of the quotient space  $\mathcal{Q} = \mathbb{P}(H^0(\mathbb{W}\mathbb{P}^3(\bar{a}), \mathcal{O}(d_{\bar{a}}))^*) / \mathrm{Aut}(\mathbb{W}\mathbb{P}^3(\bar{a}))$ , using the formula

$$\dim \mathrm{Aut}(\mathbb{W}\mathbb{P}^3(\bar{a})) = -1 + \sum h^0(\mathbb{W}\mathbb{P}^3, \mathcal{O}(a_k)).$$

The 95 weights  $\bar{a}$  are given in [Bel02, Table 3]; the degrees  $d_{\bar{a}}$  can be found in [Yon90, Table 4.6]. It turns out that this quotient has dimension  $20 - \rho$ , where  $\rho = \text{rank}(L_{\bar{a}})$  is the Picard number. There is a natural injective map from an open set in  $\mathcal{Q}$  to the moduli space  $\mathcal{M}_{\bar{a}}$ ; since  $\mathcal{M}_{\bar{a}}$  is also  $(20 - \rho)$ -dimensional, both spaces are birational.  $\square$

A direct consequence of Proposition 2.8 is the following.

**Corollary 2.9.** *The moduli spaces of the famous 95 polarized K3 surfaces are unirational.*

Comparing with Nikulin and Vinberg's lists, among the lattices  $L_{\bar{a}}$  associated to the 95 weights  $\bar{a}$ , at least 43 lattices are such that the general K3 surface  $X$  with  $\text{NS}(X) \simeq L_{\bar{a}}$  (and a model  $\bar{X} \subset \mathbb{W}\mathbb{P}^3(\bar{a})$ ) has only a finite number of automorphisms. However, there are some weights  $\bar{a}, \bar{b}$  among the 95 such that  $L_{\bar{a}} \simeq L_{\bar{b}}$  (then there are two singular models of the same K3 surface; see [KM12]). Without counting the repetitions, one get (at least) 29 lattices which are lattices among the famous 95 and are also lattices of K3 surfaces with finite automorphism group. These lattices are

$$\begin{aligned} & [2], [4], U, U(2), U \oplus \mathbf{A}_1, U(2) \oplus \mathbf{A}_1, U \oplus \mathbf{A}_2, U(2) \oplus \mathbf{D}_4, U \oplus \mathbf{D}_4, \\ & U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1, U \oplus \mathbf{D}_5, U \oplus \mathbf{A}_2 \oplus \mathbf{D}_4, U \oplus \mathbf{E}_6, U \oplus \mathbf{E}_7, U \oplus \mathbf{A}_1 \oplus \mathbf{E}_6, \\ & U \oplus \mathbf{E}_8, U \oplus \mathbf{E}_7 \oplus \mathbf{A}_1, U \oplus \mathbf{D}_4^{\oplus 2}, U \oplus \mathbf{A}_2 \oplus \mathbf{E}_6, U \oplus \mathbf{E}_8 \oplus \mathbf{A}_1, \\ & U(2) \oplus \mathbf{D}_4^{\oplus 2}, U \oplus \mathbf{A}_2 \oplus \mathbf{E}_8, U \oplus \mathbf{E}_8 \oplus \mathbf{A}_1^{\oplus 3}, U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1^{\oplus 3}, U \oplus \mathbf{D}_8 \oplus \mathbf{D}_4, \\ & U \oplus \mathbf{E}_8 \oplus \mathbf{D}_6, U \oplus \mathbf{E}_8 \oplus \mathbf{E}_7, U \oplus \mathbf{E}_8 \oplus \mathbf{E}_8, U \oplus \mathbf{E}_8 \oplus \mathbf{E}_8 \oplus \mathbf{A}_1, \end{aligned}$$

plus the lattice  $\begin{pmatrix} 2 & 1 \\ 1 & -2 \end{pmatrix}$ . For these lattices  $L_{\bar{a}}$ , we thus have a complete description of the K3 surfaces  $X$  with finite-order automorphism group with  $\text{NS}(X) \simeq L_{\bar{a}}$  as singular model(s) in weighted projective space(s).

It is worth mentioning that the 95 families have mirrors (see [Bel02]); not all mirrors are among the 95 families, but many mirrors are K3 surfaces with finite automorphism group. The families that do not already appear in the above list are

$$\begin{aligned} & U \oplus \mathbf{A}_1^{\oplus 2}, U(3) \oplus \mathbf{A}_2, U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2, U \oplus \mathbf{A}_2^{\oplus 2}, \\ & U \oplus \mathbf{A}_2 \oplus \mathbf{A}_3, U \oplus \mathbf{A}_5, U \oplus \mathbf{E}_8 \oplus \mathbf{A}_3. \end{aligned}$$

## 2.7. On the automorphism groups

In [Kon89], Kondo studies the automorphism group of a general K3 surface with a finite number of  $(-2)$ -curves and Picard number at least 3. The main result of Kondo's paper [Kon89, Table 1] is that for a certain list of 12 lattices among the 118 lattices, the automorphism group is  $(\mathbb{Z}/2\mathbb{Z})^2$ , that for  $U \oplus \mathbf{E}_8^{\oplus 2} \oplus \mathbf{A}_1$  it is  $\mathfrak{S}_3 \times \mathbb{Z}/2\mathbb{Z}$  and that otherwise, for the remaining 105 families, it is either the trivial group or  $\mathbb{Z}/2\mathbb{Z}$ .

Our study enables us to construct a hyperelliptic involution for surfaces in 90 out of 105 families, and we prove that the surfaces in the remaining 15 families have trivial automorphism group, thus completing in that way the results of Kondo. These results are summarized in the table in Section 16.

*Remark 2.10.* For the ‘‘general’’ assumption on the surface, which can be made more precise, we refer to the introduction of Kondo's paper [Kon89]. That hypothesis is important since Kondo constructed special K3 surfaces with finite automorphism group isomorphic to  $\mathbb{Z}/42\mathbb{Z}$  or  $\mathbb{Z}/66\mathbb{Z}$  (see [Kon86]). When computing the automorphism group, the K3 surfaces we consider are always supposed general.

When the K3 surface  $X$  has automorphism group isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^2$  and Picard number at most 14, we describe the two hyperelliptic involutions generating the automorphism group  $\text{Aut}(X)$ ; when the Picard number is larger, we refer to the description in [Kon89].

For computing the automorphism group of a K3 surface  $X$ , it is important to know the image of the natural map

$$\varphi: \text{Aut}(X) \rightarrow O(\text{NS}(X)).$$

It turns out that when the automorphism group is not trivial, there is always a hyperelliptic involution  $\sigma$ . Using our description of the set of  $(-2)$ -curves, one can understand when an involution  $\sigma$  is in the kernel of  $\varphi$ . One may also use Proposition 2.7 in the case when the K3 lattice is of type  $U \oplus \bigoplus \tilde{G}_j$  as follows. When a K3 surface is the double cover branched over a curve with only singularities of type

$$\mathbf{a}_1, \mathbf{d}_4, \mathbf{d}_6, \mathbf{d}_8, \mathbf{e}_7, \mathbf{e}_8,$$

the action on  $(-2)$ -curves in the star of  $U \oplus \bigoplus \tilde{G}_j$  is trivial, and if there are any singularities of type

$$\mathbf{a}_2, \dots, \mathbf{a}_5, \mathbf{d}_5, \mathbf{d}_7, \mathbf{e}_6,$$

then the involution acts non-trivially on the set of  $(-2)$ -curves in the star of  $U \oplus \bigoplus \tilde{G}_j$ . Since the Néron-Severi lattice is always generated by these  $(-2)$ -curves, one can then understand when an involution is in the kernel of  $\varphi$ . In the table in Section 16, we indicate when the action of a hyperelliptic involution on the Néron-Severi lattice is not trivial.

Let  $X$  be a K3 surface for which Kondō proved that  $\text{Aut}(X)$  is trivial or  $\mathbb{Z}/2\mathbb{Z}$ .

**Proposition 2.11.** *Suppose that  $X$  is general and the rank of  $\text{NS}(X)$  is less than or equal to 8 and*

- i) there is no  $(-2)$ -curve  $A$  and fiber  $F$  such that  $A \cdot F = 1$ ;*
- ii) there is no big and nef divisor  $D$  such that  $D \cdot F = 2$  for a fiber  $F$ ;*
- iii) there is no big and nef divisor  $D$  such that  $D^2 = 2$ .*

*Then the automorphism group of  $X$  is trivial.*

Here, a fiber  $F$  means an irreducible curve with  $F^2 = 0$ .

*Proof.* By Theorems 2.2 and 2.3, the hypotheses i), ii) and iii) imply that there is no hyperelliptic involution acting on  $X$ . By [Kon89, Table 1 and Lemma 2.3], since the rank of  $\text{NS}(X)$  is less than or equal to 8, the automorphism group is either trivial or generated by a hyperelliptic involution; thus it must be trivial.  $\square$

## 2.8. On the irreducibility of the moduli spaces

The aim of this section is to prove the following result.

**Proposition 2.12.** *The 118 moduli spaces of K3 surfaces with Picard number at least 3 and finite automorphism group are irreducible.*

Let us recall that if  $L$  is an even lattice of rank  $\rho$  and signature  $(1, \rho - 1)$ , we denote by  $\mathcal{M}_L$  the moduli space of K3 surfaces  $X$  polarized by a primitive embedding  $j_X: L \hookrightarrow \text{NS}(X)$ . The moduli space  $\mathcal{M}_L$  may depend upon the choice of the embedding of  $L$  in the K3 lattice  $\Lambda_{K3} = U^{\oplus 3} \oplus \mathbb{E}_8^{\oplus 2}$ : two non-isometric embeddings will give two different moduli spaces (see [Nik80] or [Dol96]). For that question, one can use the following result.

**Theorem 2.13** (See [Huy16, Theorem 14.1.12]). *Let  $\Lambda$  be an even unimodular lattice of signature  $(n_+, n_-)$  and  $M$  be an even lattice of signature  $(m_+, m_-)$ . If  $m_+ < n_+$ ,  $m_- < n_-$  and*

$$(2.1) \quad \ell(M) + 2 \leq \text{rk}(\Lambda) - \text{rk}(M),$$

*then there exists a primitive embedding  $M \hookrightarrow \Lambda$ , which is unique up to automorphisms of  $\Lambda$ .*

Here  $\ell(M)$  is the minimal number of generators of the discriminant group of  $M$  and  $\text{rk}(M)$  is its rank.

*Remark 2.14.* Since  $\ell(M) \leq \text{rk}(M)$ , if  $\text{rk}(M) \leq \frac{1}{2}(\text{rk}(\Lambda) - 2)$ , then condition (2.1) is verified.

In our situation, when  $\Lambda = \Lambda_{K3}$  and  $M = L$  is an even lattice of signature  $(1, \text{rk}(L) - 1)$ , one has the following.

**Corollary 2.15.** *Suppose  $\ell(L) \leq 20 - \text{rk}(L)$ . Then the primitive embedding  $L \hookrightarrow \Lambda_{K3}$  is unique up to automorphisms.*

Suppose that the embedding  $j: L \rightarrow \Lambda_{K3}$  is unique up to automorphisms. The following criterion of Dolgachev may be used to check if the moduli space  $\mathcal{M}_L$  is irreducible.

**Theorem 2.16** (See [Dol96, Proposition 5.6]). *Suppose that  $L^\perp \subset \Lambda_{K3}$  contains a sublattice isometric to  $U$  or  $U(2)$ . Then the moduli  $\mathcal{M}_L$  is irreducible.*

Let  $L$  be an even lattice of signature  $(1, \text{rk}(L) - 1)$  such that  $\ell(L) \leq 18 - \text{rk}(L)$ .

**Corollary 2.17.** *The moduli space  $\mathcal{M}_L$  is irreducible.*

*Proof.* By Theorem 2.13, one can find a primitive embedding of  $L$  into the sublattice  $U^{\oplus 2} \oplus E_8^{\oplus 2}$  of  $\Lambda_{K3}$ . Then the orthogonal complement of  $L$  in  $\Lambda_{K3}$  contains a copy of  $U$ , and therefore by Theorem 2.16, the moduli space  $\mathcal{M}_L$  is irreducible.  $\square$

Using that  $\ell(L) \leq \text{rk}(L)$ , one obtains the following.

**Corollary 2.18.** *Suppose moreover that  $L$  has rank at most 9. Then the moduli space  $\mathcal{M}_L$  is irreducible.*

The following table gives the discriminant groups of ADE lattices:

$L$	$A_n (n \geq 1)$	$D_{2n} (n \geq 2)$	$D_{2n+1} (n \geq 2)$	$E_6$	$E_7$	$E_8$
$L^*/L$	$\mathbb{Z}/(n+1)\mathbb{Z}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$\mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	$\{0\}$

There are 28 lattices  $L$  of rank at least 10 such that a K3 surface  $X$  with  $\text{NS}(X) \simeq L$  has finite automorphism group. For  $L$  among these lattices with  $L \neq U \oplus E_8^{\oplus 2} \oplus A_1$ , using the above table and Corollary 2.17, one obtains that the moduli  $\mathcal{M}_L$  is irreducible. For  $L = U \oplus E_8^{\oplus 2} \oplus A_1$ , the embedding in  $U^{\oplus 3} \oplus E_8^{\oplus 2}$  is unique up to automorphisms, and one sees that its orthogonal complement contains a copy of  $U$ ; therefore,  $\mathcal{M}_L$  is also irreducible. We have therefore proved Proposition 2.12.

### 3. Rank 3 lattices

#### 3.1. Rank 3 cases

In [Nik85], Nikulin classifies rank 3 lattices that are Néron–Severi lattices of K3 surfaces with a finite number of automorphisms. Let us describe the classification when the fundamental domain of the associated Weyl group is not compact (see [Nik85, Section 2]).

Let  $S_{1,1,1}$  be the rank 3 lattice generated by vectors  $a, b, c$  with intersection matrix

$$\begin{pmatrix} -2 & 0 & 1 \\ 0 & -2 & 2 \\ 1 & 2 & -2 \end{pmatrix}.$$

For  $r, s, t \in \mathbb{Z}$ , let  $S_{r,s,t}$  denote the sublattice of  $S_{1,1,1}$  generated by  $ra, sb, tc$ . Let also define  $S'_{4,1,2}$ , the lattice generated by  $2a + c, b, 2c$ , and  $S'_{6,1,2}$ , the lattice generated by  $6a + c, b, 2c$ . In [Nik85, Theorem 2.5], Nikulin gives a list of lattices which are Néron–Severi lattices of K3 surfaces with finite automorphism group. These lattices are of the form  $S_{r,s,t}$  or  $S'_{r,s,t}$ . In [Nik14], Nikulin observes that some lattices in the list are isometric, thus giving the same families of K3 surfaces. One has

$$S_{2,1,2} \simeq S_{4,1,1}, \quad S_{4,1,2} \simeq S_{8,1,1}, \quad S_{6,1,2} \simeq S_{12,1,1}, \quad S'_{6,1,2} \simeq S_{6,1,1},$$

and when the fundamental domain is not compact, there are only 20 distinct cases. In the compact case, there are 6 lattices  $S_1, \dots, S_6$ , see [Nik85], and their geometric description is given in [Rou19] (see also [ACL21] for their Cox ring). So the total number of (isomorphism classes of) lattices of rank 3 which are Néron–Severi lattices of K3 surfaces with finite automorphism group is 26. For each case, Nikulin gives the number and sometimes the configurations of the  $(-2)$ -curves.

### 3.2. The rank 3 and compact cases

In [Rou19], we studied the six lattices  $S_1, \dots, S_6$  of rank 3 such that the fundamental domain associated to the Weyl group is compact. These lattices are

$$\begin{aligned} S_1 &= [6] \oplus \mathbf{A}_1^{\oplus 2}, & S_2 &= [36] \oplus \mathbf{A}_2, & S_3 &= [12] \oplus \mathbf{A}_2, \\ S_4 &\subset [60] \oplus \mathbf{A}_2, & S_5 &= [4] \oplus \mathbf{A}_2, & S_6 &\subset [132] \oplus \mathbf{A}_2, \end{aligned}$$

where the two inclusions have index 3. For completeness, let us recall the obtained results.

**Theorem 3.1.** *The K3 surfaces of type  $S_1, S_4, S_5, S_6$  are double covers of the plane branched over a smooth sextic curve  $C_6$  and such that*

- *if the Néron–Severi lattice is isometric to  $S_1$ , the six  $(-2)$ -curves on  $X$  are pull-backs of three conics that are 6-tangent to the sextic  $C_6$ ;*
- *if the Néron–Severi lattice is isometric to  $S_4$ , the four  $(-2)$ -curves on  $X$  are pull-backs of a line tritangent to  $C_6$  and a conic 6-tangent to  $C_6$ ;*
- *if the Néron–Severi lattice is isometric to  $S_5$ , the four  $(-2)$ -curves on  $X$  are pull-backs of two lines tritangent to  $C_6$ ;*
- *if the Néron–Severi lattice is isometric to  $S_6$ , the six  $(-2)$ -curves on  $X$  are pull-backs of one 6-tangent conic and two cuspidal cubics that cut  $C_6$  tangentially and at their cusps.*

*Let  $X$  be a K3 surface that has a Néron–Severi lattice isometric to  $S_2$ . There are three quadrics in  $\mathbb{P}^3$  such that each intersection with  $X \hookrightarrow \mathbb{P}^3$  is the union of two smooth degree 4 rational curves. These six rational curves are the only  $(-2)$ -curves on  $X$ .*

*Let  $X$  be a K3 surface that has a Néron–Severi lattice isometric to  $S_3$ . There exist two hyperplane sections such that each hyperplane section is a union of two smooth conics. These four conics are the only  $(-2)$ -curves on  $X$ .*

The cases  $S_3$  and  $S_4$  are linked to the surfaces  $S_{1,1,4}$  and  $S_{1,1,3}$ ; see Sections 3.6 and 3.5 below, respectively. We will only add the following result.

**Proposition 3.2.** *A general K3 surface  $X$  with a Néron–Severi lattice isometric to  $S_2$  or  $S_3$  has trivial automorphism group.*


*Proof.* The Hilbert bases of the nef cones of these K3 surfaces are described in [ACL21]. Then as in Proposition 3.11 below, one can check that there is no hyperelliptic involution and conclude that the automorphism group is trivial.  $\square$

### 3.3. The lattice $S_{1,1,1}$

Let  $X$  be a K3 surface with rank 3 Néron–Severi lattice and intersection form

$$\begin{pmatrix} -2 & 0 & 1 \\ 0 & -2 & 2 \\ 1 & 2 & -2 \end{pmatrix}.$$

The surface  $X$  contains three  $(-2)$ -curves  $A_1, A_2, A_3$ , with intersection matrix as above; their dual graph is



The divisor

$$D_{22} = 3A_1 + 6A_2 + 7A_3$$

is ample, of square 22, with  $D_{22} \cdot A_1 = D_{22} \cdot A_3 = 1$ ,  $D_{22} \cdot A_2 = 2$ . The divisor

$$(3.1) \quad D_2 = A_1 + 2A_2 + 2A_3$$


is nef, of square 2, with base points since  $F = A_2 + A_3$  is a fiber of an elliptic fibration and  $D_2F = 1$ . It satisfies  $D_2 \cdot A_1 = D_2 \cdot A_2 = 0$ ,  $D_2 \cdot A_3 = 1$ . The divisor  $D_8 = 2D_2$  is base-point free and hyperelliptic. By Theorem 2.3, case i), we have the following.

**Proposition 3.3.** *The linear system  $|D_8|$  defines a map  $\varphi: X \rightarrow \mathbf{F}_4$  onto the Hirzebruch surface  $\mathbf{F}_4$  such that the branch locus of  $\varphi$  is the disjoint union of the unique section  $s$  such that  $s^2 = -4$  and a reduced curve  $B'$  in the linear system  $|3s + 12f|$ .*

By equation (3.1), the image of  $A_1$  is the curve  $s$ , and the image of  $A_3$  is the fiber through the point  $q$  onto which  $A_2$  is contracted. The curve  $B'$  has a unique singularity, which is a node at  $q$ ; its geometric genus is therefore 9.

### 3.4. The lattice $S_{1,1,2}$

Let  $X$  be a K3 surface with rank 3 Néron–Severi lattice of type  $S_{1,1,2}$ . The surface  $X$  contains three  $(-2)$ -curves  $A_1, A_2, A_3$ , with dual graph



The divisor

$$D_{14} = 2A_1 + 2A_2 + 3A_3$$

is ample, of square 14, with  $D_{14} \cdot A_1 = D_{14} \cdot A_2 = 2$  and  $D_{14} \cdot A_3 = 4$ . The divisor

$$D_2 = A_1 + A_2 + A_3$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = 0$  and  $D_2 \cdot A_3 = 2$ . Thus, the following holds.

**Proposition 3.4.** *The K3 surface is the double cover of  $\mathbb{P}^2$  branched over a sextic curve with two nodes  $p, q$ . The curves  $A_1, A_2$  are contracted to  $p, q$ , and the image of  $A_3$  is the line through  $p, q$ . That line cuts the sextic curve transversally in two other points.*

The Severi variety of plane curves of degree 6 with two nodes is rational, so the moduli space  $\mathcal{M}_{S_{1,1,2}}$  of K3 surfaces with Néron–Severi lattice isometric to  $S_{1,1,2}$  is unirational.

### 3.5. The lattice $S_{1,1,3}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,1,3}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, A_2, A_3, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 3 & 2 & 0 \\ 3 & -2 & 0 & 2 \\ 2 & 0 & -2 & 6 \\ 0 & 2 & 6 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_2 = A_1 + A_2$$

is ample, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = 1$  and  $D_2 \cdot A_3 = D_2 \cdot A_4 = 2$ . We have  $2D_2 \equiv A_3 + A_4$ ; thus we have obtained the first part of the following proposition.

**Proposition 3.5.** *The surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a smooth sextic  $C_6$  which has a tritangent line and a 6-tangent conic. For general  $X$ , the sextic has equation*

$$C_6: \ell q g - f^2 = 0,$$

where  $\ell$ ,  $q$ ,  $g$  and  $f$  are forms of degree 1, 2, 3 and 3, respectively. The line  $\ell = 0$  is the tritangent to the sextic, and the curve  $q = 0$  is the 6-tangent conic. The moduli space  $\mathcal{M}_{S_{1,1,3}}$  of K3 surfaces  $X$  with  $\text{NS}(X) \simeq S_{1,1,3}$  is unirational.

*Proof.* Let  $C_6: \ell q g - f^2 = 0$  be a sextic curve as above, and let  $Y \rightarrow \mathbb{P}^2$  be the double cover branched over  $C_6$ . Above the line  $\ell = 0$  are two (-2)-curves  $A_1, A_2$  such that  $A_1 \cdot A_2 = 3$ , and above the conic  $q = 0$  are two (-2)-curves  $A_3, A_4$  such that  $A_3 \cdot A_4 = 6$ . It remains to understand the intersections  $A_j \cdot A_k$  for  $j \in \{1, 2\}$  and  $k \in \{3, 4\}$ . This is done by computing Example 3.6 below, for which the intersection is (up to permutation of  $A_1, A_2$  and  $A_3, A_4$ ) the above intersection matrix of the four (-2)-curves on a K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,1,3}$ . The intersection numbers remain the same for the (-2)-curves in that flat family of smooth surfaces.

If the equation of  $C_6$  is general, the Picard number of  $Y$  is 3. The lattice generated by the (-2)-curves is  $S_{1,1,3}$ , of discriminant 18. The unique over-lattice containing  $S_{1,1,3}$  is  $S_{1,1,1}$ , but a surface with Néron-Severi lattice isometric to  $S_{1,1,1}$  contains only three (-2)-curves; thus  $\text{NS}(Y) \simeq S_{1,1,3}$ .

In order to prove that a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,1,3}$  is branched over a sextic curve with an equation of the form  $\ell q g - f^2 = 0$ , let us consider the map

$$\Phi: H^0(\mathbb{P}^2, \mathcal{O}(1)) \oplus H^0(\mathbb{P}^2, \mathcal{O}(2)) \oplus H^0(\mathbb{P}^2, \mathcal{O}(3))^{\oplus 2} \rightarrow H^0(\mathbb{P}^2, \mathcal{O}(6))$$

defined by

$$w := (\ell, q, g, f) \mapsto f_{6,w} := \ell q g - f^2.$$

It is invariant under the action of the transformations  $\Gamma: (\ell, q, g, f) \mapsto (\alpha\ell, \beta q, \gamma g, f)$  for  $\alpha\beta\gamma = 1$ . Suppose

$$\ell q g - f^2 = \ell' q' g' - f'^2$$

for  $f' \neq \pm f$  and that the forms are chosen general so that the double cover  $Y$  branched over the sextic  $f_{6,w} = 0$  has  $\text{NS}(Y) \simeq S_{1,1,3}$ . Since the curves  $\ell = 0$ ,  $q = 0$  are the images of the four (-2)-curves on  $Y$ , up to rescaling by using a transformation  $\Gamma$ , one can suppose  $\ell' = \ell$ ,  $q' = q$ , and then one obtains the relation

$$\ell q (g - g') = (f - f')(f + f').$$

Up to changing the sign of  $f'$ , we can suppose that  $\ell$  divides  $f + f'$ . If  $q$  does not divide  $f + f'$ , then one gets a codimension 1 family of such sextic curves; thus we can suppose that there exists a scalar  $\alpha$  such that  $\ell q = \alpha(f + f')$ , and by solving the equations, we obtain that

$$f' = \frac{1}{\alpha} \ell q - f, \quad g' = g + \frac{1}{\alpha^2} \ell q - \frac{2}{\alpha} f.$$

The dimension of the (unirational) moduli space of sextic curves with an equation of the form  $\ell q g - f^2 = 0$  is

$$(3 + 6 + 2 \cdot 10) - (9 + 2 + 1) = 17;$$

since  $\mathcal{M}_{S_{1,1,3}}$  also has dimension 17, both spaces are birational.  $\square$

*Example 3.6.* Let us take

$$\begin{aligned} \ell &= 13x + 10y + z, & q &= 4x^2 + 6xy + 26xz + 6yz + 23z^2, \\ g &= 9x^3 + 16x^2y + 5xy^2 + 24y^3 + 22x^2z + 26xyz + 4y^2z + 23xz^2 + 20yz^2 + 9z^3, \\ f &= 28x^3 + 8x^2y + 2xy^2 + 23y^3 + 2x^2z + 23xyz + 19y^2z + 24xz^2 + 20yz^2 + 17z^3. \end{aligned}$$

Let  $X$  be the associated K3 surface, and let  $X_p$  be its reduction modulo a prime  $p$ . Using the Tate conjectures, one finds that the K3 surfaces  $X_{23}$  and  $X_{29}$  have Picard number 4. By the Artin-Tate conjectures, one computes that

$$\begin{aligned} |\text{Br}(X_{23})| \cdot |\text{disc}(\text{NS}(X_{23}))| &= 3^2 \cdot 491, \\ |\text{Br}(X_{29})| \cdot |\text{disc}(\text{NS}(X_{29}))| &= 3^3 \cdot 5^2 \cdot 53, \end{aligned}$$

and using Van Luijk's trick (see [vLu07]), we conclude that the Picard number of  $X$  is 3 since the ratio of the two integers above is not a square (here  $\text{Br}$  is the Brauer group, and  $\text{disc}$  denotes the discriminant group).



*Remark 3.7.* In [Rou19], we study K3 surfaces with finite automorphism group and compact fundamental domain. The surface with Néron–Severi lattice of type  $S_4$  (Nikulin’s notation) is also a double cover of  $\mathbb{P}^2$  branched over a smooth sextic which has one tritangent line and one 6-tangent conic. But the intersection matrix of the four  $(-2)$ -curves above the line and the conic (the only  $(-2)$  curves on that surface) is

$$\begin{pmatrix} -2 & 3 & 1 & 1 \\ 3 & -2 & 1 & 1 \\ 1 & 1 & -2 & 6 \\ 1 & 1 & 6 & -2 \end{pmatrix}.$$

*Remark 3.8.* By Section 2.8, we know that the unique moduli space  $\mathcal{M}_{S_{1,1,3}}$  of  $S_{1,1,3}$ -polarized K3 surfaces is irreducible. By Proposition 3.5, there is a copy of  $\mathcal{M}_{S_{1,1,3}}$  in the moduli space  $\mathcal{M}_{[2]}$  of K3 surfaces with an ample divisor of square 2. Using the Hilbert basis of the nef cone, one can check that  $D_2$  is the unique nef divisor of square 2 in the Néron–Severi group; therefore, the copy of  $\mathcal{M}_{S_{1,1,3}}$  in  $\mathcal{M}_{[2]}$  is unique.

### 3.6. The lattice $S_{1,1,4}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,1,4}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, A_2, A_3, A_4$ , with intersection matrix

$$M_1 = \begin{pmatrix} -2 & 4 & 0 & 2 \\ 4 & -2 & 2 & 0 \\ 0 & 2 & -2 & 4 \\ 2 & 0 & 4 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisors  $F_1 = A_1 + A_4$  and  $F_2 = A_2 + A_3$  are fibers of two distinct elliptic fibrations. The divisor

$$D_4 = A_1 + A_2 \equiv A_3 + A_4$$

is ample, of square 4, with  $D_4 \cdot A_j = 2$  for  $j \in \{1, \dots, 4\}$  and is non-hyperelliptic.

**Proposition 3.9.** *The surface  $X$  is a quartic in  $\mathbb{P}^3$  which has two hyperplane sections which are unions of two conics. The general surface has a projective model of the form*

$$X: \ell_1 \ell_2 q_3 - q_1 q_2 = 0 \hookrightarrow \mathbb{P}^3,$$

where  $\ell_1, \ell_2$  are linear forms and  $q_1, q_2, q_3$  are quadrics. The moduli space  $\mathcal{M}_{S_{1,1,4}}$  of K3 surfaces  $X$  with  $\text{NS}(X) \simeq S_{1,1,4}$  is unirational.

*Proof.* Consider the map

$$\Phi: H^0(\mathbb{P}^3, \mathcal{O}(1))^{\oplus 2} \oplus H^0(\mathbb{P}^3, \mathcal{O}(2))^{\oplus 3} \rightarrow H^0(\mathbb{P}^3, \mathcal{O}(4))$$

defined by

$$w := (\ell_1, \ell_2, q_1, q_2, q_3) \mapsto Q_{4,w} := \ell_1 \ell_2 q_1 - q_2 q_3.$$

By computing the differential  $d\Phi_w$  at a randomly chosen point, we find that it has rank 33; thus the image of  $\Phi$  is 33-dimensional, and the quotient  $W$  of that image by  $GL_4(\mathbb{C})$  is at least 17-dimensional. The space  $W$  is unirational. For a general  $w$ , the quartic  $Y: Q_{4,w} = 0$  is non-singular. The curves  $A_1: \ell_1 = q_2 = 0$ ,  $A_2: \ell_1 = q_3 = 0$ ,  $A_3: \ell_2 = q_2 = 0$ ,  $A_4: \ell_2 = q_3 = 0$  are  $(-2)$ -curves on  $Y$  with (up to permutation of  $A_3$  and  $A_4$ ) intersection matrix  $M_1$ . We thus obtain an injective map from  $W$  to the moduli space  $\mathcal{M}_{S_{1,1,4}}$ . Since  $\mathcal{M}_{S_{1,1,4}}$  has dimension 17, this moduli space is unirational.  $\square$

*Remark 3.10.* a) One can construct, more geometrically, a member of  $\mathcal{M}_{S_{1,1,4}}$  by considering two disjoint conics  $C_1$  and  $C_3$  in  $\mathbb{P}^3$  and taking a general quartic among the 16-dimensional linear system of quartics containing them. We also obtain in that way a dominant map from a rational space to  $\mathcal{M}_{S_{1,1,4}}$ .

b) In [Rou19], we study K3 surfaces with finite automorphism group and compact fundamental domain. One of these surfaces, namely the surface with Néron–Severi lattice of type  $S_3$  (Nikulin’s notation) is also a quartic surface in  $\mathbb{P}^3$  with two hyperplane sections which are the union of two conics. But the intersection matrix of the four  $(-2)$ -curves (which are the only  $(-2)$  curves on that surface) is

$$\begin{pmatrix} -2 & 4 & 1 & 1 \\ 4 & -2 & 1 & 1 \\ 1 & 1 & -2 & 4 \\ 1 & 1 & 4 & -2 \end{pmatrix}.$$

By using a construction as in part a), one finds that the moduli space of these surfaces is also unirational.

**Proposition 3.11.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,1,4}$  is trivial.*

*Proof.* The walls of the nef cone are  $A_k^\perp$ ,  $k \in \{1, \dots, 4\}$ . A Hilbert basis (see Definition 3.12) of the nef cone is given in [ACL21]; it is as follows. The divisors  $H_1 = A_1 + A_4$  and  $H_2 = A_2 + A_3$  are fibers of the two distinct elliptic fibrations on the K3 surface  $X$ ; the seven remaining classes  $H_3, \dots, H_9$  in the Hilbert basis are (in the basis  $A_1, A_2, A_3$ )

$$(1, 1, 0), (1, 1, 1), (1, 2, 0), (2, 1, 0), (2, 1, 1), (2, 2, -1), (2, 3, -1).$$

The intersection matrix  $M_H = (H_i \cdot H_j)_{1 \leq i, j \leq 9}$  is

$$M_H = \begin{pmatrix} 0 & 8 & 4 & 8 & 8 & 4 & 8 & 4 & 8 \\ 8 & 0 & 4 & 4 & 4 & 8 & 8 & 8 & 8 \\ 4 & 4 & 4 & 6 & 6 & 6 & 8 & 6 & 8 \\ 8 & 4 & 6 & 6 & 10 & 8 & 8 & 12 & 16 \\ 8 & 4 & 6 & 10 & 6 & 12 & 16 & 8 & 8 \\ 4 & 8 & 6 & 8 & 12 & 6 & 8 & 10 & 16 \\ 8 & 8 & 8 & 8 & 16 & 8 & 8 & 16 & 24 \\ 4 & 8 & 6 & 12 & 8 & 10 & 16 & 6 & 8 \\ 8 & 8 & 8 & 16 & 8 & 16 & 24 & 8 & 8 \end{pmatrix}.$$

Let us check the condition of Proposition 2.11:

(i) Since the intersection matrix of the fibers  $H_1, H_2$  with  $A_1, \dots, A_4$  is

$$\begin{pmatrix} 0 & 4 & 4 & 0 \\ 4 & 0 & 0 & 4 \end{pmatrix},$$

there are no  $(-2)$ -curves  $A$  and fibers  $F$  such that  $AF = 1$ .

(ii) A nef divisor  $D$  is a positive linear combination of the elements  $H_1, \dots, H_9$  of the Hilbert basis of the nef cone. From the two first lines of the Gram matrix  $M_H$ , we see that there is no big and nef divisor  $D$  such that  $D \cdot F = 2$  for any fiber  $F = H_1$  or  $H_2$ .

(iii) From the matrix  $M_H$ , we see that there is no big and nef divisor  $D$  such that  $D^2 = 2$ .

Therefore, one can apply Proposition 2.11 and conclude that the automorphism group of  $X$  is trivial.  $\square$

**Definition 3.12.** Let us recall that if  $C \subset \mathbb{R}^d$  is a polyhedral cone generated by integral vectors, a Hilbert basis  $H(C)$  of  $C$  is a subset of integral vectors such that

- a) each element of  $C \cap \mathbb{Z}^d$  can be written as a non-negative integer combination of elements of  $H(C)$ , and
- b)  $H(C)$  has minimal cardinality with respect to all subsets of  $C \cap \mathbb{Z}^d$  for which part a) holds.

### 3.7. The lattice $S_{1,1,6}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,1,6}$ . The surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$ , with intersection matrix

$$\begin{pmatrix} -2 & 6 & 2 & 6 & 6 & 2 \\ 6 & -2 & 6 & 2 & 2 & 6 \\ 2 & 6 & -2 & 18 & 0 & 16 \\ 6 & 2 & 18 & -2 & 16 & 0 \\ 6 & 2 & 0 & 16 & -2 & 18 \\ 2 & 6 & 16 & 0 & 18 & -2 \end{pmatrix}.$$

The curves  $A_1, A_3, A_5$  generate the Néron–Severi lattice. We have

$$A_2 = A_1 - 2A_3 + 2A_5, \quad A_4 = 4A_1 - 5A_3 + 4A_5, \quad A_6 = 4A_1 - 4A_3 + 3A_5.$$

The divisor

$$D_2 = A_1 - A_3 + A_5$$

is ample, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = 2$  and  $D_2 \cdot A_j = 4$  for  $j \in \{3, \dots, 6\}$ . We have

$$2D_2 \equiv A_1 + A_2, \quad 4D_2 \equiv A_3 + A_4 \equiv A_5 + A_6.$$

Therefore, the following holds.

**Proposition 3.13.** *The K3 surface  $X$  is the double cover  $\eta: X \rightarrow \mathbb{P}^2$  of  $\mathbb{P}^2$  branched over a smooth sextic curve which has a 6-tangent conic  $C$ , and there are two rational cuspidal quartics  $Q_1, Q_2$  such that their three cusps are on the sextic curve and their remaining intersection points are tangent to the sextic (there are nine such points). The image by  $\eta$  of  $A_1 + A_2$  is  $C$ , the image of  $A_3 + A_4$  is  $Q_1$ , and the image of  $A_5 + A_6$  is  $Q_2$ .*

### 3.8. The lattice $S_{1,1,8}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,1,8}$ . The surface  $X$  contains eight  $(-2)$ -curves  $A_1, \dots, A_8$ , with intersection matrix

$$\begin{pmatrix} -2 & 2 & 0 & 8 & 16 & 8 & 18 & 14 \\ 2 & -2 & 8 & 0 & 8 & 16 & 14 & 18 \\ 0 & 8 & -2 & 14 & 18 & 2 & 16 & 8 \\ 8 & 0 & 14 & -2 & 2 & 18 & 8 & 16 \\ 16 & 8 & 18 & 2 & -2 & 14 & 0 & 8 \\ 8 & 16 & 2 & 18 & 14 & -2 & 8 & 0 \\ 18 & 14 & 16 & 8 & 0 & 8 & -2 & 2 \\ 14 & 18 & 8 & 16 & 8 & 0 & 2 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. We have

$$\begin{aligned} A_4 &= -2A_1 + 2A_2 + A_3, & A_5 &= -5A_1 + 3A_2 + 3A_3, & A_6 &= -3A_1 + A_2 + 3A_3, \\ A_7 &= -6A_1 + 3A_2 + 4A_3, & A_8 &= -5A_1 + 2A_2 + 4A_3. \end{aligned}$$

The divisor

$$D_6 = -A_1 + A_2 + A_3$$

is ample, of square 6, base-point free and non-hyperelliptic, with  $D_6 \cdot A_1 = D_6 \cdot A_2 = 4$ ,  $D_6 \cdot A_3 = D_6 \cdot A_4 = 6$ ,  $D_6 \cdot A_5 = D_6 \cdot A_6 = 10$ ,  $D_6 \cdot A_7 = D_6 \cdot A_8 = 12$ . The K3 surface  $X$  is a smooth complete intersection in  $\mathbb{P}^4$ . We remark that

$$A_3 + A_4 = 2D_6.$$

**Proposition 3.14.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,1,4}$  is trivial.*


*Proof.* We proceed as in the proof of Proposition 3.11. The Hilbert basis of the nef cone of the K3 surface  $X$  is a set of 60 classes (see [ACL21, Table 1]), among which are the fibers

$$A_2 + A_3, \quad 4A_1 + 5A_2 - 7A_3, \quad 4A_1 + A_2 - 3A_3, \quad 8A_1 + 5A_2 - 11A_3.$$

Then one can check that any big and nef divisor  $D$  on  $X$  is base-point free, with  $D^2 \geq 6$  or  $D^2 = 0$ . Moreover, there is no fiber  $F$  of an elliptic fibration such that  $FD = 2$ ; thus by Theorems 2.2 and 2.3, there are no hyperelliptic involutions acting on  $X$ . From [Kon89, Table 1 and Lemma 2.3], the automorphism group of  $X$  is therefore trivial.  $\square$

### 3.9. The lattice $S_{1,2,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,2,1}$ . The surface  $X$  contains three  $(-2)$ -curves  $A_1, \dots, A_3$ , with dual graph



These curves generate the Néron–Severi lattice. The divisor

$$D_6 = 2A_1 + 2A_2 + A_3$$

is ample, of square 6, with  $D_6 \cdot A_1 = D_6 \cdot A_2 = 1$  and  $D_6 \cdot A_3 = 2$ . Since  $D_6 \cdot (A_1 + A_2) = 2$  and  $A_1 + A_2$  is a fiber,  $D_6$  is hyperelliptic. The divisor

$$D_2 = A_1 + A_2 + A_3$$


is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = 1$  and  $D_2 \cdot A_3 = 0$ .

**Proposition 3.15.** *The K3 surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  which has a node at a point  $q$  and a line  $L$  containing the node, which is bitangent to  $C_6$  at two points  $p_1, p_2$ . The moduli space  $\mathcal{M}_{1,2,1}$  is unirational.*

*Proof.* It remains to prove the assertion on the moduli space  $\mathcal{M}_{1,2,1}$ . We identify this space with the moduli space of sextic curves with a node and a line through the node which is bitangent to the sextic. The imposition of a node and the tangency conditions are linear conditions on the space  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(6))$  of sextic curves; thus the moduli space of such curves is unirational.  $\square$

### 3.10. The lattice $S_{1,3,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,3,1}$ . The surface  $X$  contains three  $(-2)$ -curves  $A_1, A_2, A_3$ , with dual graph



These curves generate the Néron–Severi lattice. The divisor

$$D_4 = A_1 + A_2 + A_3$$

is very ample of square 4, with  $D_4 \cdot A_1 = D_4 \cdot A_2 = 1$  and  $D_4 \cdot A_3 = 2$ .

**Proposition 3.16.** *The K3 surface  $X$  is a quartic surface in  $\mathbb{P}^3$  with a hyperplane section which is the union of two lines and a conic. The general surface  $X$  with  $\text{NS}(X) \simeq S_{1,3,1}$  has an equation of the form*

$$\ell_1 f + q_1 \ell_2 \ell_3 = 0.$$

*The moduli space  $\mathcal{M}_{S_{1,3,1}}$  of such surfaces is unirational.*

*Proof.* Let  $Y: \ell_1 f + q_1 \ell_2 \ell_3 = 0$  be a quartic surface as in the proposition. The hyperplane section  $\ell_1$  cuts  $Y$  into a union of two lines  $A_1: \ell_1 = \ell_2 = 0$ ,  $A_2: \ell_1 = \ell_3 = 0$  and a conic  $A_3: \ell_1 = q_1 = 0$ . Their intersection graph is as above.

If the equation of  $Y$  is general, the Picard number of  $Y$  is 3; the lattice generated by the  $(-2)$ -curves is  $S_{1,3,1}$ , of discriminant 18. The unique over-lattice containing  $S_{1,3,1}$  is  $S_{1,1,1}$ , but the configuration of the  $(-2)$ -curves in a surface with Néron–Severi lattice isometric to  $S_{1,1,1}$  is different; thus  $\text{NS}(Y) \simeq S_{1,3,1}$ .

Let us consider the map

$$\Phi: H^0(\mathbb{P}^3, \mathcal{O}(1))^{\oplus 3} \oplus H^0(\mathbb{P}^3, \mathcal{O}(2)) \oplus H^0(\mathbb{P}^3, \mathcal{O}(3)) \rightarrow H^0(\mathbb{P}^3, \mathcal{O}(4))$$

defined by

$$w := (\ell_1, \ell_2, \ell_3, q_1, f) \mapsto Q_{4,w} := \ell_1 f + q_1 \ell_2 \ell_3.$$

It is invariant under the action of the transformations  $\Gamma: (\ell_1, \ell_2, \ell_3, q_1, f) \mapsto (\nu \ell_1, \alpha \ell_2, \beta \ell_3, \gamma q_1, \mu f)$  for  $\nu \mu = 1$  and  $\alpha \beta \gamma = 1$ . Suppose

$$(3.2) \quad \ell_1 f + q_1 \ell_2 \ell_3 = \ell'_1 f' + q'_1 \ell'_2 \ell'_3$$

and that the forms are chosen general so that the associated quartic satisfies  $\text{NS}(Y) \simeq S_{1,1,3}$ . Since the lines  $A_1: \ell_1 = \ell_2 = 0$ ,  $A_2: \ell_1 = \ell_3 = 0$  and the conic  $A_3: \ell_1 = q_1 = 0$  are the only  $(-2)$ -curves on  $X$ , we have (up to exchanging  $\ell'_2, \ell'_3$ )

$$A_1: \ell'_1 = \ell'_2 = 0, \quad A_2: \ell'_1 = \ell'_3 = 0, \quad A_3: \ell'_1 = q'_1 = 0.$$

The consequence is that—up to rescaling using a transformation  $\Gamma$ —one can suppose that there exist scalars  $\alpha_1, \alpha_2, \alpha_3$  and a linear form  $\ell$  such that

$$(3.3) \quad \ell'_1 = \ell_1, \quad \ell'_2 = \alpha_2 \ell_1 + \ell_2, \quad \ell'_3 = \alpha_3 \ell_1 + \ell_2, \quad q'_1 = \alpha_1 q_1 + \ell_1 \ell.$$

Substituting this into the formula (3.2) and taking the equation modulo  $\ell_1$ , one finds that necessarily  $\alpha_1 = 1$ . Conversely, given forms as in equation (3.3), with  $\alpha_1 = 1$ , one can find a cubic form  $f'$  such that the linear forms  $\ell'_1, \ell'_2, \ell'_3$ , the quadric  $q'_1$  and the cubic  $f'$  satisfy equation (3.2). Therefore the dimension of the (unirational) moduli space  $\mathcal{M}_4$  of quartic surfaces with an equation of type  $\ell_1 f + q_1 \ell_2 \ell_3 = 0$  is

$$(3 \cdot 4 + 10 + 20) - (16 + 1 + 2 + 6) = 17.$$

An open set of  $\mathcal{M}_4$  is a subspace of the moduli space  $\mathcal{M}_{S_{1,3,1}}$ , which also has dimension 17; thus both spaces are birational.  $\square$

**Proposition 3.17.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,3,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11. The Hilbert basis of the nef cone of the K3 surface  $X$  is a set of four classes:

$$A_2 + A_3, \quad A_1 + A_3, \quad A_1 + A_2 + A_3, \quad 2A_1 + 2A_2 + A_1;$$

the first two classes are fibers of elliptic fibrations. The intersection matrix of these four classes is

$$\begin{pmatrix} 0 & 3 & 3 & 6 \\ 3 & 0 & 3 & 6 \\ 3 & 3 & 4 & 6 \\ 6 & 6 & 6 & 4 \end{pmatrix}.$$

From that one can check that any big and nef divisor  $D$  on  $X$  is base-point free, with  $D^2 \geq 4$  or  $D^2 = 0$ , and there is no fiber  $F$  of an elliptic fibration such that  $FD = 2$ . Thus we conclude as in Proposition 3.11 that the automorphism group of  $X$  is trivial.  $\square$

### 3.11. The lattice $S_{1,4,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,4,1}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, A_2, A_3, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 3 & 2 & 1 \\ 3 & -2 & 1 & 2 \\ 2 & 1 & -2 & 11 \\ 1 & 2 & 11 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_2 = A_1 + A_2$$

is ample, of square 2. We have

$$A_3 + A_4 \equiv 3D_2;$$

therefore, using the linear system  $|D_2|$ , we obtain the following.

**Proposition 3.18.** *The K3 surface  $X$  is the double cover  $\eta: X \rightarrow \mathbb{P}^2$  of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$  which has a tritangent line and a cuspidal cubic with a cusp on  $C_6$  and such that the cuspidal cubic and  $C_6$  are tangent at every other intersection points.*

The divisor  $D_6 = A_1 + A_2 + A_3$  is very ample of square 6, with  $D_6 \cdot A_j = 3, 2, 1, 14$  for  $j = 1, \dots, 4$ , so that  $A_1, A_2, A_3$  are, respectively, a rational cubic, a conic and a line in a hyperplane of  $\mathbb{P}^4$ . That leads us to the following proposition.

**Proposition 3.19.** *The moduli space  $\mathcal{M}_{S_{1,4,1}}$  of K3 surfaces with  $\text{NS}(X) \simeq S_{1,4,1}$  is unirational.*

*Proof.* We can construct a member of  $\mathcal{M}_{S_{1,4,1}}$  as follows: Let  $H_3 \subset \mathbb{P}^4$  be a hyperplane, let  $B_3 \subset H_3$  be a line, let  $p_0, p_1, p_3$  be three point on  $B_3$ , let  $H_2 \subset H_3$  be a plane intersecting  $B_3$  at  $p_0$ , let  $B_2 \subset H_2$  be a smooth conic containing  $p_0$ , and let  $q_1, q_2, q_3$  be three general points on  $B_2$  (thus different from  $p_0$ ). The moduli space of normal cubic curves passing through the points  $p_1, p_2, q_1, q_2, q_3$  is unirational. Let  $B_1 \subset H_3$  be such a rational normal cubic curve. By construction, for  $j \neq k$  in  $\{1, 2, 3\}$ , the degree of the intersection scheme of  $B_j$  and  $B_k$  equals  $A_j \cdot A_k$ .

There is a unique quadric in  $H_3$  that contains the curves  $B_1, B_2, B_3$ . The linear system of cubics in  $H_3$  that contain these curves is 3-dimensional. Therefore, the linear systems  $L_2, L_3$  of quadrics and cubics in  $\mathbb{P}^4$  containing the three curves  $B_1, B_2, B_3$  are, respectively, 4- and 18-dimensional.

Let  $X \hookrightarrow \mathbb{P}^4$  be the degree 6 K3 surface which is the complete intersection of a general quadric in  $L_2$  and a general cubic in  $L_3$ . It has Picard number at least 3, and there is an open subspace for which the Picard number is 3 since K3 surfaces with Néron–Severi lattice  $S_{1,4,1}$  belong to that space. Thus for a general choice, the Néron–Severi lattice of  $X$  has rank 3 and contains the lattice  $S_{1,4,1}$  generated by the  $(-2)$ -curves  $B_1, B_2, B_3$ . The only over-lattices of  $S_{1,4,1}$  are  $S_{1,1,1}$  and  $S_{1,2,1}$ . The K3 surfaces with such a Néron–Severi lattice contain only three  $(-2)$ -curves which do not have the same intersection matrix as the curves  $B_k$ ,  $k = 1, \dots, 3$ . Thus  $\text{NS}(X) \simeq S_{1,4,1}$ , and by the above construction, the moduli space  $\mathcal{M}_{S_{1,4,1}}$  is unirational.  $\square$

*Remark 3.20.* a) We recall that a rational normal curve is a smooth, rational curve of degree  $n$  in projective  $n$ -space  $\mathbb{P}^n$ . Given  $n + 3$  points in  $\mathbb{P}^n$  in linear general position (that is, with no  $n + 1$  lying in a hyperplane), there is a unique rational normal curve  $C$  passing through them. From, e.g., [Har92, Theorem 1.18], the coefficients of the equations of that curve are rational functions in the coordinates of the  $n + 3$  points. In the above proof, for the unirationality of  $\mathcal{M}_{S_{1,4,1}}$ , we implicitly used the fact that the construction of a rational cubic curve passing through the four points in general position is rational in the coordinates of the points.

b) There are  $\binom{n+2}{2} - 2n - 1$  independent quadrics that generate the ideal of a degree  $n$  rational normal curve in  $\mathbb{P}^n$ .

### 3.12. The lattice $S_{1,5,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,5,1}$ . The surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$ , with intersection matrix

$$\begin{pmatrix} -2 & 6 & 4 & 2 & 2 & 14 \\ 6 & -2 & 2 & 4 & 14 & 2 \\ 4 & 2 & -2 & 11 & 1 & 23 \\ 2 & 4 & 11 & -2 & 23 & 1 \\ 2 & 14 & 1 & 23 & -2 & 66 \\ 14 & 2 & 23 & 1 & 66 & -2 \end{pmatrix}.$$

The curves  $A_1, A_3, A_5$  generate the Néron–Severi lattice. The divisor

$$D_2 = 2A_1 + 2A_3 - A_5$$

is ample, of square 2, with

$$2D_2 \equiv A_1 + A_2, \quad 3D_2 \equiv A_3 + A_4, \quad 8D_2 \equiv A_5 + A_6.$$

Thus, by using the linear system  $|D_2|$ , we obtain the following.

**Proposition 3.21.** *The surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$  which has a 6-tangent conic, a tangent cuspidal cubic and a tangent rational cuspidal octic such that the cusps are on  $C_6$ , and at each intersection point of the octic or the cubic with  $C_6$ , the multiplicity is even.*

The divisor  $D_4 = A_1 + A_3$  is very ample of square 4, with  $D_4 \cdot A_j = 2, 8, 2, 13, 3, 37$  for  $j = 1, \dots, 6$ . The divisor  $D_8 = A_1 + A_3 + A_5$  is very ample of square 8, with  $D_8 \cdot A_j = 4, 22, 3, 36, 1, 103$  for  $j = 1, \dots, 6$ . This last model enables us to construct the surfaces with  $\text{NS}(X) \simeq S_{1,5,1}$  and to obtain the following proposition.

**Proposition 3.22.** *The moduli space  $\mathcal{M}_{S_{1,5,1}}$  is unirational.*

*Proof.* Let us fix a hyperplane  $H_4 \subset \mathbb{P}^5$ , and let  $B_3 \hookrightarrow H_4$  be a normal quartic curve. Let  $H_3$  be a general hyperplane of  $H_4$ , let  $q_1, \dots, q_4$  be the intersection points of  $B_3$  with  $H_3$ . Let  $B_5 \hookrightarrow H_4$  be a line passing through two general points  $p_1, p_2$  of  $B_3$ , and let  $q_0$  the intersection point of  $B_5$  with  $H_3$ . Let  $B_1 \hookrightarrow H_3$  be a rational normal cubic curve containing the points  $q_0, q_1, \dots, q_4$ .

By construction, the curves  $B_1, B_3, B_5$  are such that the degree of the 0-cycle  $B_j \cdot B_k$  equals  $A_j \cdot A_k$  for  $j \neq k$  in  $\{1, 3, 5\}$ . The linear system of quadrics in  $H_4$  containing the curves  $B_1, B_2, B_5$  is a net (a 2-dimensional linear space); thus the linear system  $\mathcal{L}$  of quadrics in  $\mathbb{P}^5$  containing these curves is 8-dimensional. A general net of quadrics in  $\mathcal{L}$  defines a smooth K3 surface  $X$  such that  $X \cdot H_4 = B_1 + B_3 + B_5$ , and the curves  $B_k$  are  $(-2)$ -curves on  $X$ . The curves  $B_1, B_3, B_5$  generate a lattice isometric to  $S_{1,5,1}$ , and for a general choice,  $\text{NS}(X) \simeq S_{1,5,1}$ . That construction and Remark 3.20 on the parametrization of rational normal curves show that the moduli space  $\mathcal{M}_{S_{1,5,1}}$  is unirational.  $\square$

### 3.13. The lattice $S_{1,6,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,6,1}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, \dots, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 5 & 2 & 1 \\ 5 & -2 & 1 & 2 \\ 2 & 1 & -2 & 5 \\ 1 & 2 & 5 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_6 = A_1 + A_2 \equiv A_3 + A_4$$

is ample, of square 6, with  $D_6 \cdot A_j = 3$  for  $j \in \{1, \dots, 4\}$ . It is base-point free and non-hyperelliptic, and therefore the surface  $X$  is a degree 6 surface in  $\mathbb{P}^4$  with two hyperplane sections that split as the union of two rational normal cubic curves.

**Proposition 3.23.** *The linear system  $|D_6|$  gives an embedding of  $X$  as a complete intersection in  $\mathbb{P}^4$  with two hyperplanes sections which split as the unions of two rational cubic curves. The moduli space  $\mathcal{M}_{S_{1,6,1}}$  of K3 surfaces  $X$  with  $\text{NS}(X) \simeq S_{1,6,1}$  is unirational.*

*Proof.* One can construct these surfaces by taking two degree 3 rational normal curves  $C_1, C_4$  in two different hyperplanes  $H_1, H_2$  such that the curves  $C_1, C_4$  meet transversely in one point. The linear system  $\mathcal{Q}$  of quadrics containing  $C_1$  and  $C_4$  is a pencil, and the linear system  $\mathcal{C}$  of cubics containing  $C_1$  and  $C_4$  has dimension 15. Let  $X$  be the intersection of a general element in  $\mathcal{Q}$  and a general element in  $\mathcal{C}$ . The intersections of  $X$  with  $H_1, H_2$  break down into  $C_1 + C_2$  and  $C_3 + C_4$ , where  $C_2, C_3$  are two rational cubic normal curves. Using that  $C_1 + C_2, C_3 + C_4$  are hyperplane sections and that we know that  $C_1 C_4 = 1$ , one obtains that the curves  $C_1, \dots, C_4$  have the above intersection matrix and therefore generate a lattice isometric to  $S_{1,6,1}$ , which is equal to the Néron–Severi lattice for a general choice of  $X$ . The construction shows that the moduli space  $\mathcal{M}_{S_{1,6,1}}$  is unirational.  $\square$

**Proposition 3.24.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,6,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11. The Hilbert basis of the nef cone of the K3 surface  $X$  contains 17 classes (see [ACL21, Table 1]). One can check that any big and nef divisor  $D$  on  $X$  is base-point free, with  $D^2 \geq 6$  or  $D^2 = 0$ ; moreover, there is no fiber  $F$  of an elliptic fibration such that  $FD = 2$ . Thus by Theorems 2.2 and 2.3, there are no hyperelliptic involutions acting on  $X$ . From [Kon89, Table 1 and Lemma 2.3], the automorphism group of  $X$  is therefore trivial.  $\square$

### 3.14. The lattice $S_{1,9,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{1,9,1}$ . The surface  $X$  contains nine  $(-2)$ -curves  $A_1, \dots, A_9$ , with intersection matrix

$$\begin{pmatrix} -2 & 10 & 2 & 8 & 10 & 26 & 2 & 26 & 8 \\ 10 & -2 & 8 & 2 & 10 & 2 & 26 & 8 & 26 \\ 2 & 8 & -2 & 1 & 26 & 37 & 25 & 46 & 37 \\ 8 & 2 & 1 & -2 & 26 & 25 & 37 & 37 & 46 \\ 10 & 10 & 26 & 26 & -2 & 8 & 8 & 2 & 2 \\ 26 & 2 & 37 & 25 & 8 & -2 & 46 & 1 & 37 \\ 2 & 26 & 25 & 37 & 8 & 46 & -2 & 37 & 1 \\ 26 & 8 & 46 & 37 & 2 & 1 & 37 & -2 & 25 \\ 8 & 26 & 37 & 46 & 2 & 37 & 1 & 25 & -2 \end{pmatrix}.$$

The family  $(A_1, A_3, A_4)$  is a basis of the Néron–Severi lattice. In that basis, the coordinates of the other  $(-2)$ -curves are

$$\begin{aligned} A_2 &= (1, -2, 2), & A_5 &= (5, -6, 4), & A_6 &= (6, -9, 7), \\ A_7 &= (6, -5, 3), & A_8 &= (8, -11, 8), & A_9 &= (8, -8, 5). \end{aligned}$$

The divisor

$$D_4 = A_1 - A_3 + A_4$$

is ample, of square 4, and the degrees of the curves  $A_1, \dots, A_9$  are

$$4, 4, 5, 5, 10, 14, 14, 17, 17.$$

The divisor  $D_4$  is very ample, and one has

$$2D_4 \equiv A_1 + A_2.$$




The divisor  $D_{10} \equiv 2A_1 - A_3 + A_4$  is very ample of square 10.

**Proposition 3.25.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{1,9,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11. The Hilbert basis of the nef cone of the K3 surface  $X$  is a set of 72 classes (see [ACL21, Table 1]).  $\square$

### 3.15. The lattice $S_{4,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{4,1,1}$ . The surface  $X$  contains three  $(-2)$ -curves  $A_1, A_2, A_3$ , with dual graph



These curves generate the Néron–Severi lattice. The divisor

$$D_6 = A_1 + A_2 + A_3$$

is ample, base-point free, non-hyperelliptic, of square 6, with  $D_6 \cdot A_j = 2$  for  $j \in \{1, 2, 3\}$ , so that the surface  $X$  is a degree 6 surface in  $\mathbb{P}^4$  with a hyperplane section containing three conics. That leads to the following.

**Proposition 3.26.** *The K3 surface  $X$  is degree 6 complete intersection in  $\mathbb{P}^4$  with a hyperplane section which is the union of three conics. The moduli space  $\mathcal{M}_{S_{4,1,1}}$  is unirational.*

*Proof.* Let  $P^3$  be a hyperplane in  $\mathbb{P}^4$ , and let  $P_1, P_2, P_3$  be three planes in  $P^3$ . For  $\{i, j, k\} = \{1, 2, 3\}$ , let us denote by  $L_k$  the line  $P_i \cap P_j$ . For each line  $L_k$ , let us choose two general points  $p_k \neq q_k$  on  $L_k$ . For  $\{i, j, k\} = \{1, 2, 3\}$ , let  $C_k$  be a smooth conic in  $P_k$  passing through  $p_i, p_j, q_i, q_j$ . The linear system of quadrics in  $P^3$  containing the conics  $C_1, C_2, C_3$  is 0-dimensional, and the linear system of cubics containing  $C_1, C_2, C_3$  is 4-dimensional. Thus the linear system of quadrics (resp. cubics) in  $\mathbb{P}^4$  containing the three conics has dimension 5 (resp. 19). A general complete intersection of such a quadric and cubic is a K3 surface with  $\text{NS}(X) \simeq S_{4,1,1}$ . By that construction, we see that the moduli space  $\mathcal{M}_{S_{4,1,1}}$  is unirational.  $\square$

By [ACL21, Theorem 3.9], the surface  $X$  has equations of the form

$$X: q_2 = \ell_1 \ell_2 \ell_3 + \ell_4 g_2 = 0,$$

where  $q_2, g_2$  are quadrics and  $\ell_1, \ell_2, \ell_3, \ell_4$  are independent linear forms.

**Proposition 3.27.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{4,1,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11. The Hilbert basis of the nef cone of the K3 surface  $X$  is (see [ACL21, Table 1])

$$A_2 + A_3, A_1 + A_3, A_1 + A_2, A_1 + A_2 + A_3.$$

Their intersection matrix is

$$\begin{pmatrix} 0 & 4 & 4 & 4 \\ 4 & 0 & 4 & 4 \\ 4 & 4 & 0 & 4 \\ 4 & 4 & 4 & 6 \end{pmatrix};$$

the result follows.  $\square$

### 3.16. The lattice $S_{5,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{5,1,1}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, \dots, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 3 & 2 & 2 \\ 3 & -2 & 2 & 2 \\ 2 & 2 & -2 & 18 \\ 2 & 2 & 18 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_2 = A_1 + A_2$$

is ample, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = 1$  and  $D_2 \cdot A_3 = D_2 \cdot A_4 = 4$ . We have

$$4D_2 \equiv A_3 + A_4;$$

thus, by using the linear system  $|D_2|$ , we obtain the following.

**Proposition 3.28.** *The surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$  which has a tritangent line and such that there is a rational cuspidal quartic curve  $Q_4$  such that its three cusps are on  $C_6$  and the intersection points of  $Q_4$  and  $C_6$  have even multiplicities. The moduli space  $\mathcal{M}_{S_{5,1,1}}$  of K3 surfaces  $X$  with  $\text{NS}(X) \simeq S_{5,1,1}$  is unirational.*

*Proof.* The divisor  $D_8 = A_1 + A_2 + A_3$  is very ample of square 8, with  $D_8 \cdot A_j = 3, 3, 2, 22$ . Thus the curves  $A_1, A_2, A_3$  are, respectively, two rational normal cubics and a conic. The rational normal cubics cut each others in three points and cut the conic in two points.

Let  $P_4 \subset \mathbb{P}^5$  be a hyperplane, and let  $P_3, P'_3$  be 3-dimensional projective subspaces of  $H_4$ . Let  $P_0$  be the plane  $P_0 = P_3 \cap P'_3$ . Let  $P_2 \subset P_4$  be a plane such that  $P_0$  and  $P_2$  meet at one point only. Let us define the lines  $L_1 = P_2 \cap P_3, L'_1 = P_2 \cap P'_3$ , and let us fix two points  $p_1, p_2$  (resp.  $p'_1, p'_2$ ) on  $L_1$  (resp.  $L'_1$ ) and three points  $q_1, q_2, q_3$  in  $P_0$ . We now fix a smooth rational normal cubic curve  $B_1$  (resp.  $B_2$ ) on  $P_3$  (resp.  $P'_3$ ) passing through  $p_1, p_2$  (resp.  $p'_1, p'_2$ ) and  $q_1, q_2, q_3$ . We also fix an irreducible conic  $B_3$  in  $P_2$  passing through  $p_1, p_2, p'_1, p'_2$ , so that

$$\text{Degree}(B_j \cap B_k) = A_j \cdot A_k$$

for  $j \neq k$  in  $\{1, 2, 3\}$ . The linear system  $\mathcal{Q}$  of quadrics containing the curve  $B_1 + B_2 + B_3$  is 11-dimensional; the complete intersection surface obtained by the intersection of the quadrics in a general net of  $\mathcal{Q}$  is a K3 surface  $X$  containing the  $(-2)$ -curves  $B_1, B_2, B_3$  which generate the lattice  $S_{5,1,1}$ . In fact, one has  $\text{NS}(X) \simeq S_{5,1,1}$  since the only over-lattice containing  $S_{5,1,1}$  is  $S_{1,1,1}$ . From that construction and Remark 3.20 on the construction of rational normal curves, we obtain that the moduli space  $\mathcal{M}_{S_{5,1,1}}$  is unirational.  $\square$

### 3.17. The lattice $S_{6,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{6,1,1}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, \dots, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 4 & 2 & 2 \\ 4 & -2 & 2 & 2 \\ 2 & 2 & -2 & 10 \\ 2 & 2 & 10 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_4 = A_1 + A_2$$

is ample, of square 4, with  $D_4 \cdot A_1 = D_4 \cdot A_2 = 2$  and  $D_4 \cdot A_3 = D_4 \cdot A_4 = 4$ . We have  $2D_4 = A_3 + A_4$ . The linear system  $|D_4|$  is base-point free, non-hyperelliptic. This leads to the following.

**Proposition 3.29.** *The surface  $X$  is a quartic in  $\mathbb{P}^3$  with a hyperplane section which is the union of two conics and a quadric section which is the union of two degree 4 smooth rational curves. The moduli space  $\mathcal{M}_{S_{6,1,1}}$  of K3 surfaces  $X$  with  $\text{NS}(X) \simeq S_{6,1,1}$  is unirational.*

*Proof.* Let us construct these K3 surfaces. Let  $A_3$  be a smooth degree 4 rational curve in  $\mathbb{P}^3$ , let  $p_1, p_2$  be two points on it, let  $P$  be a general plane containing the points  $p_1, p_2$ , and let  $A_1$  be an irreducible conic contained in  $P$  and containing the points  $p_1, p_2$ , so that the degree of the intersection  $A_1 \cdot A_3$  is 2. The linear system of quartics containing  $A_1$  and  $A_3$  is 10-dimensional. Let  $X$  be such a general quartic; the intersection of  $X$  and  $P$  contains  $A_1$ ; the residual curve  $A_2$  is another smooth conic. By [Har77, Exercise IV.6.1], there exists a unique quadric  $Q_2$  (which is moreover smooth) containing the curve  $A_3$ . The intersection of  $X$  and  $Q_2$  is the union of  $A_3$  and another degree 4 smooth rational curve  $A_4$ . Since

$$8 = 2HA_3 = (A_3 + A_4)A_3,$$

we get  $A_3 \cdot A_4 = 10$ , and therefore from  $(A_3 + A_4)^2 = 16$ , one obtains  $A_4^2 = -2$ . From the construction, the curves  $A_1, \dots, A_4$  have the above intersection matrix; thus the Néron-Severi lattice of  $X$  contains the lattice  $S_{6,1,1}$ , and by the general assumption  $\text{NS}(X) \simeq S_{6,1,1}$ . That construction shows that the moduli space  $\mathcal{M}_{S_{6,1,1}}$  is unirational.  $\square$

**Proposition 3.30.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{6,1,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11.  $\square$

### 3.18. The lattice $S_{7,1,1}$

Let  $X$  be a K3 surface with Néron-Severi lattice of type  $S_{7,1,1}$ . The surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$ , with intersection matrix

$$\begin{pmatrix} -2 & 5 & 2 & 5 & 16 & 2 \\ 5 & -2 & 2 & 5 & 2 & 16 \\ 2 & 2 & -2 & 16 & 26 & 26 \\ 5 & 5 & 16 & -2 & 2 & 2 \\ 16 & 2 & 26 & 2 & -2 & 26 \\ 2 & 16 & 26 & 2 & 26 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron-Severi lattice, and

$$A_4 \equiv 3A_1 + 3A_2 - 2A_3, \quad A_5 \equiv 4A_1 + 6A_2 - 3A_3, \quad A_6 \equiv 6A_1 + 4A_2 - 3A_3.$$

The divisor

$$D_6 = A_1 + A_2$$

is very ample of square 6. For  $j \in \{1, \dots, 6\}$ , we have  $D_6 \cdot A_j$  equal to, respectively, 3, 3, 4, 10, 18. The divisors

$$2A_1 + 2A_2 - A_3, \quad 3A_1 + 4A_2 - 2A_3, \quad 4A_1 + 3A_2 - 2A_3$$

are also very ample of square 6.

**Proposition 3.31.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{6,1,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11.  $\square$

### 3.19. The lattice $S_{8,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{8,1,1}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, \dots, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 6 & 2 & 2 \\ 6 & -2 & 2 & 2 \\ 2 & 2 & -2 & 6 \\ 2 & 2 & 6 & -2 \end{pmatrix}.$$

The curves  $A_1, A_2, A_3$  generate the Néron–Severi lattice. The divisor

$$D_8 = A_1 + A_2 \equiv A_3 + A_4$$

is ample, of square 8, base-point free, non-hyperelliptic, with  $D_8 \cdot A_j = 4$  for  $j \in \{1, \dots, 4\}$ .

**Proposition 3.32.** *The K3 surface is a complete intersection in  $\mathbb{P}^5$  with two hyperplane sections which are each the union of two degree 4 rational curves. The moduli space  $\mathcal{M}_{S_{8,1,1}}$  is unirational.*

*Proof.* One can construct these surfaces by taking two degree 4 rational normal curves  $C_1, C_3$  in two different hyperplanes  $H_1, H_3$  but such that the curves  $C_1, C_3$  meet transversely in two fixed points. Let  $X$  be a general quartic that contains  $C_1$  and  $C_3$ ; then the intersections of  $X$  with  $H_1, H_2$  are  $C_1 + C_2$  and  $C_3 + C_4$ , where  $C_2, C_4$  are two degree 4 rational normal curves. Curves  $C_1, \dots, C_4$  generate a lattice isometric to  $S_{8,1,1}$ . That construction shows that the moduli space  $\mathcal{M}_{S_{8,1,1}}$  is unirational.  $\square$

**Proposition 3.33.** *The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{8,1,1}$  is trivial.*

*Proof.* We proceed as in the proof of Proposition 3.11.  $\square$

### 3.20. The lattice $S_{10,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{10,1,1}$ . The surface  $X$  contains eight  $(-2)$ -curves  $A_1, \dots, A_8$ , with intersection matrix

$$\begin{pmatrix} -2 & 18 & 8 & 8 & 2 & 22 & 2 & 22 \\ 18 & -2 & 8 & 8 & 22 & 2 & 22 & 2 \\ 8 & 8 & -2 & 18 & 2 & 22 & 22 & 2 \\ 8 & 8 & 18 & -2 & 22 & 2 & 2 & 22 \\ 2 & 22 & 2 & 22 & -2 & 38 & 18 & 18 \\ 22 & 2 & 22 & 2 & 38 & -2 & 18 & 18 \\ 2 & 22 & 22 & 2 & 18 & 18 & -2 & 38 \\ 22 & 2 & 2 & 22 & 18 & 18 & 38 & -2 \end{pmatrix}.$$

The curves  $A_1, A_3, A_5$  generate the Néron–Severi lattice. The divisor

$$D_2 = A_1 + A_3 - A_5$$

is ample, base-point free, of square 2, with  $D_2 \cdot A_j = 4$  for  $j \in \{1, 2, 3, 4\}$  and  $D_2 \cdot A_j = 6$  for  $j \in \{5, 6, 7, 8\}$ . We have

$$4D_2 \equiv A_1 + A_2 \equiv A_3 + A_4, \quad 6D_2 \equiv A_5 + A_6 \equiv A_7 + A_8.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 3.34.** *The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$  such that there are two quartic cuspidal rational curves  $Q_4, Q'_4$  and two sextic cuspidal rational curves  $Q_6, Q'_6$  such that the cusps are on  $C_6$  and the intersection multiplicities of these curves with  $C_6$  are even at all intersection points.*

The divisor  $D_8 = 2A_1 + A_3 - A_5$  is very ample, of square 8, with  $D_8 \cdot A_j = 2, 22, 12, 12, 8, 28, 8, 28$  for  $j = 1, \dots, 8$ .

### 3.21. The lattice $S_{12,1,1}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S_{12,1,1}$ . The surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$ , with intersection matrix

$$\begin{pmatrix} -2 & 14 & 2 & 10 & 10 & 2 \\ 14 & -2 & 10 & 2 & 2 & 10 \\ 2 & 10 & -2 & 14 & 2 & 10 \\ 10 & 2 & 14 & -2 & 10 & 2 \\ 10 & 2 & 2 & 10 & -2 & 14 \\ 2 & 10 & 10 & 2 & 14 & -2 \end{pmatrix}.$$

The curves  $A_1, A_3, A_5$  generate the Néron–Severi lattice. The divisor

$$D_6 = A_1 - A_3 + A_5$$

is very ample, of square 6, with  $D_2 \cdot A_j = 6$  for  $j \in \{1, \dots, 6\}$ . We have

$$2D_6 \equiv A_1 + A_2 \equiv A_3 + A_4 \equiv A_5 + A_6;$$

thus we obtain the first part of the following proposition.

**Proposition 3.35.** *The surface  $X$  is a degree 6 surface in  $\mathbb{P}^4$  such that there are three quadric sections, each of which splits as the union of two degree 6 smooth rational curves.*

*The automorphism group of a general K3 surface  $X$  with  $\text{NS}(X) \simeq S_{12,1,1}$  is trivial.*

*Proof.* For the second part, we proceed as in the proof of Proposition 3.11. □

### 3.22. The lattice $S'_{4,1,2}$

Let  $X$  be a K3 surface with Néron–Severi lattice of type  $S'_{4,1,2}$ . The surface  $X$  contains four  $(-2)$ -curves  $A_1, \dots, A_4$ , with intersection matrix

$$\begin{pmatrix} -2 & 6 & 2 & 2 \\ 6 & -2 & 2 & 2 \\ 2 & 2 & -2 & 6 \\ 2 & 2 & 6 & -2 \end{pmatrix}.$$

This is the same intersection matrix as the four  $(-2)$ -curves in Section 3.19, but here the curves  $A_1, A_2, A_3, A_4$  generate only an index 2 subgroup of the Néron–Severi lattice. There is a basis  $e_1, e_2, e_3$  of the Néron–Severi lattice such that the intersection matrix of  $e_1, e_2, e_3$  is

$$\begin{pmatrix} -6 & 2 & 0 \\ 2 & -2 & 4 \\ 0 & 4 & -8 \end{pmatrix}.$$

In that basis, the classes of the curves are

$$\begin{aligned} A_1 &= (2, 3, 1), & A_2 &= (0, 1, 1), \\ A_3 &= (2, 3, 2), & A_4 &= (0, 1, 0), \end{aligned}$$

and the divisor

$$D_2 = (1, 2, 1)$$

is ample, of square 2, base-point free, with  $D_2 \cdot A_j = 2$  for  $j \in \{1, 2, 3, 4\}$ . We have

$$2D_2 \equiv A_1 + A_2 \equiv A_3 + A_4.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 3.36.** *The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has two 6-tangent conics.*

Let  $D_4$  be the divisor  $D_4 = (1, 3, 1)$ . It is nef, base-point free, with  $D_4 \cdot A_j = 4, 4, 8, 0$  for  $j = 1, \dots, 4$ . The linear system  $|D_4|$  defines a singular model  $Y$  of  $X$  which is a quartic in  $\mathbb{P}^3$  with a node. Since

$$2D_4 \equiv A_1 + A_2 \equiv A_3 + A_4,$$

there are two quadric sections  $Q_1, Q_2$  such that  $Q_1$  is the union of two smooth rational degree 4 curves which are the images of  $A_1, A_2$  and  $Q_2$  is a degree 8 rational curve (the image of  $A_3$ ) which contains the node (the image of  $A_4$ ).

## 4. Rank 4 lattices

### Vinberg's classification

The reference for the classification of rank 4 lattices that are Néron–Severi lattices of K3 surfaces with finite automorphism group is the article of Vinberg [Vin07]. There are two lattices such that the fundamental domain of the Weyl group is compact, and 12 for which the domain is not compact. Geometrically, the fundamental domain is compact if and only if the K3 surface has no elliptic fibration.

#### 4.1. The rank 4 and compact cases

We studied in [Roul9] the two lattices  $L(24), L(27)$  of rank 4 such that the K3 surfaces have no elliptic fibrations. The Gram matrices of the lattices  $L(24), L(27)$  are, respectively,

$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & -2 & 0 & 0 \\ 1 & 0 & -2 & 0 \\ 1 & 0 & 0 & -2 \end{pmatrix}, \begin{pmatrix} 12 & 2 & 0 & 0 \\ 2 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & -2 \end{pmatrix}.$$

The K3 surfaces with such Néron–Severi lattices are double covers of the plane branched over a smooth sextic curve  $C_6$ . For completeness, let us recall the results obtained in [Roul9].

**Theorem 4.1.** *Let  $X$  be a K3 surface with Néron–Severi lattice isometric to  $L(24)$ . The six  $(-2)$ -curves on  $X$  are pull-backs of three lines tritangent to  $C_6$ .*

*Let  $X$  be a K3 surface with Néron–Severi lattice isometric to  $L(27)$ . The eight  $(-2)$ -curves on  $X$  are pull-backs of one line tritangent to  $C_6$  and three conics 6-tangent to  $C_6$ .*

The K3 surfaces with Néron–Severi lattice isometric to  $L(24)$  have some connection with K3 surfaces of type  $S_0 \oplus \mathbf{A}_2$ , where  $S_0 = \begin{bmatrix} 0 & -3 \\ -3 & -2 \end{bmatrix}$ ; see [ACR20].

#### 4.2. The rank 4 and non-compact cases

This is the subject of another paper [ACR20]. The lattices are

$$\begin{aligned} & [8] \oplus \mathbf{A}_1^{\oplus 3}, U \oplus \mathbf{A}_1^{\oplus 2}, U(2) \oplus \mathbf{A}_1^{\oplus 2}, U(3) \oplus \mathbf{A}_1^{\oplus 2}, U(4) \oplus \mathbf{A}_1^{\oplus 2}, \\ & U \oplus \mathbf{A}_2, U(2) \oplus \mathbf{A}_2, U(3) \oplus \mathbf{A}_2, U(6) \oplus \mathbf{A}_2, \\ & S_0 \oplus \mathbf{A}_2, [4] \oplus [-4] \oplus \mathbf{A}_2, U(4) \oplus \mathbf{A}_3. \end{aligned}$$

We give the number of  $(-2)$ -curves in the different cases in the table in Section 16.

For the Hirzebruch surface  $\mathbb{F}_n$ ,  $n \geq 1$ , there is a unique negative curve  $s$  which is such that  $s^2 = -n$ . We denote by  $f$  a fiber of the natural fibration  $\mathbb{F}_n \rightarrow \mathbb{P}^1$ . For completeness, we summarize the constructions obtained in [ACR20].

**Theorem 4.2.**

- A K3 surface with  $\text{NS}(X) \simeq [8] \oplus \mathbf{A}_1^{\oplus 3}$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has six 6-tangent conics.
- A K3 surface with  $\text{NS}(X) \simeq U \oplus \mathbf{A}_1^{\oplus 2}$  is a double cover of a Hirzebruch surface,  $\eta: X \rightarrow \mathbf{F}_2$ . It is branched over the section  $s$  and a curve  $B \in |3s + 8f|$ , so that  $sB = 2$ .
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U(2) \oplus \mathbf{A}_1^{\oplus 2}$  is the minimal desingularization of the double cover of  $\mathbb{P}^2$  branched over a sextic curve with three nodal singularities.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U(3) \oplus \mathbf{A}_1^{\oplus 2}$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve with two tritangent lines and two 6-tangent conics.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U(4) \oplus \mathbf{A}_1^{\oplus 2}$  is a quartic in  $\mathbb{P}^3$  which has four hyperplane sections that decompose into the union of two conics.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U \oplus \mathbf{A}_2$  is the minimal desingularization of the double cover of  $\mathbf{F}_3$  with branch locus the section  $s$  and a reduced curve in  $B \in |3s + 10f|$ , so that  $Bs = 1$ .
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U(2) \oplus \mathbf{A}_2$  is the minimal resolution of the double cover of  $\mathbb{P}^2$  branched over a sextic with two nodes such that the line through the two nodes is tangent to the sextic curve in a third point.
- A K3 surface with  $\text{NS}(X) \simeq U(3) \oplus \mathbf{A}_2$  is a quartic in  $\mathbb{P}^3$  which has a hyperplane section that is the union of four lines.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq U(6) \oplus \mathbf{A}_2$  is a quartic in  $\mathbb{P}^3$  which has three hyperplane sections that decompose into the union of two conics.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq \begin{bmatrix} 0 & -3 \\ -3 & -2 \end{bmatrix} \oplus \mathbf{A}_2$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has three tritangent lines.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq [4] \oplus [-4] \oplus \mathbf{A}_2$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has two tritangent lines and one 6-tangent conic.
- A K3 surface  $X$  with  $\text{NS}(X) \simeq [4] \oplus \mathbf{A}_3$  is the minimal resolution of a double cover of  $\mathbb{P}^2$  branched over a sextic curve with one node; through that node go two lines that are tangent to the sextic at every other intersection point.

We will only add the following result.

**Proposition 4.3.** *A general K3 surface  $X$  with a Néron–Severi lattice isometric to one of the lattices*

$$U(4) \oplus \mathbf{A}_1^{\oplus 2}, U(3) \oplus \mathbf{A}_2, U(6) \oplus \mathbf{A}_2$$

*has trivial automorphism group.*

*Proof.* For each of these lattices, the Hilbert basis of its nef cone is described in [ACR20]. Then as in Proposition 3.11, one can check that there are no hyperelliptic involutions and conclude that the automorphism group is trivial.  $\square$


## 5. Rank 5 lattices

### Nikulin’s classification for higher ranks

The list of lattices  $L$  of rank at least 5 such that the K3 surfaces with  $\text{NS}(X) \simeq L$  have finite automorphism group is given in [Nik14]; that list is obtained from the paper [Nik83].

#### 5.1. The lattice $U \oplus \mathbf{A}_1^{\oplus 3}$

There exist seven  $(-2)$ -curves  $A_1, \dots, A_7$  on  $X$ , with dual graph



The curves  $A_1, \dots, A_5$  generate the Néron-Severi lattice. The divisor

$$D_{18} = 3A_1 + 5A_2 + A_3 + A_4 + 4A_5$$

is ample, with  $D_{18} \cdot A_j = 1$  for  $j \in \{1, 2, 3, 4\}$  and  $D_{18} \cdot A_j = 2$  for  $j \in \{5, 6, 7\}$ . The divisor

$$D_2 = 2A_1 + 2A_2 + A_3 + A_4 + A_5$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_j = 0$  for  $j \in \{1, 2, 3, 4\}$  and  $D_2 \cdot A_j = 2$  for  $j \in \{5, 6, 7\}$ . We also have

$$D_2 \equiv 2A_1 + A_2 + 2A_3 + A_4 + A_6 \equiv 2A_1 + A_2 + A_3 + 2A_4 + A_7.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 5.1.** *The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve with a  $\mathbf{d}_4$  singularity  $q$ .*


The three curves  $A_5, A_6, A_7$  are mapped by the double cover map to the three lines that are the tangents to the three branches of the singularity  $q$ .

## 5.2. The lattice $U(2) \oplus A_1^{\oplus 3}$ and the del Pezzo surface of degree 5

The K3 surface  $X$  is the minimal resolution of the double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with four nodes  $p_1, \dots, p_4$  in general position. It is also the double cover branched over the strict transform of  $C_6$  in the degree 5 del Pezzo surface  $Z$  which is the blow-up at  $p_1, \dots, p_4$ . The surface  $X$  contains 10  $(-2)$ -curves denoted by  $A_{i,j}$ , for  $\{i, j\} \subset \{1, \dots, 5\}$  with  $i < j$  (the pull-back of the 10  $(-1)$ -curves on the del Pezzo surface). One has

$$A_{ij} \cdot A_{st} = 2$$

if and only if  $\#\{i, j, s, t\} = 4$ ; else  $A_{ij} \cdot A_{st} = 0$  or  $-2$ . The dual graph of the configuration is the Petersen graph



with weight 2 on the edges.

*Remark 5.2.* If  $C$  is a general curve of genus 6, then there exists a map  $C \rightarrow \mathbb{P}^2$  with image a sextic curve with four nodes. Using that property, Artebani and Kondō describe in [AK11] the moduli space of genus 6 curves and their link with K3 surfaces and the quintic del Pezzo surface. They study that moduli space as a quotient of a bounded symmetric domain.



### 5.3. The lattice $U(4) \oplus A_1^{\oplus 3}$

The K3 surface  $X$  contains 24  $(-2)$ -curves  $A_1, \dots, A_{24}$ . Up to permutation, one can suppose that  $A_1, A_3, A_5, A_7, A_9$  have the following intersection matrix:

$$\begin{pmatrix} -2 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 & 4 \\ 0 & 0 & -2 & 2 & 2 \\ 0 & 0 & 2 & -2 & 2 \\ 0 & 4 & 2 & 2 & -2 \end{pmatrix}.$$

These five curves generate  $NS(X)$ . The divisor

$$D_2 = -A_1 + A_3 + A_9$$

is ample, of square 2, with  $D_2 \cdot A_j = 2$  for  $j \in \{1, \dots, 24\}$  and, up to permutation of the indices,

$$2D_2 \equiv A_{2k-1} + A_{2k}$$


for any  $k \in \{1, \dots, 12\}$ . The classes of  $A_{11}, A_{13}, \dots, A_{23}$  in the basis  $A_1, A_3, A_5, A_7, A_9$  are

$$\begin{aligned} A_{11} &= (0, -1, 1, 1, 0), & A_{13} &= (-1, 0, 1, 1, 0), \\ A_{15} &= (-1, 0, 0, 1, 1), & A_{17} &= (-2, 1, 0, 1, 1), \\ A_{19} &= (-2, 0, 1, 1, 1), & A_{21} &= (-1, 0, 1, 0, 1), \\ A_{23} &= (-2, 1, 1, 0, 1), \end{aligned}$$

so that we know the 24 classes of  $(-2)$ -curves in  $X$ . By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 5.3.** *The K3 surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has 12 6-tangent conics.*

There exists a partition of the 24  $(-2)$ -curves into three sets  $S_1, S_2, S_3$  of 8 curves each such that for curves  $B, B'$  in two different sets  $S, S'$ , one has  $BB' = 0$  or 4 and for any  $B \in S$ , there are exactly 4 curves  $B'$  in  $S'$  such that  $BB' = 4$ , and symmetrically for  $B'$ . Therefore,  $S$  and  $S'$  form an  $8_4$  configuration. This is the so-called Möbius configuration (see [Cox50]). The following graph is the Levi graph of that  $8_4$  configuration; this is the graph of the 4-dimensional hypercube (see [Cox50]). Vertices in red are curves in  $S$ , vertices in blue are curves in  $S'$ , and an edge links a red curve to a blue curve if and only if their intersection number is 4.



From that graph, we can moreover read the intersection numbers of the curves in  $S$  (and  $S'$ ) as follows. For any red curve  $B$ , there are four blue curves linked to it by an edge. Consider the complementary set of blue curves; this is another set of four blue curves, all linked through an edge to the same red curve  $B'$ . Then we have  $BB' = 6$ , and for any other red curve  $B'' \notin \{B, B'\}$ , we have  $BB'' = 2$ . Symmetrically, the intersection numbers between the blue curves follow the same rule.

In [Nik83, Proof of Theorem 8.1.1], Nikulin studies the lattice  $U(4) \oplus \mathbf{A}_1^{\oplus 3}$  in detail, obtaining that it contains only 24  $(-2)$ -curves, and he describes the fundamental polygon of the action of the Weyl group by an embedding in the 4-dimensional Euclidian space. There, the polyhedron formed by the 24 vertices is the dual of the polyhedron formed by the roots of type  $\mathbf{D}_4$ .

There is a second geometric model which is as follows. The divisor

$$D_4 = -A_1 + 2A_3 + A_9$$


is a nef, non-hyperelliptic divisor of square 4, with  $D_4 \cdot A_j = 0$  if and only if  $j = 3$ . We have, moreover,

$$D_4 \equiv A_1 + A_{10} \equiv A_5 + A_{22} \equiv A_7 + A_{16} \equiv A_{13} + A_{20},$$

and the intersection number of these eight curves with  $D_4$  is 2. Therefore, the linear system  $|D_4|$  gives a singular model of  $X$  as a quartic in  $\mathbb{P}^3$  with a unique node and four hyperplane sections which are unions of two conics. One can check that the eight  $(-2)$ -curves which are mapped to the conics and the  $(-2)$ -curve which is contracted to the node generate the lattice  $U(4) \oplus \mathbf{A}_1^{\oplus 3}$ .

#### 5.4. The lattice $U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2$

The K3 surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$ ; their configuration is



These curves generate the Néron–Severi lattice. The divisor

$$D_{20} = 5A_1 + 5A_2 + 6A_3 + 3A_4 + A_5$$

is ample, of square 20, with  $D_{20} \cdot A_j = 1$  for  $j \leq 5$  and  $D_{20} \cdot A_6 = 2$ . The divisor

$$D_4 = 2A_1 + 2A_2 + 3A_3 + 2A_4 + A_5$$

is nef, of square 4, base-point free and hyperelliptic since

$$D_4(A_1 + A_2 + A_3) = 2,$$

and  $A_1 + A_2 + A_3 \equiv A_5 + A_6$  is a fiber of an elliptic fibration. One has  $D_4 \cdot A_1 = D_4 \cdot A_2 = 1$ ,  $D_4 \cdot A_6 = 2$  and  $D_4 \cdot A_j = 0$  for  $j \in \{3, 4, 5\}$ . Moreover,


$$D_4 \equiv A_3 + 2A_4 + 3A_5 + 2A_6.$$

**Proposition 5.4.** *The double cover induced by  $|D_4|$  factors through  $\eta: X \rightarrow \mathbf{F}_2$ . The image by  $\eta$  of  $A_4$  is  $s$  (where  $s$  is the section such that  $s^2 = -2$ ). The branch locus is the union of the section  $s$  and  $B \in |3s + 8f|$ , so that  $Bs = 2$ . Let  $f_1, f_2$  be the fibers through the points  $p$  and  $q$ , the intersection points of  $s$  and  $B$ . The curve  $f_1$  cuts  $B$  with multiplicity 2 at another point, and the image by  $\eta$  of the curves  $A_1, A_2$  is  $f_1$ , the image of  $A_6$  is  $f_2$ , and  $A_3, A_5$  are contracted to  $p$  and  $q$ , respectively.*

*Proof.* We apply Theorem 2.3, case a) iii) v). □

#### 5.5. The lattice $U \oplus \mathbf{A}_3$

The K3 surface  $X$  contains five  $(-2)$ -curves  $A_1, \dots, A_5$ ; their configuration is



These curves generate the Néron–Severi lattice. In that basis, the divisor

$$D_{50} = (5, 11, 9, 9, 8)$$

is ample, of square 50, with  $D_{50} \cdot A_j = 1$  for  $j \leq 4$  and  $D_{50} \cdot A_5 = 2$ . We have  $D_{50} \cdot A_j = 0$  for  $j \leq 4$  and  $D_{50} \cdot A_5 = 2$ . The divisor

$$D_4 = 2A_1 + 4A_2 + 3A_3 + 3A_4 + 2A_5$$


is nef, of square 4, and  $|D_4|$  is base-point free hyperelliptic since  $D_4(A_2 + A_3 + A_4 + A_5) = 2$ ; one has  $D_4 \cdot A_j = 0$  for  $j \leq 4$  and  $D_4 \cdot A_5 = 2$ .

**Proposition 5.5.** *The double cover induced by  $|D_4|$  factors through  $\eta: X \rightarrow \mathbf{F}_2$ . The image by  $\eta$  of  $A_1$  is the section  $s$ . The branch locus is the union of the section  $s$  and  $B \in |3s + 8f|$ , so that  $Bs = 2$ . The intersection of  $B$  and  $s$  is tangent at one point  $q$ , forming an  $\mathfrak{a}_3$  singularity. The curve  $A_5$  is mapped onto the fiber through  $q$ , and the curves  $A_2, A_3, A_4$  are mapped to  $q$ .*

*Proof.* We apply Theorem 2.3, case a) iii) v). □

### 5.6. The lattice $[4] \oplus D_4$

The K3 surface  $X$  contains five  $(-2)$ -curves  $A_1, \dots, A_5$ ; their configuration is



These curves generate the Néron–Severi lattice. The divisor

$$D_{22} = A_1 + 3 \sum_{j=1}^5 A_j$$

is ample, of square 22, with  $D_{22} \cdot A_3 = 3$  and  $D_{22} \cdot A_j = 1$  for  $j \neq 3$ . The divisor

$$D_2 = \sum_{j=1}^5 A_j$$

is nef, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_3 = 1$ ,  $D_2 \cdot A_j = 0$  for  $j \in \{2, 4, 5\}$ . By using the linear system  $|D_2|$ , we obtain the following.


**Proposition 5.6.** *The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve with three nodes on a line.*

The image of  $A_1, A_3$  is the line through the three nodes; the curves  $A_2, A_4, A_5$  are contracted to the nodes.

The divisor  $D_4 = A_1 + 2A_2 + 3A_3 + 2A_4 + 2A_5$  is nef, non-hyperelliptic, with  $D_4 \cdot A_1 = 4$  and  $D_4 \cdot A_k = 0$  for  $k \geq 2$ . It defines a model of the K3 surface as a quartic in  $\mathbb{P}^3$  with a  $D_4$  singularity.

### 5.7. The lattice $[8] \oplus D_4$

The K3 surface  $X$  contains seven  $(-2)$ -curves  $A_1, \dots, A_7$ ; their configuration is



The curves  $A_1, A_2, A_3, A_4, A_7$  generate the Néron–Severi lattice. The divisor

$$D_6 = 2A_1 + 2A_4 + A_7$$

is ample, of square 6, with  $D_6 \cdot A_j = 1$  for  $j \leq 6$  and  $D_6 \cdot A_7 = 2$ . The divisor

$$D_2 = A_1 + A_4 + A_7$$

is nef, of square 2, with  $D_2 \cdot A_j = 1$  for  $j \leq 6$  and  $D_2 \cdot A_7 = 0$ . We have

$$D_2 \equiv A_2 + A_5 + A_7 \equiv A_3 + A_6 + A_7.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 5.7.** *The surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve with one node, and there are three lines through that node such that each line is tangent to the sextic at its other intersection points.*


One can obtain another interesting geometric model of the K3 surface  $X$  as follows. The divisor  $D_8 = 3A_1 + 2A_2 + 2A_3 + A_4 + 4A_7$  is big and nef, base-point free, non-hyperelliptic, with  $D_8 \cdot A_k = 0, 0, 0, 8, 8, 8, 0$  for  $k = 1, \dots, 7$ . The image of the K3 surface by the map associated to  $|D_8|$  is a degree 8 surface in  $\mathbb{P}^5$  with a  $D_4$  singularity.

### 5.8. The lattice $[16] \oplus D_4$

The K3 surface  $X$  contains eight  $(-2)$ -curves  $A_1, \dots, A_8$ ; their intersection matrix is

$$\begin{pmatrix} -2 & 3 & 0 & 1 & 0 & 1 & 0 & 1 \\ 3 & -2 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 3 & 0 & 1 & 0 & 1 \\ 1 & 0 & 3 & -2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -2 & 3 & 0 & 1 \\ 1 & 0 & 1 & 0 & 3 & -2 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & -2 & 3 \\ 1 & 0 & 1 & 0 & 1 & 0 & 3 & -2 \end{pmatrix}.$$

Their configuration is



where the thick lines have weight 3. The curves  $A_1, A_2, A_3, A_5, A_7$  generate  $\text{NS}(X)$ . The divisor

$$D_2 = A_1 + A_2$$

is ample of square 2, with  $D_2 \cdot A_j = 1$  for  $j \in \{1, \dots, 8\}$ . We have also

$$D_2 \equiv A_3 + A_4 \equiv A_5 + A_6 \equiv A_7 + A_8.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 5.8.** *The surface is a double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has four tritangent lines.*


One can obtain another interesting geometric model of the K3 surface  $X$  as follows. The divisor  $D_{16} = 4A_1 + A_2 + 5A_4 + 3A_5 + 3A_7$  is nef, non-hyperelliptic, of square 16, with  $D_{16} \cdot A_k = 0$  for curves  $A_1, A_4, A_5, A_7$  and  $D_{16} \cdot A_k = 16$  for the other curves. The image of the K3 surface under the map associated to  $|D_{16}|$  is a degree 16 surface in  $\mathbb{P}^9$  with a  $\mathbf{D}_4$  singularities, which is the image of the curves  $A_1, A_4, A_5, A_7$ .

### 5.9. The lattice $[6] \oplus \mathbf{A}_2^{\oplus 2}$

The K3 surface  $X$  contains 10  $(-2)$ -curves  $A_1, \dots, A_{10}$ ; their intersection matrix is

$$\begin{pmatrix} -2 & 3 & 1 & 0 & 0 & 1 & 0 & 1 & 2 & 0 \\ 3 & -2 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 2 \\ 1 & 0 & -2 & 3 & 0 & 1 & 0 & 1 & 2 & 0 \\ 0 & 1 & 3 & -2 & 1 & 0 & 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 1 & -2 & 3 & 1 & 0 & 0 & 2 \\ 1 & 0 & 1 & 0 & 3 & -2 & 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & -2 & 3 & 0 & 2 \\ 1 & 0 & 1 & 0 & 0 & 1 & 3 & -2 & 2 & 0 \\ 2 & 0 & 2 & 0 & 0 & 2 & 0 & 2 & -2 & 6 \\ 0 & 2 & 0 & 2 & 2 & 0 & 2 & 0 & 6 & -2 \end{pmatrix}.$$

The configuration of the first eight curves is



where the thick lines have weight 3, the curves  $A_k, k \in \{1, 3, 6, 8\}$  (with a red vertex) are such that  $A_k \cdot A_9 = 2, A_k \cdot A_{10} = 0$ , and the curves  $A_k, k \in \{2, 4, 5, 7\}$  (with a black vertex) are such that  $A_k \cdot A_{10} = 2, A_k \cdot A_9 = 0$ . The curves  $A_1, A_3, A_5, A_7, A_9$  generate  $\text{NS}(X)$ . The divisor

$$D_2 = A_1 + A_3 - A_5 - A_7 + A_9$$

is ample, of square 2, with  $D_2 \cdot A_j = 1$  for  $j \leq 8, D_2 \cdot A_9 = D_2 \cdot A_{10} = 2$ . We have

$$D_2 \equiv A_{2k-1} + A_{2k}, k \in \{1, 2, 3, 4\}$$

and  $2D_2 \equiv A_9 + A_{10}$ . By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 5.9.** *The surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a smooth sextic curve which has four tritangent lines and one 6-tangent conic.*

*Remark 5.10.* The Néron–Severi lattice is also generated by  $A_1, A_2, A_3, A_5, A_7$ .

It is interesting to compare this case with the previous lattice case, where four tritangent lines are also involved.

**Proposition 5.11.** *Let  $X$  be a K3 surface with  $\text{NS}(X) \simeq [6] \oplus \mathbf{A}_2^{\oplus 2}$ . There exist linear forms  $\ell_1, \dots, \ell_4 \in H^0(\mathbb{P}^2, \mathcal{O}(1))$ , a quadric  $q_2 \in H^0(\mathbb{P}^2, \mathcal{O}(2))$  and a cubic  $f_3 \in H^0(\mathbb{P}^2, \mathcal{O}(3))$  such that  $X$  is the double cover of  $\mathbb{P}^2$  branched over the curve*

$$C_6: \ell_1 \ell_2 \ell_3 \ell_4 q_2 - f_3^2 = 0.$$

The moduli space of K3 surfaces  $X$  with  $\text{NS}(X) \simeq [6] \oplus \mathbf{A}_2^{\oplus 2}$  is unirational.

*Proof.* Let us consider the map

$$\Phi: H^0(\mathbb{P}^2, \mathcal{O}(1))^{\oplus 4} \oplus H^0(\mathbb{P}^2, \mathcal{O}(2)) \oplus H^0(\mathbb{P}^2, \mathcal{O}(3)) \rightarrow H^0(\mathbb{P}^2, \mathcal{O}(6))$$

defined by

$$w := (\ell_1, \ell_2, \ell_3, \ell_4, q_2, f_3) \mapsto f_{6,w} := \ell_1 \ell_2 \ell_3 \ell_4 q_2 - f_3^2.$$

Suppose that  $w$  is general, so that the double cover branched over  $C_w = \{f_{6,w} = 0\}$  is a K3 surface. The pull-backs of  $\ell_k = 0$ ,  $k = 1, \dots, 4$ , and  $q_2 = 0$  are pairs of  $(-2)$ -curves, for which Example 5.12 below shows that (for a suitable order) these curves intersect according to the above matrix since intersection numbers are preserved for flat families of surfaces.

The map  $\Phi$  is invariant under the action of the transformations

$$(\ell_1, \ell_2, \ell_3, \ell_4, q_2) \mapsto (\alpha \ell_1, \beta \ell_2, \gamma \ell_3, \delta \ell_4, \epsilon q_2, f_3)$$

for  $\alpha\beta\gamma\delta\epsilon = 1$ . Since the curves  $\ell_j = 0$ ,  $j \in \{1, \dots, 4\}$ , and  $q_2 = 0$  are the images of the 10  $(-2)$ -curves in the double cover  $X$  ramified over  $f_{6,w}$ , we have

$$\Phi(w) = \Phi(w')$$

if and only if, up to permutation and the action of the above transformations,  $w = \lambda w'$  for some  $\lambda \neq 0$ , where

$$\lambda \cdot (\ell_1, \ell_2, \ell_3, \ell_4, q_2, f_3) = (\lambda \ell_1, \lambda \ell_2, \lambda \ell_3, \lambda \ell_4, \lambda^2 q_2, \lambda^3 f_3).$$

The image of  $\Phi$  modulo the action of  $PGL_3$  is therefore a unirational space of dimension

$$4 \cdot 3 + 6 + 10 - 4 - 9 = 15,$$

with an open set which is birational to the moduli space of K3 surfaces  $X$  with  $\text{NS}(X) \simeq [6] \oplus \mathbf{A}_2^{\oplus 2}$ . □

*Example 5.12.* Let us take

$$\begin{aligned} \ell_1 &= 16y + 29z, \ell_2 = 22x + y + 27z, \ell_3 = 17y + 29z, \ell_4 = 25x + 29y + 11z, \\ q_2 &= 31x^2 + 16xy + 11y^2 + 11xz + 9yz + 17z^2, \\ f_3 &= 6x^3 + 8x^2y + 17xy^2 + 5y^3 + 17x^2z + 18xyz + 19y^2z + 16xz^2 + 8yz^2 + 23z^3. \end{aligned}$$

Let  $X$  be the associated K3 surface, and let  $X_q$  be its reduction over the field  $\mathbb{F}_q$ . Using the Tate conjecture, one computes that the Picard numbers of  $X_{17^2}$  and  $X_{23^2}$  are both 6, and using the Artin–Tate conjecture and Van Luijk’s trick, one obtains that  $X$  has Picard number 5. One can check that the 10  $(-2)$ -curves above the lines  $\ell_k = 0$  and the conic  $q_2 = 0$  have the above intersection matrix and  $\text{NS}(X) \simeq [6] \oplus \mathbf{A}_2^{\oplus 2}$ .


## 6. Rank 6 lattices

### 6.1. The lattice $U(3) \oplus \mathbf{A}_2^{\oplus 2}$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U(3) \oplus \mathbf{A}_2^{\oplus 2}.$$

The surface contains 12  $(-2)$ -curves  $A_1, \dots, A_{12}$ , with configuration



where for clarity we represented the same curve several times.

The Néron–Severi lattice is generated by  $A_1, A_3, A_5, A_7, A_9, A_{11}$ . The divisor

$$D_2 = -A_1 + A_3 + A_5 + A_7 + A_9 - A_{11}$$

is ample, base-point free, of square 2, and the 12  $(-2)$ -curves  $A_1, \dots, A_{12}$  on  $X$  are of degree 1 for  $D_2$ . We have

$$D_2 \equiv A_{2k-1} + A_{2k}$$

for  $k \in \{1, \dots, 6\}$ , and by using the linear system  $|D_2|$ , we obtain the following.

**Proposition 6.1.** *The surface  $X$  is a double cover  $\pi: X \rightarrow \mathbb{P}^2$  of  $\mathbb{P}^2$  branched over a smooth sextic curve  $C_6$ . The 12  $(-2)$ -curves on  $X$  are pull-backs of 6 lines that are tritangent to the sextic curve. There exist linear forms  $\ell_1, \dots, \ell_6 \in H^0(\mathbb{P}^2, \mathcal{O}(1))$  and a cubic  $f_3 \in H^0(\mathbb{P}^2, \mathcal{O}(3))$  such that the curve  $C_6$  is given by*

$$\ell_1 \ell_2 \ell_3 \ell_4 \ell_5 \ell_6 - f_3^2 = 0.$$

*The moduli space of K3 surfaces  $X$  with  $\text{NS}(X) \simeq U(3) \oplus \mathbf{A}_2^{\oplus 2}$  is unirational.*

*Proof.* Let us consider the map

$$\Phi: H^0(\mathbb{P}^2, \mathcal{O}(1))^{\oplus 6} \oplus H^0(\mathbb{P}^2, \mathcal{O}(3)) \rightarrow H^0(\mathbb{P}^2, \mathcal{O}(6))$$

defined by

$$w := (\ell_1, \ell_2, \ell_3, \ell_4, \ell_5, \ell_6, f_3) \mapsto f_{6,w} := \ell_1 \ell_2 \ell_3 \ell_4 \ell_5 \ell_6 - f_3^2.$$

When the sextic curve  $C_w: f_{6,w} = 0$  is smooth, the K3 surface  $Y_w$  which is the double cover of  $\mathbb{P}^2$  branched over  $C_w$  contains 12  $(-2)$ -curves over the 6 tritangent lines  $\{\ell_k = 0\}$ ,  $k = 1, \dots, 6$ . Example 6.2 below shows that these  $(-2)$ -curves generate a sublattice isometric to  $U(3) \oplus \mathbf{A}_2^{\oplus 2}$ . The map  $\Phi$  is invariant under the action of the transformations

$$(\ell_1, \ell_2, \ell_3, \ell_4, \ell_5, \ell_6) \mapsto (\alpha \ell_1, \beta \ell_2, \gamma \ell_3, \delta \ell_4, \epsilon \ell_5, \mu \ell_6, f_3)$$

for  $\alpha \beta \gamma \delta \epsilon \mu = 1$ . Since the curves  $\ell_j = 0$ ,  $j \in \{1, \dots, 6\}$ , are the images of the 12  $(-2)$ -curves in the double cover ramified over  $f_6$ , we have

$$\Phi(w) = \Phi(w')$$

if and only if, up to permutation and the action of the above transformations,  $w = \lambda \cdot w'$  for some  $\lambda \neq 0$ , where

$$\lambda \cdot (\ell_1, \ell_2, \ell_3, \ell_4, \ell_5, \ell_6, f_3) = (\lambda \ell_1, \lambda \ell_2, \lambda \ell_3, \lambda \ell_4, \lambda \ell_5, \lambda \ell_6, \lambda^3 f_3).$$

The image of  $\Phi$  modulo the action of  $PGL_3$  is therefore a unirational space of dimension

$$6 \cdot 3 + 10 - 5 - 9 = 14,$$

with an open set which is birational to the moduli space of K3 surfaces  $X$  with  $\text{NS}(X) \simeq U(3) \oplus \mathbf{A}_2^{\oplus 2}$ .  $\square$

*Example 6.2.* As an example, one can take

$$\begin{aligned} \ell_1 &= 15x + 5y + 2z, \ell_2 = 20x + 12y + 17z, \ell_3 = 22x + 22y + 16z, \\ \ell_4 &= 8x + y + 7z, \ell_5 = 10x + 15y + 15z, \ell_6 = 6x + 22y + 20z, \\ f_3 &= 18x^2y + 13xy^2 + 16y^3 + 16x^2z + 17xyz + 22y^2z + 6xz^2 + 10yz^2 + 20z^3; \end{aligned}$$

this gives a smooth K3 surface  $X$ . Its reduction modulo 23 is a smooth surface  $X_{23}$ . Using the Artin–Tate conjecture, one finds that its Picard number is 4 and the discriminant of the Néron–Severi lattice has order  $3^4$ . Using the pull-back of the six lines  $\{\ell_k = 0\}$ , one can check that the Néron–Severi lattice contains the lattice  $U(3) \oplus \mathbf{A}_2^{\oplus 2}$  with discriminant  $3^4$ . Thus, one has  $\text{NS}(X_{23}) \simeq U(3) \oplus \mathbf{A}_2^{\oplus 2}$ , and that implies that the Néron–Severi lattice of  $X$  is also isometric to  $U(3) \oplus \mathbf{A}_2^{\oplus 2}$ .


*Remark 6.3.* In [Nik83, Proof of Theorem 6.4.1], Nikulin constructs some surfaces  $X$  with Néron–Severi lattice  $U(3) \oplus \mathbf{A}_2^{\oplus 2}$  as the minimal desingularization of a triple cover of a smooth quadric  $Q \subset \mathbb{P}^3$  branched over a  $(3, 3)$ -curve  $C$  with two singularities  $\mathbf{a}_1$ . In particular, the automorphism group of such a surface  $X$  has order at least 6. According to [Kon89], the general surface with  $\text{NS}(X) \simeq U(3) \oplus \mathbf{A}_2^{\oplus 2}$  has automorphism group isomorphic to  $\mathbb{Z}/2\mathbb{Z}$ .

## 6.2. The lattice $U(4) \oplus \mathbf{D}_4$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U(4) \oplus \mathbf{D}_4.$$

The surface  $X$  contains nine  $(-2)$ -curves  $A_1, \dots, A_9$ , with configuration



The class

$$D_6 = A_1 + 2A_2 + 2A_6$$

is ample with  $D_6^2 = 6$ . The divisor  $E = A_2 + A_6$  is the class of a fiber, and  $D_6 = 2E + A_9$ . For  $j > 1$ , one has  $EA_j = 0$  and  $D_6 \cdot A_j = 1$ ; moreover,  $EA_1 = D_6 \cdot A_1 = 2$ . The curves  $A_2, \dots, A_9$  are such that the divisors

$$A_k + A_{k+4}, \quad k \in \{2, 3, 4, 5\}$$

are the reducible fibers of the elliptic fibration induced by  $E$ .

The divisor

$$D_2 = A_1 + A_2 + A_6 = E + A_1$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = 0$  and  $D_2 \cdot A_j = 1$  for  $j \geq 2$ . We have

$$D_2 \equiv A_1 + A_3 + A_7 \equiv A_1 + A_4 + A_8 \equiv A_1 + A_5 + A_9.$$

By using the linear system  $|D_2|$ , we obtain the following.


**Proposition 6.4.** *The K3 surface  $X$  is a double cover of the plane branched over a nodal sextic curve such that there exist four lines through the node that are tangent to the sextic at other intersection points.*

## 6.3. The lattice $U \oplus \mathbf{A}_4$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U \oplus \mathbf{A}_4.$$

The surface  $X$  contains six  $(-2)$ -curves  $A_1, \dots, A_6$  with dual graph





These six curves generate the Néron–Severi lattice. The class

$$D_{50} = (8, 9, 11, 5, 9, 8)$$

in the basis  $A_1, \dots, A_6$  is ample, and  $D_{50}^2 = 50$ . The six  $(-2)$ -curves have degree 1 for  $D_{50}$ . The class

$$F = A_1 + A_2 + A_3 + A_5 + A_6$$

is the class of a fiber of type  $I_5$ . The curve  $A_4$  is a section of the elliptic fibration. The divisor

$$D_2 = 2F + A_4$$

is nef, of square 2, with  $D_2 \cdot A_j = 0$  for  $j \neq 3$ . It has base points, but  $D_8 = 2D_2$  is base-point free.

**Proposition 6.5.** *The linear system  $|D_8|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  that is branched over  $A_4$ ; the image of  $A_4$  is the unique section  $s$  of the Hirzebruch surface  $\mathbf{F}_4$  with  $s^2 = -4$ . The branch curve of  $\varphi$  is the disjoint union of  $s$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a unique singularity  $q$  of type  $\mathbf{a}_4$  which is the image of  $A_1, A_2, A_5, A_6$ , and the image of  $A_3$  by  $\varphi$  is the fiber  $f$  through  $q$ . The local intersection number of  $f$  and  $B$  at  $q$  is 2; in other words,  $f$  is transverse to the branch of  $B$  at  $q$ .*


*Proof.* We apply Theorem 2.3, case a) i). □

#### 6.4. The lattice $U \oplus \mathbf{A}_1 \oplus \mathbf{A}_3$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\mathrm{NS}(X) = U \oplus \mathbf{A}_1 \oplus \mathbf{A}_3.$$

The surface contains seven  $(-2)$ -curves  $A_1, \dots, A_7$  with configuration



The class

$$D_{42} = 6A_1 + 8A_2 + 5A_3 + 3A_4 + A_5 + A_6$$

is ample, of square 42. The divisors  $F_1 = A_4 + A_5 + A_6 + A_7$  and  $F_2 = A_1 + A_2$  are two fibers of an elliptic fibration for which the class  $A_3$  is a section. The class

$$D_2 = A_1 + 2A_2 + 2A_3 + 2A_4 + A_5 + A_6 \equiv A_2 + 2A_3 + 3A_4 + 2A_5 + 2A_6 + A_7$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_7 = 2$  and  $D_2 \cdot A_j = 0$  for  $j \in \{2, 3, 4, 5, 6\}$ . Let  $\eta: X \rightarrow \mathbb{P}^2$  be the associated double cover and  $C_6$  be the branch curve.


**Proposition 6.6.** *The curve  $C_6$  has a  $\mathbf{d}_5$  singularity  $q$  onto which the curves  $A_j$ ,  $j = 2, \dots, 6$  are contracted. From the above equivalence relation between divisors, we see that the images of  $A_1$  and  $A_7$  are the two tangents to the branches of the singularity  $q$ .*

#### 6.5. The lattice $U \oplus \mathbf{A}_2^{\oplus 2}$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\mathrm{NS}(X) = U \oplus \mathbf{A}_2^{\oplus 2}.$$

The surface contains seven  $(-2)$ -curves  $A_1, \dots, A_7$  with configuration



The class

$$D_{20} = 5A_1 + 5A_2 + 6A_3 + 3A_4 + A_5$$

is ample, of square 20, and the curves  $A_j$  have degree 1 for  $D_{20}$ . The divisor

$$D_4 = 2A_1 + 2A_2 + 3A_3 + 2A_4 + A_5 \equiv A_3 + 2A_4 + 3A_5 + 2A_6 + 2A_7$$

is nef, of square 4, base-point free, with  $D_4 \cdot A_j = 1$  for  $j \in \{1, 2, 6, 7\}$  and  $D_4 \cdot A_j = 0$  for  $j \in \{3, 4, 5\}$ . Since  $D_4 F_1 = 2$ , the linear system  $|D_4|$  is hyperelliptic. By using the linear system  $|D_4|$ , we obtain the following.

**Proposition 6.7.** *The surface is a double cover of  $\mathbf{F}_2$  branched over the unique section  $s$  with  $s^2 = -2$  and a curve  $B \in |3s + 8f|$ . The fibers  $f_p, f_q$  through the two intersection points  $p, q$  of  $s$  and  $B$  are tangent to the curve  $B$  at another intersection point with  $B$ . The image of  $A_1, A_2$  is  $f_p$ , the image of  $A_6, A_7$  is  $f_q$ , and the curves  $A_3, A_5$  are mapped to the points  $p, q$ .*


*Proof.* This comes from Theorem 2.3, case a) iii) v). □

## 6.6. The lattice $U \oplus \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_2$

Let  $X$  be a K3 surface with Néron-Severi lattice

$$\text{NS}(X) = U \oplus \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_2.$$

The surface contains eight  $(-2)$ -curves  $A_1, \dots, A_8$ , with configuration



The class

$$D_{18} = 4A_1 + 4A_2 + 5A_3 + 3A_4 + A_5 + A_6$$

is ample, of square 18, with  $D_{18} \cdot A_j = 1$  for  $j \leq 6$  and  $D_{18} \cdot A_7 = D_{18} \cdot A_8 = 2$ . The three divisors

$$F = A_1 + A_2 + A_3 \equiv A_5 + A_7 \equiv A_6 + A_8$$

are classes of fibers of type  $I_3$  or  $IV$  and of type  $I_2 + I_2$  or  $III$ ; the  $(-2)$ -curve  $A_4$  is a section of the elliptic fibration. The class

$$D_2 = A_1 + A_2 + 2A_3 + 2A_4 + A_5 + A_6$$

is of square 2 and base-point free. Let  $\eta: X \rightarrow \mathbb{P}^2$  be the associated double cover map, and let  $C_6$  be the branch curve. For  $j = 1, \dots, 8$ , one has

$$D_2 \cdot A_j = 1, 1, 0, 0, 0, 0, 2, 2,$$

respectively. This leads to the following.

**Proposition 6.8.** *The morphism  $\eta$  contracts the curves  $A_3, A_4, A_5, A_6$  to a  $\mathbf{d}_4$  singularity  $q$ , the curve  $A_1 + A_2$  is the pull-back of the tangent line of a branch of the singularity, a line which is tangent to the sextic at another intersection point, and the curves  $A_7, A_8$  are pull-backs of the lines tangent to other branches, each of which lines meets  $C_6$  in two other points.*

### 6.7. The lattice $U(2) \oplus \mathbf{A}_1^{\oplus 4}$ and del Pezzo surfaces of degree 4

The class of square 8

$$D_8 = (2, 2, -1, -1, -1, -1) \in U(2) \oplus \mathbf{A}_1^{\oplus 4}$$

has no  $(-2)$ -vectors perpendicular to it; thus we have a marking  $\text{NS}(X) \simeq U(2) \oplus \mathbf{A}_1^{\oplus 4}$  that maps  $D_8$  to an ample class. There are 16  $(-2)$ -curves  $A_1, \dots, A_{16}$  on  $X$ ; all have degree 2 with respect to  $D_8$ . The class  $D_8$  is base-point free and not hyperelliptic; moreover, the linear system  $|D_8|$  embeds  $X$  as a degree 8 surface in  $\mathbb{P}^5$ . One can describe the configuration of the  $(-2)$ -curves on  $X$  as follows:

Let  $p_1, \dots, p_5$  be five points in general position in  $\mathbb{P}^2$  (by which we mean that no three are on a line), and let  $C_6$  be a sextic curve that has only nodal singularities at each point  $p_j$ . Let  $\pi: X \rightarrow \mathbb{P}^2$  be the minimal desingularization of the double cover  $Y \rightarrow \mathbb{P}^2$  branched over  $C_6$ . This is a smooth K3 surface which contains the following 16  $(-2)$ -curves:

- the  $(-2)$ -curves  $A_1, \dots, A_5$  above the points  $p_1, \dots, p_5$ ,
- the strict transform  $A_{ij}$  in  $X$  of the line through  $p_i$  and  $p_j$  for  $1 \leq i < j \leq 5$ ,
- the strict transform  $A_0$  of the unique conic passing through the five points  $p_1, \dots, p_5$ .

From that description, one understands easily the configuration of the 16  $(-2)$ -curves  $A_j$ ,  $0 \leq j \leq 5$ , and  $A_{ij}$ ,  $1 \leq i < j \leq 5$ , on  $X$ . The pull-backs of a line  $L$  and the  $(-2)$ -curves  $A_0, \dots, A_5$  generate a lattice which is

$$[2] \oplus \mathbf{A}_1^{\oplus 5} \simeq U(2) \oplus \mathbf{A}_1^{\oplus 4}.$$

The involution from the double cover fixes a smooth curve of genus  $10 - 5 = 5$ .

Let  $p_1, \dots, p_5$  be five points in general position in the plane, and let  $C_6$  be a general sextic curve with nodes at the points  $p_i$ .


**Proposition 6.9.** *The double cover of the blow-up of  $\mathbb{P}^2$  at the points  $p_i$  branched over the strict transform of  $C_6$  is a K3 surface with  $\text{NS}(X) \simeq U(2) \oplus \mathbf{A}_1^{\oplus 4}$ . The moduli space of these K3 surfaces is therefore unirational.*

### 6.8. The lattice $U \oplus \mathbf{A}_1^{\oplus 4}$

Let  $X$  be a K3 surface with Néron-Severi lattice

$$\text{NS}(X) = U \oplus \mathbf{A}_1^{\oplus 4}.$$

The surface contains nine  $(-2)$ -curves  $A_1, \dots, A_9$ , with dual graph



The  $(-2)$ -curves  $A_1, \dots, A_6$  generate the Néron-Severi lattice, and the classes of the remaining  $(-2)$ -curves are

$$A_7 \equiv A_2 + A_3 - A_6, \quad A_8 \equiv A_2 + A_3 - A_5, \quad A_9 \equiv A_2 + A_3 - A_4.$$

The class

$$D_{16} = 3A_1 + 4A_2 + 3A_3 + A_4 + A_5 + A_6$$

is ample and of square 16. The divisors

$$E_1 = A_2 + A_3, A_4 + A_9, A_5 + A_8, A_6 + A_7$$

are the reducible fibers of an elliptic fibration such that  $A_1$  is a section since one has

$$A_j \cdot A_1 = 1 \text{ for } j \in \{2, 4, 5, 6\}.$$

The class

$$E_2 = 2A_1 + A_2 + A_4 + A_5 + A_6$$

is the unique reducible fiber of another fibration. The effective divisor

$$D_2 = 2A_1 + A_2 + A_3 + A_4 + A_5 + A_6$$

is nef, of square  $D_2^2 = 2$ . The system  $|D_2|$  is base-point free, and  $D_2 \cdot A_j = 0$  if and only if  $j \in \{1, 3, 4, 5, 6\}$ ; else  $D_2 \cdot A_j = 2$ . Let  $\pi: X \rightarrow \mathbb{P}^2$  be the double cover map associated to  $|D_2|$ . Since  $D_2 \cdot A_j = 0$  for  $j \in \{1, 3, 4, 5, 6\}$ , the image of  $A_2$  by the double cover map is a line  $L_2$  such that  $D_2 = \pi^*L_2$ . The intersection matrix of the curves  $A_j$  for  $j \in \{1, 3, 4, 5, 6\}$  reveals that the sextic branch curve has a  $\mathbf{d}_4$  singularity and a node  $\mathbf{a}_1$ , and  $L_2$  is the line through these two singularities. The node is resolved on the double cover by  $A_3$ ; the  $\mathbf{d}_4$  singularity is resolved by the union of the curves  $A_1, A_4, A_5, A_6$  with  $A_1 \cdot A_4 = A_1 \cdot A_5 = A_1 \cdot A_6 = 1$ . We have

$$\begin{aligned} D_2 &\equiv 2A_1 + A_4 + A_5 + 2A_6 + A_7, \\ D_2 &\equiv 2A_1 + A_4 + 2A_5 + A_6 + A_8, \\ D_2 &\equiv 2A_1 + 2A_4 + A_5 + A_6 + A_9; \end{aligned}$$

thus the images of  $A_7, A_8, A_9$  are the three tangent lines  $L_7, L_8, L_9$  of the  $\mathbf{d}_4$  singularity.

Let  $C_6$  be a general sextic plane curve with a  $\mathbf{d}_4$  singularity and a node. We denote by  $L_2$  the line through the node and the  $\mathbf{d}_4$  singularity. Let  $L_7, L_8, L_9$  be the three lines tangent to the sextic at the  $\mathbf{d}_4$  singularity. Let  $Z \rightarrow \mathbb{P}^2$  be the embedded desingularization of  $C_6$ : this is the blow-up of  $\mathbb{P}^2$  at the node of  $C_6$  (with the exceptional divisor denoted by  $L_3$ ), and over the  $\mathbf{d}_4$  singularity, there are four blow-ups, producing four exceptional curves  $L_1, L_4, L_5, L_6$  such that

$$L_1^2 = -4, \quad L_1 \cdot L_j = 1, \quad L_j^2 = -1, \quad L_j \cdot L_k = 0 \quad \forall j, k \in \{4, 5, 6\} \text{ with } j \neq k.$$

The strict transforms of the  $L_j$  ( $j \in \{2, 7, 8, 9\}$ ) are disjoint curves  $\bar{L}_j$ , and (up to re-ordering)

$$L_4 \cdot \bar{L}_9 = 1, \quad L_5 \cdot \bar{L}_8 = 1, \quad L_6 \cdot \bar{L}_7 = 1.$$

From the above discussion, we obtain the following.


**Proposition 6.10.** *The curve  $\bar{C}_6 + L_1$  is the branch locus of a double cover  $Y \rightarrow Z$ . The surface  $Y$  is a smooth K3 surface, and the pull-backs of the curves  $L_1, \bar{L}_2, L_3, L_4, L_5, L_6, \bar{L}_7, \bar{L}_8, \bar{L}_9$  are  $(-2)$ -curves  $A_1, \dots, A_9$  such that the lattice generated by these curves is  $U \oplus \mathbf{A}_1^{\oplus 4}$ .*

### 6.9. The lattice $U(2) \oplus \mathbf{D}_4$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U(2) \oplus \mathbf{D}_4.$$

The surface  $X$  contains six  $(-2)$ -curves, with dual graph



These six  $(-2)$ -curves generate the Néron–Severi lattice. The class

$$D_{22} = 4A_6 + 3 \sum_{j=1}^6 A_j$$

is ample of square 22, and the curves  $A_j$  have degree 1 for that polarization.

For  $j \in \{1, \dots, 5\}$ , let  $F_j = A_6 + \sum_{k \neq j} A_k$ ; one has  $F_j^2 = 0$ , and  $F_j$  is a singular fiber of type  $I_0^*$  of an elliptic fibration  $\phi_j$ . Moreover,  $F_j \cdot A_j = 2$ , so that there are no sections.

The class

$$D_2 = A_6 + \sum_{j=1}^6 A_j$$

has square 2, with  $D_2 \cdot A_j = 0$  if and only if  $j \in \{1, \dots, 5\}$  and  $D_2 \cdot A_6 = 1$ . By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 6.11.** *The K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve which is the union of a line  $L$  and a smooth quintic  $Q$  such that  $L \cup Q$  has normal crossings.*


The divisor  $D_2$  is the pull-back of the line  $L$ . The six  $(-2)$ -curves on  $X$  come from the pull-backs of the line and the five exceptional divisors above the nodes. Conversely, the six  $(-2)$ -curves on a K3 surface which is the minimal desingularization of the double cover of the plane branched over the union of a line and a quintic necessarily have the same dual graph. The moduli space of K3 surfaces with  $\text{NS}(X) = U(2) \oplus \mathbf{D}_4$  is unirational.

### 6.10. The lattice $U \oplus \mathbf{D}_4$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U \oplus \mathbf{D}_4.$$

The dual graph of the six  $(-2)$ -curves on  $X$  is



The divisor

$$D_{70} = 21A_1 + 10(A_2 + A_3 + A_4) + 13A_5 + 6A_6$$

is ample, of square 70; the curves  $A_1, \dots, A_6$  have degree 1 for  $D_{70}$ . The divisor

$$F = 2A_1 + A_2 + A_3 + A_4 + A_5$$

is a fiber of type  $I_0^*$  of an elliptic fibration of  $X$  for which  $A_6$  is the unique section. The divisor

$$D_2 = 2F + A_6$$

is nef, of square 2, and has base points. The divisor  $D_8 = 2D_2$  is base-point free and hyperelliptic.

**Proposition 6.12.** *The linear system  $|D_8|$  defines a double cover  $\varphi: X \rightarrow \mathbf{F}_4$ , where the branch locus is the disjoint union of the unique section  $s$  with  $s^2 = -4$  and  $B' \in |3s + 12f|$ . The curve  $B'$  has a  $\mathbf{d}_4$  singularity  $q$ ; the curves  $A_1, A_2, A_3, A_4$  are contracted to  $q$  by  $\varphi$ ; the image of the curve  $A_5$  is the fiber through  $q$ .*

*Proof.* We apply Theorem 2.3, case a) i). □


## 7. Rank 7 lattices

### 7.1. The lattice $U \oplus D_4 \oplus A_1$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U \oplus D_4 \oplus A_1.$$

There exist eight  $(-2)$ -curves  $A_1, \dots, A_8$  on  $X$  with dual graph



The divisor

$$D_{62} = (8, 17, 8, 8, 11, 6, 2)$$

in the basis  $A_1, \dots, A_7$  is ample, of square 62. The classes

$$F_1 = A_1 + 2A_2 + A_3 + A_4 + A_5, \quad F_2 = A_7 + A_8$$

are singular fibers of type  $I_0^*$  and  $I_2$  or  $III$ , respectively, of an elliptic fibration for which  $A_6$  is a section (that determines the class of  $A_8$  in  $\text{NS}(X)$ ). The divisor

$$D_2 = A_1 + 4A_2 + 2A_3 + 2A_4 + 3A_5 + 2A_6 + A_7$$

is base-point free, of square 2, with  $D_2 \cdot A_k = 0$  if and only if  $k \in \{2, 3, 4, 5, 6, 7\}$ ,  $D_2 \cdot A_1 = D_2 \cdot A_8 = 2$  and

$$D_2 \equiv 2A_2 + A_3 + A_4 + 2A_5 + 2A_6 + 2A_7 + A_8.$$


**Proposition 7.1.** *The linear system  $|D_2|$  defines a double cover of the plane  $\pi: X \rightarrow \mathbb{P}^2$  branched over a sextic curve  $C_6$  with a  $\mathbf{d}_6$  singularity. The image of the curve  $A_1$  is the line that is tangent to two branches of the  $\mathbf{d}_6$  singularity, and the image of  $A_8$  is tangent to the third branch.*

### 7.2. The lattice $U \oplus A_1^{\oplus 5}$

Let  $X$  be a K3 surface with a Néron–Severi lattice

$$\text{NS}(X) = U \oplus A_1^{\oplus 5}.$$

There exist 12  $(-2)$ -curves  $A_1, \dots, A_{12}$  on  $X$ , with dual graph



The divisor

$$D_{14} = 2A_1 + 2A_2 + A_7 + \sum_{i=1}^7 A_i$$

is ample of square 14. The divisors

$$A_{2+j} + A_{7+j}, \quad j \in \{0, 1, 2, 3, 4\},$$

are irreducible fibers of an elliptic fibration  $\phi_0$  of  $X$ . The curve  $A_1$  is a section of  $\phi_0$ , and the curve  $A_{12}$  (of class  $(2, 0, 1, 1, 1, 1, -1)$ ) is a 2-section.

The divisor

$$D_2 = 2A_1 + \sum_{j=2}^6 A_j$$

has square 2, and  $|D_2|$  is base-point free. It defines a 2-to-1 cover  $\pi: X \rightarrow \mathbb{P}^2$  of the plane branched over a sextic curve  $C_6$ . For the curves  $A_j$ ,  $j \in \{2, 3, 4, 5, 6, 12\}$ , one has  $D_2 \cdot A_j = 0$ ; thus the images of these disjoint curves are six points in  $\mathbb{P}^2$ , and the sextic has nodes at these points. We have  $D_2 \cdot A_1 = 1$  and

$$D_2 \cdot A_7 = D_2 \cdot A_8 = D_2 \cdot A_9 = D_2 \cdot A_{10} = D_2 \cdot A_{11} = 2.$$

Since  $D_2 = 2A_1 + \sum_{j=2}^6 A_j$ , there is a line  $L_1$  in the branch locus such that  $D_2 = \pi^*L_1$ , and  $C_6$  is the union of  $L_1$  and a nodal quintic. Conversely, we have the following.

**Proposition 7.2.** *The minimal resolution of a surface which is the double cover of the plane branched over the union of a line and a general quintic with a node is a K3 surface of type  $U \oplus \mathbf{A}_1^{\oplus 5}$ .*

Such surfaces are studied in [AK11, Section 3.3]. Clearly, the moduli space of K3 surfaces with  $\text{NS}(X) = U \oplus \mathbf{A}_1^{\oplus 5}$  is unirational.

### 7.3. The lattice $U(2) \oplus \mathbf{A}_1^{\oplus 5}$ and cubic surfaces

Let  $C_6$  be a sextic curve in  $\mathbb{P}^2$  with six nodes in general position. Let  $Z \rightarrow \mathbb{P}^2$  be the blow-up of the nodes; it is a degree 3 del Pezzo surface and contains 27  $(-1)$ -curves:

- the 6 exceptional divisors  $E_i$ ,  $i = 1, \dots, 6$ ,
- the strict transforms  $L_{ij}$  of the 15 lines through  $p_i, p_j$  ( $i \neq j$ ),
- the strict transforms  $Q_j$  of the 6 conics that go through points in  $\{p_1, \dots, p_6\} \setminus \{p_j\}$ .

The Néron–Severi lattice of  $Z$  is generated by the pull-back  $L'$  of a line and  $E_1, \dots, E_6$ ; it is the unimodular rank 7 lattice

$$I_1 \oplus I_{-1}^{\oplus 6}.$$

The anti-canonical divisor of the degree 3 del Pezzo surface  $Z$  is given by

$$-K_Z = 3L' - (E_1 + \dots + E_6);$$

this is an ample divisor. The linear system  $|-K_Z|$  is base-point free (see [Dem76, Section 3, Theorem 1]), and each of the 30 divisors

$$Q_j + L_{ij} + E_i \quad \text{with } i \neq j, \quad i, j \in \{1, \dots, 6\}$$

and 15 divisors

$$L_{ij} + L_{kl} + L_{mn} \quad \text{with } \{i, j, k, l, m, n\} = \{1, \dots, 6\}$$

belongs to the linear system  $|-K_Z|$ .

The double cover  $f: Y \rightarrow Z$  branched over the strict transform  $C'_6$  of  $C_6$  is a smooth K3 surface. The pull-backs of the 27  $(-1)$ -curves are  $(-2)$ -curves. We denote by  $A_1, \dots, A_6$  the pull-backs on  $Y$  of the curves  $E_i$  and by  $L$  the pull-back of  $L'$ . Naturally, the lattice  $f^*\text{NS}(Z)$  is  $(I_1 \oplus I_{-1}^{\oplus 6})(2)$ , which is also the lattice generated by  $L, A_1, \dots, A_6$  and is isometric to  $U(2) \oplus \mathbf{A}_1^{\oplus 5}$ .

Since  $f: Y \rightarrow Z$  is finite, its pull-back  $D_6 = f^*(-K_Z)$  is ample, base-point free, non-hyperelliptic, with  $D_6^2 = 6$ . We thus have the following.

**Proposition 7.3.** *The image of  $Y$  under the map  $\pi: Y \rightarrow \mathbb{P}^4$  obtained from  $|D_6|$  is a degree 6 complete intersection surface. There exist 45 hyperplane sections of  $Y$  which are each the union of 3 conics.*


These 45 hyperplane sections correspond to the 45 tritangent planes of a cubic, *i.e.*, to the planes containing 3 lines in the cubic. In fact, by the above discussion, the strict transform in the cubic surface  $Z$  of  $C_6$  is a degree 6 curve in  $\mathbb{P}^3$  which is the complete intersection of the cubic surface  $Y = \{f_3(x, y, z, t) = 0\}$  and a quadric  $\{q_2(x, z, y, t) = 0\}$ . The K3 surface  $Y$  is the complete intersection of  $\{f_3(X, Y, Z, T) = 0\}$  and the quadric  $\{q_2(X, Y, Z, T) - W^2 = 0\}$  in  $\mathbb{P}^4$  (with coordinates  $X, Y, Z, T, W$ ). From that discussion, we see that the moduli space of K3 surfaces with  $\text{NS}(X) \simeq U(2) \oplus \mathbf{A}_1^{\oplus 5}$  is unirational.

#### 7.4. The lattice $U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2^{\oplus 2}$

Let  $X$  be a K3 surface with Néron–Severi lattice

$$\text{NS}(X) = U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2^{\oplus 2}.$$

The dual graph of the nine  $(-2)$ -curves on  $X$  is



The curves  $A_1, A_2, A_3, A_4, A_5, A_7, A_8$  generate the Néron–Severi lattice. The divisor

$$D_{18} = (5, 5, 6, -1, -1, 3, 1)$$

in the above basis is ample, of square 18. One has  $D_{18} \cdot A_j = 1$  for  $j < 8$  and  $D_{18} \cdot A_9 = 2$ . The divisors

$$F = A_1 + A_2 + A_3, A_4 + A_5 + A_6, A_8 + A_9$$

are the reducible fibers of an elliptic fibration of the surface. The divisor

$$D_2 = 2F + A_7$$

is nef, of square 2, with base points and  $D_2 \cdot A_3 = D_2 \cdot A_6 = D_2 \cdot A_8 = 1$ ,  $D_2 \cdot A_j = 0$  for  $j \neq 0$ . By Theorem 2.3, case a) i), the base-point free linear system  $|2D_2|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  which is branched over the unique section  $s$  of  $\text{mathbf{F}}_4$  with  $s^2 = -4$  and a disjoint curve  $B \in |2s + 12f|$ . The curve  $B$  has a node  $q$  (which is the image of  $A_9$ ) and two  $\mathbf{a}_2$  singularities  $p, p'$  which are the images of  $A_1, A_2$  and  $A_4, A_5$ . The images of  $A_3, A_6, A_8$  are the fibers through  $p, p'$  and  $q$ , respectively.

One can give another description as follows. The divisor

$$D'_2 = A_9 + (A_3 + A_6 + 2A_7 + 2A_8)$$

is nef of square 2, and  $|D'_2|$  is base-point free. Moreover,  $D_2 \cdot A_j = 0$  if and only if  $j \in \{3, 6, 7, 8\}$ , and  $D_2 \cdot A_9 = 2$  and  $D_2 \cdot A_j = 1$  for  $j \in \{1, 2, 4, 5\}$ . We have

$$\begin{aligned} D'_2 &\equiv A_1 + A_2 + (2A_3 + A_6 + 2A_7 + A_8), \\ D'_2 &\equiv A_4 + A_5 + (A_3 + 2A_6 + 2A_7 + A_8). \end{aligned}$$

By using the linear system  $|D'_2|$ , we obtain the following.


**Proposition 7.4.** *The K3 surface  $X$  is the double cover  $\eta: X \rightarrow \mathbb{P}^2$  of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with a  $\mathbf{d}_4$  singularity  $q$  onto which the curves  $A_3, A_6, A_7, A_8$  are contracted.*

The image of the curve  $A_9$  is a line that is tangent to a branch of the singularity  $q$  and that meets the sextic in two other points. The two lines  $L_1, L_2$  that are tangent to the two other branches of  $q$  are tangent to another point of the sextic. The image of  $A_1, A_2$  is  $L_1$ , and the image of  $A_4, A_5$  is  $L_2$ .



### 7.5. The lattice $U \oplus A_1^{\oplus 2} \oplus A_3$

The dual graph of the nine  $(-2)$ -curves  $A_1, \dots, A_9$  on  $X$  is



The curves  $A_1, \dots, A_7$  generate the Néron-Severi lattice. The divisor

$$D_{34} = (6, 4, 2, 5, 3, 1, 1)$$

in the basis  $A_1, \dots, A_7$  is ample of square 34. The divisors

$$F_1 = A_1 + A_2, \quad F_2 = A_3 + A_9, \quad F_3 = A_5 + A_6 + A_7 + A_8$$

are the reducible fibers of an elliptic fibration  $\phi_1$  of  $X$  for which  $A_4$  is a section (for that, one can deduce the classes of curves  $A_8, A_9$ ). The divisor


$$D_2 = 2F_1 + A_4$$

is nef, of square 2, with base points. By Theorem 2.3, case i) a), we have the following.

**Proposition 7.5.** *The linear system  $|D_2|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has two nodes  $p, p'$  and an  $\mathfrak{a}_3$  singularity  $q$ . The pull-backs of the fibers through  $p, p', q$  are the fibers  $F_1, F_2, F_3$ .*

### 7.6. The lattice $U \oplus A_2 \oplus A_3$

The dual graph of the eight  $(-2)$ -curves  $A_1, \dots, A_8$  on  $X$  is



The seven  $(-2)$ -curves  $A_1, \dots, A_7$  generate  $\text{NS}(X)$ , and the divisor

$$D_{42} = (6, 6, 8, 5, 3, 1, 1)$$

in the basis  $A_1, \dots, A_7$  is ample, of square 42. The surface has an elliptic fibration  $\phi_1$  with reducible fibers the curves

$$F_1 = A_1 + A_2 + A_3, \quad F_2 = A_5 + A_6 + A_7 + A_8,$$

and  $A_4$  is a section. The divisor


$$D_2 = 2F_1 + A_4$$

is nef, of square 2, with base points. By Theorem 2.3, case i) a), we have the following.

**Proposition 7.6.** *The linear system  $|D_2|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a cusp  $p$  and an  $\mathfrak{a}_3$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 7.7. The lattice $U \oplus A_1 \oplus A_4$

The dual graph of the eight  $(-2)$ -curves  $A_1, \dots, A_8$  on  $X$  is



In the basis  $A_1, \dots, A_7$ , the divisor

$$D_{42} = (7, 9, 5, 2, 0, -1, -1)$$

is ample, of square 42. The curves

$$F_1 = A_1 + A_2, \quad F_2 = A_4 + A_5 + A_6 + A_7 + A_8$$

are the reducible fibers of an elliptic fibration for which  $A_3$  is a section. The divisor


$$D_2 = 2F_1 + A_4$$

is nef, of square 2, with base points. By Theorem 2.3, case i) a), we have the following.

**Proposition 7.7.** *The linear system  $|2D_2|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and an  $\mathbf{a}_4$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 7.8. The lattice $U \oplus A_5$

The dual graph of the seven  $(-2)$ -curves  $A_1, \dots, A_7$  on  $X$  is



The seven  $(-2)$ -curves  $A_1, \dots, A_7$  generate the Néron–Severi lattice, and in that basis, the divisor

$$D_{84} = (7, 15, 12, 10, 9, 10, 12)$$

is ample, of square 84. We have  $D_{84} \cdot A_j = 1$  for  $j \in \{1, \dots, 7\} \setminus \{5\}$  and  $D_{84} \cdot A_5 = 2$ . The divisor

$$F = \sum_{j=2}^7 A_j$$

is the unique reducible fiber of an elliptic fibration of  $X$ ; the curve  $A_1$  is a section.

The divisor

$$D'_2 = (2, 4, 3, 2, 1, 2, 3)$$


is nef, base-point free, of square 2. One has  $D_2 \cdot A_j = 0$  for  $j \in \{1, \dots, 7\} \setminus \{5\}$  and  $D_2 \cdot A_5 = 2$ . The surface  $X$  is a double cover of  $\mathbb{P}^2$ . The branch locus is a sextic curve with an  $\mathbf{e}_6$  singularity; there exists a line  $L$  through the  $\mathbf{e}_6$  singularity which cuts the sextic transversely in two points such that the image of  $A_5$  is  $L$ .

One may also describe that surface as a double cover of  $\mathbf{F}_4$ .

**Proposition 7.8.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $\mathbf{a}_5$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

### 7.9. The lattice $U \oplus D_5$

The dual graph of the seven  $(-2)$ -curves  $A_1, \dots, A_7$  on  $X$  is



The divisor

$$D_{114} = (8, 17, 27, 13, 25, 12, 12)$$

in the basis  $A_1, \dots, A_7$  is ample, of square 114, and every  $(-2)$ -curve on  $X$  has degree 1 with respect to  $D_{114}$ . The divisor

$$F = A_2 + 2A_3 + A_4 + 2A_5 + A_6 + A_7$$


is the unique reducible fiber of an elliptic fibration of  $X$  for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 7.9.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_5$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

## 8. Rank 8 lattices

### 8.1. The lattice $U \oplus D_6$

The dual graph of the eight  $(-2)$ -curves  $A_1, \dots, A_8$  on  $X$  is



The divisor

$$D_{220} = (11, 24, 18, 38, 35, 33, 16, 16)$$

in the basis  $A_1, \dots, A_8$  is ample, of square 220, with  $D_{220} \cdot A_j = 1$  if  $j \notin \{1, 3\}$  and  $D_{220} \cdot A_j = 2$  if  $j \in \{1, 3\}$ . The divisor


$$F = A_2 + A_3 + 2A_4 + 2A_5 + 2A_6 + A_7 + A_8$$

is a fiber of an elliptic fibration for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.1.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_6$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

### 8.2. The lattice $U \oplus D_4 \oplus A_1^{\oplus 2}$

The dual graph of the 10  $(-2)$ -curves  $A_1, \dots, A_{10}$  on  $X$  is



The first eight curves generate the Néron-Severi lattice. The divisor

$$D_{54} = (6, 13, 6, 6, 9, 6, 2, 2)$$

of square 54 is ample with  $D_{54} \cdot A_j = 1$  for  $j \leq 6$ ,  $D_{54} \cdot A_j = 2$  for  $j \in \{7, 8\}$  and  $D_{54} \cdot A_j = 4$  for  $j \in \{9, 10\}$ . The divisors

$$F_1 = A_2 + \sum_{j=1}^5 A_j, \quad F_2 = A_7 + A_9, \quad F_3 = A_8 + A_{10}$$

are fibers of an elliptic fibration  $\varphi_2$  such that  $A_6$  is a section. For  $j \in \{1, 3, 4\}$ , the divisors

$$-A_j + A_2 + A_5 + A_6 + \sum_{k=1}^8 A_k$$

are fibers of elliptic fibrations  $\varphi_j$  for which  $A_j$  and  $A_9, A_{10}$  are 2-sections. The divisor

$$D_2 = A_1 + 2A_2 + A_3 + A_4 + 2A_5 + 2A_6 + A_7 + A_8$$

is nef, of square 2 and base-point free. The linear system  $|D_2|$  contracts the curves  $A_j$  for  $j \in \{1, 3, \dots, 8\}$ ; moreover,  $D_2 \cdot A_2 = 1$  and  $D_2 \cdot A_9 = D_2 \cdot A_{10} = 2$ .

**Proposition 8.2.** *The branch curve is the union of a line  $L$  and a nodal quintic  $Q$ ; the line goes through the node transversely (forming a  $\mathbf{d}_4$  singularity on the sextic  $L \cup Q$ , resolved by  $A_5, A_6, A_7, A_8$ ) and cuts the quintic in three other points (resolved by  $A_1, A_3, A_4$ ).*

We have  $D_2 = \pi^*L$ , and the equivalences

$$D_2 \equiv A_5 + 2A_6 + A_8 + A_9$$

and


$$D_2 \equiv A_5 + 2A_6 + A_7 + A_{10}$$

give that the images of  $A_9$  and  $A_{10}$  are lines that are tangent to the branches of the node of the quintic  $Q$ .

*Remark 8.3.* One may also describe that surface as the desingularization of the double cover of  $\mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$  with a  $\mathbf{d}_4$  singularity and two nodes.

### 8.3. The lattice $U \oplus A_1^{\oplus 6}$

Let  $X$  be a K3 surface with  $\text{NS}(X) \simeq U \oplus A_1^{\oplus 6}$ . The surface  $X$  contains 19  $(-2)$ -curves; their dual graph is represented by



where we draw the graph of the configuration of the curves  $A_8, \dots, A_{19}$  in another part in order for it to be more readable. The first eight curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. The divisor

$$D_{12} = 3A_1 + 2A_2 + A_3 + A_4 + A_5 + A_6 + A_7 + A_8$$

is ample, of square 12, with  $D_{12} \cdot A_j = 1$  for  $j \leq 7$ ,  $D_{12} \cdot A_j = 2$  for  $8 \leq j \leq 13$  and  $D_{12} \cdot A_j = 4$  for  $14 \leq j \leq 19$ . Let  $j \neq k$  be two elements of  $\{2, \dots, 7\}$ ; the effective divisor

$$F_{jk} = 2A_1 - A_j - A_k + \sum_{i=2}^7 A_i$$

is a fiber of type *IV* of an elliptic fibration  $\varphi_{jk}$ . The divisors

$$A_{j+6} + A_{k+12}, A_{k+6} + A_{j+12}$$

are the other reducible fibers of that fibration. The divisors

$$A_{2+k} + A_{8+k}, \quad k \in \{0, \dots, 5\}$$

are reducible fibers of an elliptic fibration  $\varphi_1$ . The divisors

$$A_{2+k} + A_{14+k}, \quad k \in \{0, \dots, 5\}$$

are reducible fibers of an elliptic fibration  $\varphi_2$ . The curve  $A_1$  is a section of  $\varphi_1$  and  $\varphi_2$ . The divisor

$$D_2 = 2A_1 + A_2 + A_3 + A_4 + A_5 + A_6$$

has square 2; it is base-point free, and it contracts  $A_2, \dots, A_6, A_{13}, A_{19}$ . Moreover,  $D_2 \cdot A_1 = 1$ , and for the remaining curves,  $D_2 \cdot A_j = 2$ . From the equivalence relations obtained by the elliptic fibration with fiber  $A_2 + A_8$  and the elliptic fibration  $\varphi_{27}$ , we obtain

$$D_2 \equiv A_j + A_{j+6} + A_{19} \equiv A_j + A_{12+6} + A_{13}, \quad \forall j \in \{2, \dots, 7\}.$$

Thus we see that there is a line  $L$  in the branch locus which is the image of  $A_1$ , and the curves  $A_2, \dots, A_6$  are mapped to points  $p_2, \dots, p_6$  in the intersection of  $L$  with residual quintic  $Q$  in the branch locus. The curve  $Q$  has two nodes  $p, q$  above which are  $A_{13}$  and  $A_{19}$ . The curve  $A_7$  is the strict transform of the line through  $p$  and  $q$ . For  $j \in \{2, 3, 4, 5, 6\}$ , the curve  $A_{j+6}$  (resp.  $A_{j+12}$ ) is the pull-back of the line through  $p, p_j$  (resp.  $q, p_j$ ). This leads to the following.

**Proposition 8.4.** *The K3 surface  $X$  is the minimal resolution of the double cover of  $\mathbb{P}^2$  branched over the union of a line and a quintic with two nodes.*

From that description, the moduli space of K3 surfaces  $X$  with  $\text{NS}(X) \simeq U \oplus \mathbf{A}_1^{\oplus 6}$  is unirational.

### 8.4. The lattice $U(2) \oplus \mathbf{A}_1^{\oplus 6}$ and degree 2 del Pezzo surfaces

Let  $C_6$  be a sextic curve in  $\mathbb{P}^2$  with nodes through seven points  $p_1, \dots, p_7$  in general position. Let  $Z \rightarrow \mathbb{P}^2$  the blow-up of the nodes; it is a degree 2 del Pezzo surface, and it contains 56  $(-1)$ -curves:

- the 7 exceptional divisors  $E_i$ ,  $i = 1, \dots, 7$ ,
- the strict transforms  $L_{ij}$  of the 21 lines through  $p_i, p_j$  ( $i \neq j$ ),
- the strict transforms  $Q_{rs}$  of the 21 conics that go through points in  $\{p_1, \dots, p_7\} \setminus \{p_r, p_s\}$ ,
- the strict transforms  $CU_j$  of the 7 cubics that go through the 7 points  $p_k$ , with a double point at one of these points  $p_j$ .

The Néron–Severi lattice of  $Z$  is generated by the pull-back  $L'$  of a line and  $E_1, \dots, E_7$ ; it is the unimodular rank 7 lattice

$$I_1 \oplus I_{-1}^{\oplus 7}.$$

The anti-canonical divisor of the del Pezzo surface  $Z$  of degree 2 is given by

$$-K_Z = 3L' - (E_1 + \dots + E_7);$$

this is an ample divisor. The linear system  $|-K_Z|$  is base-point free (see [Dem76, Section 3, Theorem 1]), and we remark that each of the 28 divisors

$$E_i + CU_i, L_{ij} + Q_{ij}, \quad i, j \in \{1, \dots, 7\}, i \neq j$$

belongs to the system  $|-K_Z|$ .

Let  $Y \rightarrow Z$  be the double cover branched over the strict transform  $C'$  of  $C_6$  in  $Z$ . Since  $C' \equiv -2K_Z$ , the surface  $Y$  is a smooth K3 surface. Since for any  $(-1)$ -curve  $B$ ,  $C'B = -2K_Z B = 2$ , the pull-backs on  $Y$  of the curves  $E_i, L_{ij}, Q_{rs}, CU_j$  are  $(-2)$ -curves. We denote these curves by  $A_i, B_{ij}, C_{rs}, D_j$ , respectively. Let  $L$  be the pull-back in  $Y$  of  $L'$ . The lattice generated by  $L, A_1, \dots, A_7$  is (isometric to)  $U(2) \oplus \mathbf{A}_1^{\oplus 6} = f^* \text{NS}(Z) = (I_1 \oplus I_{-1}^{\oplus 7})(2)$ . Since  $f: Y \rightarrow Z$  is finite and the linear system  $|-K_Z|$  is ample and base-point free, its pull-back  $D_4 = f^*(-K_Z)$  is ample and base-point free, with  $D_4^2 = 4$ . The linear system  $|D_4|$  is 3-dimensional, and the invariant part is the pencil  $f^*|-K_Z|$ . We thus have the following.

**Proposition 8.5.** *The image of  $Y$  under the map  $\pi: Y \rightarrow \mathbb{P}^3$  induced from  $|D_4|$  is a smooth quartic surface. There exist 28 hyperplane sections of  $Y$  which are each the union of 2 conics.*


*Remark 8.6.* The moduli space of K3 surfaces of type  $U(2) \oplus \mathbf{A}_1^{\oplus 6}$  is 12-dimensional. In [Kon00], Kondo studies the 6-dimensional moduli space of curves of genus 3 via the periods of the K3 surfaces  $X = \{t^4 - f_4(x, y, z) = 0\}$  which are quadruple covers of the plane branched over smooth quartic curves  $C = \{f_4(x, y, z) = 0\}$  (thus of genus 3). The double cover of  $\mathbb{P}^2$  branched over  $C$  is a degree 2 del Pezzo surface. For such K3 surfaces, the pull-backs of the 28 bitangents of the curve  $C$  give the 28 hyperplane sections of Theorem 8.5.

**Proposition 8.7.** *The moduli space of  $U(2) \oplus \mathbf{A}_1^{\oplus 6}$ -polarized K3 surfaces is unirational.*

*Proof.* Seven points  $p_1, \dots, p_7$  in general position in  $\mathbb{P}^2$  form an open subscheme of  $(\mathbb{P}^2)^7$ . The space of sextics with the points  $p_1, \dots, p_7$  as seven nodes is a 14-dimensional linear subspace of the space of sextics. Therefore, the moduli space of K3 surfaces that are double covers of  $\mathbb{P}^2$  branched over a 7-nodal sextic is unirational.  $\square$

### 8.5. The lattice $U \oplus \mathbf{A}_2^{\oplus 3}$

The K3 surface contains 10  $(-2)$ -curves  $A_1, \dots, A_{10}$  with dual graph



The first eight curves generate the Néron–Severi lattice. The divisor

$$D_{18} = 3A_1 + 5A_2 + A_3 + A_4 + 4A_5 + 4A_6$$

is ample, of square 18, with  $D_{18} \cdot A_j = 1$  for  $j \in \{1, \dots, 10\}$ . The divisors

$$A_2 + A_5 + A_6, A_3 + A_7 + A_9, A_4 + A_8 + A_{10}$$

are fibers of an elliptic fibration of the K3 surface such that  $A_1$  is a section. For  $i \in \{5, 6\}$ ,  $j \in \{7, 9\}$ ,  $k \in \{8, 10\}$ , the divisor

$$F_{ijk} = 3A_1 + 2A_2 + 2A_3 + 2A_4 + A_i + A_j + A_k$$

is the unique reducible fiber of type  $IV^*$  of an elliptic fibration  $\varphi_{ijk}$ . The divisor

$$D_2 = 2A_1 + 2A_2 + A_3 + A_4 + A_5 + A_6$$

is nef, base-point free, of square 2; it contracts  $A_1, A_2, A_3, A_4$ , and the other  $(-2)$ -curves have degree 1 for that divisor. We also have

$$D_2 \equiv 2A_1 + A_2 + 2A_3 + A_4 + A_7 + A_9, \\ D_2 \equiv 2A_1 + A_2 + A_3 + 2A_4 + A_8 + A_{10}.$$


By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 8.8.** *The surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve with a  $\mathbf{d}_4$  singularity. The three tangents to the three branches of the singularity are tangent to another point of the sextic curve, so that the pull-back of a tangent splits into two  $(-2)$ -curves. The  $6 = 3 \cdot 2$   $(-2)$ -curves above the tangent lines are  $A_5 + A_6$ ,  $A_7 + A_9$  and  $A_8 + A_{10}$ .*

One can also construct this surface as a double cover of  $\mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$  with three cusps singularities.

### 8.6. The lattice $U \oplus A_3^{\oplus 2}$

The K3 surface contains nine  $(-2)$ -curves; the dual graph is



The eight  $(-2)$ -curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. The divisor

$$D_{40} = (5, 6, 6, 8, 5, 3, 1, 1)$$

in the basis  $A_1, \dots, A_8$  is ample, of square 40, with  $D_{40} \cdot A_1 = D_{40} \cdot A_9 = 2$  and  $D_{40} \cdot A_j = 1$  for  $j \in \{2, \dots, 8\}$ . The divisors


$$F_1 = A_1 + A_2 + A_3 + A_4, \quad F_2 = A_6 + A_7 + A_8 + A_9$$

are reducible fibers of an elliptic fibration, where  $A_5$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.9.** *The linear system  $|4F_1 + 2A_5|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has two  $\mathbf{a}_3$  singularities  $p, q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 8.7. The lattice $U \oplus \mathbf{A}_2 \oplus \mathbf{A}_4$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



The curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. The divisor

$$D_{42} = (7, 6, 6, 7, 9, 5, 2, 0)$$

in the basis  $A_1, \dots, A_8$  is ample, of square 42, with  $D_{42} \cdot A_j = 1$  for  $j \in \{1, \dots, 7\}$  and  $D_{42} \cdot A_8 = D_{42} \cdot A_9 = 2$ . The divisors


$$F_1 = A_1 + \dots + A_5, \quad F_2 = A_7 + A_8 + A_9$$

are reducible fibers of an elliptic fibration for which  $A_6$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.10.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{a}_4$  singularity  $p$  and a cusp  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 8.8. The lattice $U \oplus \mathbf{A}_1 \oplus \mathbf{A}_5$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



The curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. The divisor

$$D_{76} = (10, 12, 7, 3, 0, -2, -3, -2)$$

in the basis  $A_1, \dots, A_8$  is ample, of square 76. The divisors

$$F_1 = A_1 + A_2, \quad F_2 = A_4 + A_5 + A_6 + A_7 + A_8 + A_9$$


are the reducible fibers of an elliptic fibration such that  $A_3$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.11.** *The linear system  $|4F_1 + 2A_3|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and one  $\mathbf{a}_5$  singularity  $q$ . The pull-back of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*



### 8.9. The lattice $U \oplus A_6$

The surface  $X$  contains eight  $(-2)$ -curves  $A_1, \dots, A_8$ , with dual graph



These eight curves generate the Néron–Severi lattice; the divisor

$$D_{84} = (7, 15, 12, 10, 9, 9, 10, 12)$$

in the basis  $A_1, \dots, A_8$  has square 84, and the curves  $A_j$  have degree 1 for  $D_{84}$ . The divisor

$$F = A_2 + \dots + A_8$$

is a fiber of an elliptic fibration for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.12.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $\mathbf{a}_6$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*


*Remark 8.13.* The divisor

$$D_2 = (2, 4, 3, 2, 1, 1, 2, 3)$$

in the basis  $A_1, \dots, A_8$  is base-point free of square 2. It defines a double cover branched over a sextic with an  $\mathbf{e}_6$  singularity.

### 8.10. The lattice $U \oplus A_2 \oplus D_4$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



The curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. In that basis, the divisor

$$D_{56} = (7, 7, 7, 15, 10, 6, 3, 1)$$

is ample of square 56, with  $D_{56} \cdot A_j = 1$  for  $j \in \{1, \dots, 8\}$  and  $D_{56} \cdot A_9 = 4$ . The divisors


$$F_1 = A_7 + A_8 + A_9, \quad F_2 = A_1 + A_2 + A_3 + A_5 + 2A_4$$

are fibers of an elliptic fibration for which  $A_6$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.14.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a cusp  $p$  and one  $\mathbf{d}_4$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 8.11. The lattice $U \oplus A_1 \oplus D_5$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



The curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. In that basis, the divisor

$$D_{96} = (9, 12, 8, 5, 3, 1, 1, 0)$$

is ample of square 96. We have  $D_{96} \cdot A_1 = 6$ ,  $D_{96} \cdot A_2 = 2$  and  $D_{96} \cdot A_j = 1$  for  $j \geq 3$ . The divisors


$$F_1 = A_1 + A_2, \quad F_2 = A_4 + A_6 + 2A_5 + 2A_7 + A_8 + A_9$$

are fibers of an elliptic fibration for which  $A_3$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.15.** *The linear system  $|4F_1 + 2A_3|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and one  $\mathbf{d}_5$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 8.12. The lattice $U \oplus E_6$

The surface  $X$  contains eight  $(-2)$ -curves, with dual graph



These eight curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. In that basis, the divisor

$$D_{234} = (12, 25, 39, 54, 35, 35, 17, 17)$$

is ample, of square 234, with  $D_{234} \cdot A_j = 1$  for  $j \in \{1, \dots, 8\}$ . The divisor

$$F = A_2 + 2A_3 + 3A_4 + 2A_5 + 2A_6 + A_7 + A_8$$


is a fiber of an elliptic fibration for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 8.16.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $\mathbf{e}_6$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

## 9. Rank 9 lattices

### 9.1. The lattice $U \oplus E_7$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



These nine curves  $A_1, \dots, A_9$  generate the Néron-Severi lattice. In that basis, the divisor

$$D_{532} = (19, 39, 60, 82, 105, 52, 77, 50, 24)$$

is ample, of square 532, with  $D_{532} \cdot A_j = 1$  for  $j \leq 8$  and  $D_{532} \cdot A_9 = 2$ . The divisor


$$F = A_2 + 2A_3 + 3A_4 + 4A_5 + 2A_6 + 3A_7 + 2A_8 + A_9$$

is a fiber of an elliptic fibration, for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 9.1.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $e_7$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

### 9.2. The lattice $U \oplus D_6 \oplus A_1$

The surface  $X$  contains 10  $(-2)$ -curves, with dual graph



The nine curves  $A_1, \dots, A_9$  generate the Néron-Severi lattice. In that basis, the divisor

$$D_{148} = (10, 15, 11, 8, 6, 2, 3, 1, 0)$$

is ample, of square 148, with  $D_{148} \cdot A_1 = 10$ ,  $D_{148} \cdot A_6 = 2$  and  $D_{148} \cdot A_j = 1$  for  $j \neq 1, 6$ . The divisors


$$F_1 = A_1 + A_2, \quad F_2 = A_4 + A_6 + 2A_5 + 2A_7 + 2A_8 + A_9 + A_{10}$$

are fibers of an elliptic fibration. By Theorem 2.3, case i) a), we have the following.

**Proposition 9.2.** *The linear system  $|4F_1 + 2A_3|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and one  $d_6$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 9.3. The lattice $U \oplus D_4 \oplus A_1^{\oplus 3}$

The surface  $X$  contains 15  $(-2)$ -curves  $A_1, \dots, A_{15}$ , with dual graph



The curves  $A_1, \dots, A_9$  generate the Néron-Severi lattice. In that basis, the divisor

$$D_{44} = (6, 7, 3, 3, 3, 7, 3, 3, 3)$$

is ample, of square 44, with  $D_{44} \cdot A_j = 1$  for  $j \in \{2, \dots, 9\}$ ,  $D_{44} \cdot A_1 = 2$  and  $D_{44} \cdot A_j = 6$  for  $j \in \{10, \dots, 15\}$ . The divisor

$$D_2 = (1, 2, 1, 1, 1, 2, 1, 1, 1)$$

is nef, base-point free, with  $D_2 \cdot A_j = 0$  for  $j \in \{2, \dots, 9\}$  and  $D_2 \cdot A_j = 2$  for  $j \notin \{2, \dots, 9\}$ . Let  $\eta: X \rightarrow \mathbb{P}^2$  be the map associated to the linear system  $|D_2|$ . One thus has the following.

**Proposition 9.3.** *The branch curve of  $\eta$  is a sextic with two  $\mathbf{d}_4$  singularities  $p, q$ . The image of the curve  $A_1$  by the double cover map is the line through  $p, q$ . The six lines  $C_{10}, \dots, C_{15}$  that are tangent to the six branches of the two  $\mathbf{d}_4$  singularities are the images of  $A_{10}, \dots, A_{15}$ .*

#### 9.4. The lattice $U \oplus \mathbf{A}_1^{\oplus 7}$

The K3 surface contains 37  $(-2)$ -curves, which we denote by  $A_0, A_1, \dots, A_8$  and  $A_{ij}$  for  $i, j \in \{1, \dots, 8\}$  with  $i < j$ . We have  $A_0 \cdot A_j = 1$  for  $j \in \{1, \dots, 8\}$ ,  $A_0 \cdot A_{ij} = 0$  for  $j \in \{1, \dots, 8\}$  and  $i < j$ , and we have  $A_j \cdot A_{st} = 2$  if and only if  $j \in \{s, t\}$ ; else  $A_j \cdot A_{st} = 0$ . Moreover, for  $\{u, v\} \neq \{s, t\}$ , we have

$$A_{uv} \cdot A_{st} = 2(1 - |\{u, v\} \cap \{s, t\}|).$$

**Proposition 9.4.** *The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve which is the union of a smooth conic  $C_0$  and a quartic  $Q$ .*

The curves  $A_1, \dots, A_8$  are mapped to the 8 nodes of the sextic curve, the 28 divisors  $A_{ij}$  are mapped onto the 28 lines through the 8 nodes, and  $A_0$  is mapped onto the conic  $C_0$ . The moduli space of such surfaces is unirational.

#### 9.5. The lattice $U(2) \oplus \mathbf{A}_1^{\oplus 7}$ and degree 1 del Pezzo surfaces

Let  $C_6$  be a sextic curve in  $\mathbb{P}^2$  with 8 nodes in general position. Let  $Z \rightarrow \mathbb{P}^2$  the blow-up of the nodes; it is a degree 1 del Pezzo surface and contains 240  $(-1)$ -curves:

- the 8 exceptional divisors  $E_i$ ,  $i = 1, \dots, 8$ ,
- the strict transform  $L_{ij}$  of the 28 lines through  $p_i, p_j$  ( $i \neq j$ ),
- the strict transforms  $QO_{rst}$  of the 56 conics that go through points in  $\{p_1, \dots, p_8\} \setminus \{p_r, p_s, p_t\}$ ,
- the strict transforms  $CU_{rj}$  of the 56 cubics that go through 7 points  $p_k$  (with  $k \neq r$ ) with a double point at  $p_j$  ( $j \neq r$ ),
- the strict transforms  $QA_{rst}$  of the 56 quartics through the 8 points  $p_j$  with double points at  $p_r, p_s, p_t$ ,
- the strict transforms  $QI_{ij}$  of the 28 quintics through the 8 points  $p_j$  with double points at  $p_i, p_j$  ( $i \neq j$ ),
- the strict transforms  $S_j$  of the 8 sextics through 8 points with double points at all except a single point  $p_j$  with multiplicity 3.

The Néron–Severi lattice of  $Z$  is generated by the pull-back  $L'$  of a line and  $E_1, \dots, E_8$ ; it is the unimodular rank 9 lattice

$$I_1 \oplus I_{-1}^{\oplus 8}.$$

The anti-canonical divisor of the del Pezzo surface  $Z$  of degree 1 is given by

$$-K_Z = 3L' - (E_1 + \dots + E_8);$$

this is an ample divisor. Moreover, we remark that each of the 120 divisors

$$E_j + S_j, L_{ij} + QI_{ij}, CU_{ij} + CU_{ji}, QO_{ijk} + QA_{ijk}, \quad \{i, j, k\} \subset \{1, \dots, 8\} \text{ with } |\{i, j, k\}| = 3,$$

belongs to the base-point free (see [Dem76, Section 3, Theorem 1]) linear system  $| -2K_Z |$ .

The double cover  $f: Y \rightarrow Z$  branched over the strict transform of  $C_6$  is a smooth K3 surface. The pull-backs of the 240  $(-1)$ -curves are  $(-2)$ -curves (the only negative curves on a K3 surface are  $(-2)$ -curves). We denote by  $A_1, \dots, A_8$  the pull-backs on  $Y$  of the curves  $E_i$  and by  $L$  the pull-back of  $L'$ . Naturally, the lattice  $f^* \text{NS}(Z)$  is  $(I_1 \oplus I_{-1}^{\oplus 8})(2)$ , which is also the lattice generated by  $L, A_1, \dots, A_8$  and is isometric to  $U(2) \oplus \mathbf{A}_1^{\oplus 7}$ .

Since  $f: Y \rightarrow Z$  is finite, its pull-back  $D_2 = f^*(-K_Z)$  is ample, with  $D_2^2 = 2$ . Since  $|-2K_Z|$  is base-point free, the system  $|D_2|$  is non-hyperelliptic; thus it defines a double cover

$$\pi: Y \rightarrow \mathbb{P}^2$$

branched over a sextic curve  $\tilde{C}_6$ , which is smooth since  $D_2$  is ample. Let  $A$  be the pull-back on  $Y$  of a  $(-1)$ -curve  $E$ ; this is a  $(-2)$ -curve. We have  $D_2A = 2(-K_Z E) = 2$ . Moreover, we see from the above description of the 120 divisors in  $|-2K_Z|$  that there exists a  $(-2)$ -curve  $B$  such that  $A + B \equiv 2D_2$ ; in particular, the image of  $A + B$  by  $\pi$  is a conic which is 6-tangent to the sextic  $\tilde{C}_6$ . We thus obtain the following result.

**Proposition 9.5.** *The branch curve  $\tilde{C}_6$  of the morphism  $\pi$  is a smooth sextic curve which possesses 120 6-tangent conics.*

*Remark 9.6.* Let  $B$  be the strict transform in  $Z$  of  $C_6$ ; this is a smooth genus 2 curve such that  $B \equiv E_j + S_j$ . Thus the ramification locus  $R$  of the double cover  $f: Y \rightarrow Z$  is a smooth genus 2 curve with  $2R \equiv f^*(E_j + S_j) \equiv 2D_2$ , and (since  $\text{NS}(X) = \text{Pic}(X)$ ) the divisor  $R$  is in the linear system  $|D_2|$ , and its image by  $\pi$  is a line in  $\mathbb{P}^2$ .


*Remark 9.7.* The morphism  $\pi$  is branched over a smooth sextic curve  $\tilde{C}_6$  (thus of genus 10), whereas  $f$  is branched over a smooth genus 2 curve  $B$ , so that there exist (at least) two distinct non-symplectic involutions  $\iota_1, \iota_2$  on the same K3 surface. In fact, according to Kondo [Kon89], the automorphism group of such a K3 surface is  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ , so that we know the generators of  $\text{Aut}(X)$ .

The curve  $\tilde{C}_6$  thus has 120 conics that are 6-tangent. As far as we know, this is the record for a smooth sextic.

*Remark 9.8.* By a result of Degtyarev [Deg19], a smooth plane sextic curve can have at most 72 tritangent lines. Moreover, there is an example of such a sextic curve by Mukai (see [Muk88b]). In [Elk09], Elkies gives an example of an irreducible sextic curve and 1240 conics that are 6-tangent to it; that curve has a unique node.

### 9.6. The lattice $U \oplus A_7$

The surface  $X$  contains nine  $(-2)$ -curves  $A_1, \dots, A_9$ , with dual graph



These nine curves generate the Néron–Severi lattice; the divisor


$$D_{120} = (9, 19, 15, 12, 10, 9, 10, 12, 15)$$

in that basis has square 120, is ample, with  $D_{120} \cdot A_j = 1$  for  $j \neq 6$  and  $D_{120} \cdot A_6 = 2$ . The divisor  $F = \sum_{j=2}^9 A_j$  is a fiber of an elliptic fibration for which  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 9.9.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $\mathfrak{a}_7$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

### 9.7. The lattice $U \oplus D_4 \oplus A_3$

The surface  $X$  contains 10  $(-2)$ -curves, with dual graph



The curves  $A_1, \dots, A_8$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{56} = (7, 7, 7, 15, 10, 6, 3, 1, 0)$$

is ample, with  $D_{56} \cdot A_j = 1$  for  $j \in \{1, \dots, 10\} \setminus \{9\}$  and  $D_{56} \cdot A_9 = 3$ . The divisors

$$F_1 = A_4 + \sum_{j=1}^5 A_j, \quad F_2 = \sum_{j=7}^{10} A_j$$

are fibers of an elliptic fibration with section  $A_6$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 9.10.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_4$  singularity  $p$  and one  $\mathbf{a}_3$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

The divisor

$$D_2 = A_1 + 2A_2 + 2A_3 + 4A_4 + 3A_5 + 2A_6 + A_7$$


is nef, of square 2 and base-point free. We have  $D_2 \cdot A_1 = 2$ ,  $D_2 \cdot A_8 = D_2 \cdot A_9 = 1$  and  $D_2 \cdot A_j = 0$  for  $j \notin \{1, 8, 9\}$ . Moreover,

$$D_2 \equiv A_2 + A_3 + 2A_4 + 2A_5 + 2A_6 + 2A_7 + A_8 + A_9 + A_{10}.$$

Thus the K3 surface is a double cover of the plane branched over a sextic curve with a  $\mathbf{d}_6$  singularity and a node  $\mathbf{a}_1$ .

### 9.8. The lattice $U \oplus D_5 \oplus A_2$

The surface  $X$  contains 10  $(-2)$ -curves, with dual graph



The Néron–Severi lattice is generated by  $A_1, \dots, A_9$ ; moreover,

$$A_{10} = (1, 1, 2, 2, 1, 1, 0, -1, -1)$$

in that base. The divisor

$$D_{88} = (8, 8, 17, 19, 9, 13, 8, 4, 1)$$

is ample, of square 88, with  $D_{88} \cdot A_j = 1$  for  $j \leq 8$ ,  $D_{88} \cdot A_9 = 2$  and  $D_{88} \cdot A_{10} = 5$ . The divisors

$$F_1 = A_1 + A_2 + 2A_3 + 2A_4 + A_5 + A_6, \quad F_2 = A_8 + A_9 + A_{10}$$

are fibers of an elliptic fibration with section  $A_7$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 9.11.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the negative section  $s$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_5$  singularity  $p$  and one  $\mathbf{a}_2$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

The divisor

$$D_2 = A_1 + A_2 + 3A_3 + 4A_4 + 2A_5 + 3A_6 + 2A_7 + A_8$$


is nef, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = D_2 \cdot A_9 = D_2 \cdot A_{10} = 1$  and  $D_2 \cdot A_j = 0$  for  $j \in \{1, 2, 9, 10\}$ . Since

$$D_2 \equiv A_3 + 2A_4 + A_5 + 2A_6 + 2A_7 + 2A_8 + A_9 + A_{10},$$

we get that the K3 surface is the minimal resolution of a double cover of  $\mathbb{P}^2$  branched over a sextic curve with a  $\mathbf{d}_6$  singularity.

### 9.9. The lattice $U \oplus D_7$

The surface  $X$  contains nine  $(-2)$ -curves, with dual graph



The nine curves generate the Néron-Severi lattice, and in that basis, the divisor

$$D_{260} = (13, 27, 42, 20, 38, 35, 33, 16, 16)$$

is ample, of square 260, with  $D_{260} \cdot A_2 = 2$  and  $D_{260} \cdot A_j = 1$  for  $j \neq 4$ . The divisor

$$F = A_2 + A_4 + 2(A_3 + A_5 + A_6 + A_7) + A_8 + A_9$$

is a fiber of an elliptic fibration with section  $A_1$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 9.12.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_7$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*


One can also construct that surface as a double plane: the divisor

$$D_2 = 2A_1 + 4A_2 + 6A_3 + 3A_4 + 5A_5 + 4A_6 + 3A_7 + A_8 + A_9$$

is nef, of square 2, with  $D_2 \cdot A_j = 0$  for  $j \leq 7$  and  $D_2 \cdot A_8 = D_2 \cdot A_9 = 1$ . The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve with an  $\mathbf{e}_7$  singularity.

### 9.10. The lattice $U \oplus A_1 \oplus E_6$

The surface  $X$  contains 10  $(-2)$ -curves, with dual graph



The nine curves  $A_1, \dots, A_9$  generate the Néron-Severi lattice, and in that basis, the divisor

$$D_{184} = (12, 17, 12, 8, 5, 3, 1, 0, 1)$$

is ample, of square 184, with  $D_{184} \cdot A_1 = 10$ ,  $D_{184} \cdot A_2 = 2$  and  $D_{184} \cdot A_j = 1$  for  $j \geq 3$ . The divisors

$$F_1 = A_1 + A_2, \quad F_2 = A_4 + A_8 + A_{10} + 2(A_5 + A_7 + A_9) + 3A_6$$

are fibers of an elliptic fibration of  $X$  with a section  $A_3$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 9.13.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and one  $\mathbf{e}_6$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

One can also construct that surface as a double plane: the divisor


$$D_2 = A_1 + 2(A_2 + A_3 + A_4 + A_5 + A_6) + A_7 + A_9$$

is nef, base-point free, of square 2. We have  $D_2 \cdot A_1 = 2$ ,  $D_2 \cdot A_8 = D_2 \cdot A_{10} = 1$  and  $D_2 \cdot A_j = 0$  for the other curves. The K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve with a  $\mathbf{d}_7$  singularity.

## 10. Rank 10 lattices

### 10.1. The lattice $U \oplus \mathbf{E}_8$

The K3 surface  $X$  contains 10  $(-2)$ -curves with dual graph



These curves generate the Néron–Severi lattice. In that base, the divisor

$$D_{1240} = (30, 61, 93, 126, 160, 195, 231, 115, 153, 76)$$

is ample, of square 1240, with  $D_{1240} \cdot A_j = 1$  for  $j \in \{1, \dots, 10\}$ . The divisor

$$F = A_2 + 2A_3 + 3A_4 + 4A_5 + 5A_6 + 6A_7 + 3A_8 + 4A_9 + 2A_{10}$$


is a fiber of an elliptic fibration and  $A_1$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 10.1.** *The linear system  $|4F + 2A_1|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has an  $\mathbf{e}_8$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

*Remark 10.2.* Using that  $U \oplus \mathbf{E}_8$  is unimodular, one can prove that if  $D$  is an ample divisor, then  $D^2 \geq 1240$ .

### 10.2. The lattice $U \oplus \mathbf{D}_8$

The K3 surface  $X$  contains 10  $(-2)$ -curves with dual graph



These curves generate the Néron–Severi lattice. In that base, the divisor

$$D_{280} = (15, 15, 31, 33, 36, 40, 45, 22, 29, 14)$$

is ample, with  $D_{280} \cdot A_j = 1$  for  $j \in \{1, \dots, 10\}$ . The divisor

$$F = A_1 + A_2 + 2(A_3 + \dots + A_7) + A_8 + A_9$$


is a fiber of an elliptic fibration with section  $A_{10}$ . By Theorem 2.3, case i) a), we have the following.



**Proposition 10.3.** *The linear system  $|4F + 2A_{10}|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a  $\mathbf{d}_8$  singularity  $q$ . The pull-back of the fiber through  $q$  is the fiber  $F$ .*

### 10.3. The lattice $U \oplus E_7 \oplus A_1$

The K3 surface  $X$  contains 11  $(-2)$ -curves with dual graph



The curves  $A_1, \dots, A_{10}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{370} = (15, 24, 19, 15, 12, 10, 9, 4, 5, 2)$$

is ample, of square 370, with  $D_{370} \cdot A_j = 1$  for  $j \in \{2, \dots, 10\}$ ,  $D_{370} \cdot A_1 = 18$  and  $D_{370} \cdot A_{11} = 2$ . The divisors


$$F_1 = A_1 + A_2, \quad F_2 = A_4 + 2A_5 + 3A_6 + 4A_7 + 2A_8 + 3A_9 + 2A_{10} + A_{11}$$

are fibers of an elliptic fibration with section  $A_3$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 10.4.** *The linear system  $|4F_1 + 2A_3|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has a node  $p$  and one  $\mathbf{e}_7$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 10.4. The lattice $U \oplus D_4^{\oplus 2}$

The K3 surface  $X$  contains 11  $(-2)$ -curves with dual graph



The curves  $A_1, \dots, A_{10}$  generate the Néron–Severi lattice. In that base, we have

$$A_{11} = (1, 2, 1, 1, 1, 0, -1, -2, -1, -1).$$

The divisor

$$D_{56} \equiv (7, 15, 7, 7, 10, 6, 3, 1, 0, 0)$$

is ample of square 56, and the  $(-2)$ -curves on  $X$  have degree 1 for  $D_{56}$ . The divisors

$$F_1 = A_2 + \sum_{j=1}^5 A_j, \quad F_2 = A_8 + \sum_{j=7}^{11} A_j,$$

are fibers of an elliptic fibration with section  $A_6$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 10.5.** *The linear system  $|4F_1 + 2A_6|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has two  $\mathbf{d}_4$  singularities. The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

One can give another construction as follows. The divisor

$$D_2 = A_1 + 4A_2 + 2A_3 + 2A_4 + 3A_5 + 2A_6 + A_7$$


is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = 2$ ,  $D_2 \cdot A_8 = 1$  and  $D_2 \cdot A_j = 0$  for  $j \notin \{2, 8\}$ . We have

$$D_2 \equiv 2A_2 + A_3 + A_4 + 2A_5 + 2A_6 + 2A_7 + 2A_8 + A_9 + A_{10} + A_{11}.$$

The branch curve of the double cover  $\varphi: X \rightarrow \mathbb{P}^2$  associated to  $D_2$  is a sextic with three nodes and a  $\mathbf{d}_6$  singularity  $q$ . The curve  $A_8$  is in the ramification locus; its image by  $\varphi$  is a line  $L$ . Let  $Q$  be the residual quintic curve of the sextic branch locus. The line  $L$  cuts  $Q$  transversally in an  $\mathbf{a}_3$  singularity and in three other points  $p_1, p_2, p_3$ . The curves  $A_9, A_{10}, A_{11}$  are mapped to these three points. The image by  $\varphi$  of  $A_1$  is the line that is the branch of the  $\mathbf{a}_3$  singularity. The curves  $A_2, \dots, A_7$  are mapped to  $q$ .

**10.5. The lattice  $U \oplus D_6 \oplus A_1^{\oplus 2}$**

The K3 surface  $X$  contains 14  $(-2)$ -curves with dual graph



The curves  $A_1, \dots, A_{10}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{98} = (7, 16, 13, 11, 5, 5, 13, 11, 5, 5)$$

is ample, of square 98, with  $D_{98} \cdot A_j = 1$  for  $j \in \{2, \dots, 10\}$ ,  $D_{98} \cdot A_1 = 2$  and  $D_{98} \cdot A_j = 10$  for  $j \in \{11, \dots, 14\}$ . The divisor

$$D_2 = (1, 2, 2, 2, 1, 1, 2, 2, 1, 1)$$


is nef, base-point free, of square 2, with  $D_2 \cdot A_2 = 1$ ,  $D_2 \cdot A_j = 2$  for  $j \in \{11, \dots, 14\}$  and  $D_2 \cdot A_j = 0$  for the remaining curves. By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 10.6.** *The branch curve has a node and two  $\mathbf{d}_4$  singularities. It is the union of a quintic curve with two nodes  $q, q'$  and the line through the points  $q, q'$ . The line cuts the quintic in a third point  $p$ , forming a node.*

The curve  $A_1$  is sent to  $p$ , the curves  $A_3, A_4, A_5, A_6$  are sent to  $q$ , and the curves  $A_7, A_8, A_9, A_{10}$  are sent to  $q'$ . The curves  $A_{11}$  and  $A_{12}$  are sent to the two branches of the singularity  $q$  that are distinct from  $L$ ; the curves  $A_{13}$  and  $A_{14}$  are sent to the two branches of the singularity  $q'$  that are distinct from  $L$ .

**10.6. The lattice  $U(2) \oplus D_4^{\oplus 2}$**

**10.6.1. First involution.**— There exist 18  $(-2)$ -curves  $A_1, \dots, A_{18}$  on  $X$ , with dual graph



The curves  $A_1, \dots, A_8, A_{10}, A_{18}$  generate the Néron–Severi lattice. The divisor

$$D_8 = A_1 + 3A_2 + 3A_{10} + A_{18}$$

is ample, base-point free, of square 8, with  $D_8 \cdot A_j = 1$  for all  $j \in \{1, \dots, 18\}$ . The divisors  $A_k + A_{8+k} \equiv F$ ,  $k \in \{2, \dots, 9\}$ , are singular fibers of an elliptic fibration, for which  $A_1$  and  $A_{18}$  are sections. Since  $FD_8 = 2$ , we see that  $D_8$  is hyperelliptic; thus the image by the associated map  $\varphi$  of  $X$  is a rational normal scroll of degree 4 in  $\mathbb{P}^5$ , *i.e.*, the Hirzebruch surface  $F_2$ . The pull-back by  $\varphi$  of the unique negative curve  $s$  (the section) is the union of two disjoint  $(-2)$ -curves. The branch curve  $B$  must satisfy  $Bs = 0$ ; thus  $B = v(s + 2f)$ , where  $f$  is a fiber of the unique fibration of  $F_2$ . We have  $F = \varphi^*f$ ; thus the branch curve cuts a general fiber  $f$  in four points and  $v = 4$ .

**Proposition 10.7.** *The K3 surface  $X$  is the double cover of  $F_2$  branched on a smooth curve  $B$  of genus 9 in  $|4s + 8f|$ . The curve  $B$  is such that there are 8 fibers  $f_1, \dots, f_8$  that meet  $B$  with even multiplicities at each intersection point. The 18 curves on  $X$  are in the pull-back of  $s$  (which is  $A_1 + A_{18}$ ) and the pull-backs of the fibers  $f_i$ ,  $i \in \{1, \dots, 8\}$ .*

**10.6.2. Second involution.**— The divisor

$$D_2 = 2A_1 + 2A_2 + A_3 + A_4 + A_{10}$$

is nef, of square 2, with  $D_2 \cdot A_j = 2$  for  $j \in \{5, \dots, 12\}$ ,  $D_2 \cdot A_{18} = 1$  and else  $D_2 \cdot A_j = 0$ . We have

$$D_2 \equiv 2A_1 + A_2 + A_3 + A_4 + A_k + A_{8+k}, \quad k \in \{2, \dots, 9\},$$

and

$$D_2 \equiv A_{13} + A_{14} + A_{15} + A_{16} + A_{17} + 2A_{18}.$$


**Proposition 10.8.** *The K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with a  $\mathbf{d}_4$  singularity  $q$  (onto which  $A_1, A_2, A_3, A_4$  are mapped) and five nodes (onto which the five curves  $A_{13}, \dots, A_{17}$  are mapped).*

The curve  $C_6$  is the union of a line  $L$  and a quintic with a  $\mathbf{d}_4$  singularity. The double cover is ramified over  $A_{18}$ , and the image of  $A_{18}$  is  $L$ . The images of  $A_{10}, A_{11}, A_{12}$  are conics tangent to the three branches of the  $\mathbf{d}_4$  singularity; the images of  $A_{13}, \dots, A_{17}$  are the five lines through the five nodes (intersection of  $L$  and the quintic) and  $q$ .

*Remark 10.9.* According to [Kon89], the automorphism group of  $X$  is  $(\mathbb{Z}/2\mathbb{Z})^2$ . The fixed loci of the two involutions we described are distinct; thus we obtained generators of the automorphism group.

### 10.7. The lattice $U \oplus D_4 \oplus A_1^{\oplus 4}$

The K3 surface  $X$  contains 27  $(-2)$ -curves  $A_1, \dots, A_{27}$ ; the curves  $A_1, \dots, A_{11}$  have dual graph



and  $A_1, \dots, A_{10}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{26} = (3, 3, 3, 3, 7, 3, 1, 0, 0, 0)$$

is ample, of square 26, with  $D_{26} \cdot A_j = 1$  for  $j \in \{1, \dots, 11\}$ ,  $j \neq 5$ ,  $D_{26} \cdot A_5 = 2$  and  $D_{26} \cdot A_k = 6$  for the remaining  $(-2)$ -curves. The divisors

$$A_1 + A_2 + A_3 + A_4 + 2A_5, \quad 2A_7 + A_8 + A_9 + A_{10} + A_{11}$$

are fibers of the same fibration. That gives the class of  $A_{11}$  in the basis  $A_1, \dots, A_{10}$ . The divisor

$$F_5 = A_6 + 2A_7 + A_8 + A_9 + A_{10}$$

is a fiber of an elliptic fibration. For  $j \in \{1, \dots, 4\}$ , one has  $A_{11+j} = F_5 - A_j$  (as a class). The divisor

$$F_1 = A_2 + A_3 + A_4 + 2A_5 + A_6$$

is a fiber of an elliptic fibration. For  $j \in \{1, \dots, 3\}$ , one has  $A_{15+j} = F_1 - A_{7+j}$  (as a class). The divisor

$$F_2 = A_1 + A_3 + A_4 + 2A_5 + A_6$$

is a fiber of an elliptic fibration. For  $j \in \{1, \dots, 3\}$ , one has  $A_{18+j} = F_2 - A_{7+j}$  (as a class). The divisor

$$F_3 = A_1 + A_2 + A_4 + 2A_5 + A_6$$

is a fiber of an elliptic fibration. For  $j \in \{1, \dots, 3\}$ , one has  $A_{21+j} = F_3 - A_{7+j}$  (as a class). The divisor

$$F_4 = A_1 + A_2 + A_3 + 2A_5 + A_6$$

is a fiber of an elliptic fibration. For  $j \in \{1, \dots, 3\}$ , one has  $A_{24+j} = F_4 - A_{7+j}$  (as a class).

The divisor

$$D_2 = A_1 + A_2 + A_3 + A_4 + 2A_5 + A_6$$

is nef, of square 2, with  $D_2 \cdot A_5 = D_2 \cdot A_7 = 1$ ,  $D_2 \cdot A_k = 0$  for  $k \in \{1, \dots, 4, 6, 8, \dots, 11\}$  and  $D_2 \cdot A_j = 2$  for the remaining curves. We also have

$$D_2 \equiv A_6 + 2A_7 + A_8 + A_9 + A_{10} + A_{11}.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 10.10.** *The surface  $X$  is the double cover of the plane branched over a sextic curve which is the union of two lines  $L, L'$  and a quartic  $Q$ .*

The images of  $A_5, A_7$  are the two lines. We observe that

$$\begin{aligned} D_2 &\equiv A_{16} + A_1 + A_8 \equiv A_{17} + A_1 + A_9 \equiv A_{18} + A_1 + A_{10}, \\ D_2 &\equiv A_{19} + A_2 + A_8 \equiv A_{20} + A_2 + A_9 \equiv A_{21} + A_2 + A_{10}, \\ D_2 &\equiv A_{22} + A_3 + A_8 \equiv A_{23} + A_3 + A_9 \equiv A_{24} + A_3 + A_{10}, \\ D_2 &\equiv A_{25} + A_4 + A_8 \equiv A_{26} + A_4 + A_9 \equiv A_{27} + A_4 + A_{10}. \end{aligned}$$

Thus  $A_1, A_2, A_3, A_4$  are contracted to the nodes in the intersection of  $Q$  and  $L$ , and so are  $A_8, A_9, A_{10}$ . Since

$$D_2 \equiv A_1 + A_{11} + A_{12} \equiv A_2 + A_{11} + A_{13} \equiv A_3 + A_{11} + A_{14} \equiv A_4 + A_{11} + A_{15},$$

the curve  $A_{11}$  is contracted to the fourth intersection point of  $L'$  and  $Q$ . Moreover, we see from these equivalences that the 16  $(-2)$ -curves  $A_{12}, \dots, A_{27}$  are sent to the 16 lines between points in  $L \cap Q$  and points in  $L' \cap Q$ . The curve  $A_6$  is contracted onto the intersection point of  $L$  and  $L'$ .

Proposition 10.10 implies that the moduli space of K3 surfaces with Néron–Severi group isometric to  $U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 4}$  is unirational.

## 10.8. The lattice $U \oplus \mathbf{A}_1^{\oplus 8}$

**10.8.1. First involution.**— Let us denote by  $f_1, f_2, e_1, \dots, e_8$  the canonical basis of  $U \oplus \mathbf{A}_1^{\oplus 8}$ . In that basis, let

$$D_6 = (7, 5, -2, -2, -2, -2, -2, -2, -2, -2).$$

It has square 6, and no  $(-2)$ -classes are perpendicular to it. We thus have a marking such that  $U \oplus \mathbf{A}_1^{\oplus 6} \simeq \text{NS}(X)$  which maps  $D_6$  to an ample class. The K3 surface  $X$  contains 145  $(-2)$ -curves; with respect to  $D_6$ , the curves  $A_1, \dots, A_{16}$  have degree 1, the curve  $A_0$  has degree 2, and the remaining curves have degree 4. Let us describe these curves.

For  $j \in \{1, \dots, 8\}$ , one has  $A_j = f_1 - e_j$ ; the divisors

$$A_1 + A_9, \dots, A_8 + A_{16}$$

are fibers of an elliptic fibration  $\varphi: X \rightarrow \mathbb{P}^1$ , where the class of a fiber  $F$  is

$$F = (4, 2, -1, -1, -1, -1, -1, -1, -1, -1)$$

in the canonical basis. The curve  $A_0$  is  $A_0 = -f_1 + f_2$ , and  $A_0 \cdot A_k = 1$  for  $1 \leq k \leq 16$ .

For the choice of any three elements  $\{i, j, k\}$  in  $\{1, \dots, 8\}$  (56 possibilities), the classes

$$\begin{aligned} A_{i,j,k} &= 4f_1 + 4f_2 - e_i - e_j - e_k - \sum_{l=1}^8 e_l, \\ B_{i,j,k} &= 2f_1 + 2f_2 + e_i + e_j + e_k - \sum_{l=1}^8 e_l \end{aligned}$$

are classes of  $(-2)$ -curves. The 16 classes

$$C_j = 6f_1 + 6f_2 - e_j - 2\sum_{l=1}^8 e_l \quad \text{and} \quad E_j = e_j, \quad j \in \{1, \dots, 8\}$$

are classes of  $(-2)$ -curves. Thus in total, we get 145  $(-2)$ -curves.

The divisor

$$D_2 = (3, 3, -1, -1, -1, -1, -1, -1, -1, -1) = F + A_0$$

has square 2, and  $D_2 \cdot A_j = 1$  for  $j \in \{1, \dots, 16\}$ ,  $D_2 \cdot A_0 = 0$  and  $D_2 A = 2$  for the remaining  $(-2)$ -curves  $A$ . Let  $C_6$  be the sextic branch curve of the associated double cover  $X \rightarrow \mathbb{P}^2$ . The curve  $C_6$  has a node  $q$  onto which the curve  $A_0$  is contracted. For  $j \in \{1, \dots, 8\}$ , we have

$$A_j + A_{8+j} + A_0 \equiv D_2;$$

thus the divisor  $A_j + A_{8+j} + A_0$  is the pull-back of a line through the nodal point  $q$ , and that line is tangent to  $C_6$  at every other intersection points. For a set of three elements  $\{i, j, k\}$  in  $\{1, \dots, 8\}$ , we have

$$A_{i,j,k} + B_{i,j,k} \equiv 2D_2;$$

thus the 56 divisors  $A_{i,j,k} + B_{i,j,k}$  are pull-backs of 56 conics that are 6-tangent to  $C_6$  (and not containing the nodal point of  $C_6$ ). We have

$$C_j + E_j \equiv 2D_2;$$

thus the eight divisors  $C_j + E_j$ ,  $j \in \{1, \dots, 8\}$ , are also pull-backs of eight conics that are 6-tangent to  $C_6$ . Summing up, we have the following.

**Proposition 10.11.** *The K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a nodal sextic curve. Through the node of the sextic, there are eight lines that are tangent to the sextic at other intersection points. Moreover, there are 64 conics that are 6-tangent to the sextic.*

The Néron–Severi lattice is generated by the classes of  $A_0, A_1, \dots, A_8$  and  $E_8$ .

*Remark 10.12.* Let  $S$  be the set of the 128  $(-2)$ -curves that are above the 64 conics. One can prove that there exists a partition of  $S$  into eight sets  $S_1, \dots, S_8$  of 16 curves such that for any such set  $S_i = \{B_1, \dots, B_{16}\}$ , one has (up to permutation of the indices)  $B_{2k-1} \cdot B_{2k} = 6$  for  $k \in \{1, \dots, 8\}$ , and for  $s, t$  ( $s \neq t$ ) such that  $\{s, t\} \neq \{2k-1, 2k\}$ , one has  $B_s \cdot B_t = 4$  if  $s+t = 1 \pmod{2}$ ; else  $B_s \cdot B_t = 0$ .

For  $j \neq t$  in  $\{1, \dots, 8\}$ , if  $B$  is a curve in  $S_j$ , there are 10  $(-2)$ -curves  $B'$  in  $S_t$  such that  $BB' = 2$ , 3  $(-2)$ -curves such that  $BB' = 0$  and 3  $(-2)$ -curves such that  $BB' = 4$ .

*Remark 10.13.* The ample divisor  $D_6$  satisfies  $D_6 \equiv 2F + A_0$  and  $FD_6 = 2$ ; thus it is hyperelliptic. According to Theorem [Sai74], the morphism associated to  $|D_6|$  is a degree 2 map onto a degree 3 rational normal scroll in  $\mathbb{P}^4$ . That surface is the Hirzebruch surface  $\mathbb{F}_1$ , which is the blow-up of  $\mathbb{P}^2$  in one point, embedded by  $|2L - E_0|$ , where  $L$  is the pull-back of a line and  $E_0$  is the exceptional divisor. The composite of  $X \rightarrow \mathbb{F}_1$  with the natural map  $\mathbb{F}_1 \rightarrow \mathbb{P}^2$  is given by the linear system  $|D_2|$ .

**10.8.2. Second involution: A more geometric interpretation of the  $(-2)$ -curves.**— Let  $f_1, f_2, e_1, \dots, e_8$  be the canonical basis of  $U \oplus \mathbf{A}_1^{\oplus 8}$ . The divisor  $D'_2 = f_1 + f_2$  is nef, of square 2. We recall that  $A_0 = -f_1 + f_2$  is a  $(-2)$ -curve; moreover, the divisor  $F' = f_1$  is a fiber of an elliptic fibration; thus

$$D'_2 = 2F' + A_0.$$

One has  $FA_0 = 1$ ,  $D'_2F' = 1$ , and therefore the linear system  $|D'_2|$  has base points. We have  $D'_2 \cdot A_0 = 0$  and

$$\begin{aligned} D'_2 \cdot A_j &= 1, \quad D'_2 \cdot A_{j+8} = 5, & \forall j \in \{1, \dots, 8\}, \\ D'_2 \cdot A_{i,j,k} &= 8, & \forall \{i, j, k\} \subset \{1, \dots, 8\} \text{ of order } 3, \\ D'_2 \cdot B_{i,j,k} &= 4, & \forall \{i, j, k\} \subset \{1, \dots, 8\} \text{ of order } 3, \\ D'_2 \cdot C_j &= 12, \quad D'_2 \cdot E_j = 0, & \forall k \in \{1, \dots, 8\}. \end{aligned}$$

Let us define  $D_8 = 2D'_2$ . The linear system  $|D_8|$  is base-point free, and it is hyperelliptic since  $D_8F' = 2$ . By Theorem 2.3, case iii) v), the associated map  $\varphi_{D_8}: X \rightarrow \mathbb{P}^5$  has image a cone over a rational normal curve in  $\mathbb{P}^4$ ; it factorizes through a surjective map  $\varphi': X \rightarrow \mathbf{F}_4$ , where  $\mathbf{F}_4$  is the Hirzebruch surface with a section  $s$  such that  $s^4 = -4$ . The section  $s$  is mapped to the vertex of the cone by the map  $\mathbf{F}_4 \rightarrow \mathbb{P}^5$ . Let  $f$  denote a fiber of the unique fibration  $\mathbf{F}_4 \rightarrow \mathbb{P}^1$ . The branch locus of  $\varphi'$  is the union of  $s$  and a curve  $C$  such that  $Cs = 0$  and  $C \in |3(s+4f)|$  (so that  $s+C \in |-2K_{\mathbf{F}_4}|$ ). The curves  $E_1, \dots, E_8$  being contracted by  $\varphi'$ , the curve  $C$  has eight nodes  $p_1, \dots, p_8$ , which are the images of  $E_1, \dots, E_8$ . We have

$$|D_8| = \varphi'^*|4f + s|.$$

We moreover have the relations

$$A_j + E_j \equiv F' = \varphi'^*f, \quad \forall j \in \{1, \dots, 8\};$$

therefore, the curves  $A_1, \dots, A_8$  are mapped by  $\varphi'$  to the eight fibers going through the nodes  $p_1, \dots, p_8$ . Since  $D_8 \equiv 4(A_1 + E_1) + 2A_0$ , the curve  $A_0$  is in the ramification locus, with image  $s$ . Since

$$A_{8+k} + \sum_{j=1, j \neq k}^{j=8} E_j \equiv D_8 + F', \quad \forall k \in \{1, \dots, 8\},$$

the image of  $A_{8+k}$  belongs in  $|5f + s|$ ; it is a curve which goes through the seven points  $\{p_1, \dots, p_8\} \setminus \{p_k\}$ . We moreover have the relations

$$(10.1) \quad \begin{aligned} D_8 &\equiv B_{ijk} - (E_i + E_j + E_k) + \sum_{t=1}^8 E_t, \\ 2D_8 &\equiv A_{ijk} + E_i + E_j + E_k + \sum_{t=1}^8 E_s, \\ 3D_8 &\equiv C_k + E_k + 2 \sum_{t=1}^8 E_s, \end{aligned}$$

and therefore

- the image of  $B_{ijk}$  is a curve in the linear system  $|4f + s|$  which goes through the five points distinct from  $p_i, p_j, p_k$ ;
- the image of  $A_{ijk}$  is a curve in the linear system  $|2(4f + s)|$  which goes through the eight points  $p_1, \dots, p_8$  with double points at  $p_i, p_j, p_k$ ;
- the image of  $C_k$  is a curve in the linear system  $|3(4f + s)|$  which goes through eight points with double points at all except at the single point  $p_k$  with multiplicity 3.

Let  $J, J'$  be subsets of order 3 of  $\{1, \dots, 8\}$ . Using the relations in (10.1) and  $2D_2 \equiv A_J + B_J \equiv C_k + E_k$ , one finds that


$$\begin{aligned} A_J \cdot A_{J'} &= B_J \cdot B_{J'} = 4 - 2\#(J \cap J'), \\ A_J \cdot B_{J'} &= 2\#(J \cap J'), \\ C_k \cdot C_{k'} &= E_k E_{k'} = -2\delta_{kk'}, \\ C_k \cdot E_{k'} &= 4 + 2\delta_{kk'}, \\ C_k \cdot B_J &= 4 \text{ if } k \in J, \quad C_k \cdot B_J = 2 \text{ if } k \notin J, \\ E_k \cdot B_J &= 0 \text{ if } k \in J, \quad E_k \cdot B_J = 2 \text{ if } k \notin J. \end{aligned}$$

The situation is very much similar to what happens for K3 surfaces  $Y$  with Néron-Severi lattice  $U(2) \oplus A_1^{\oplus 7}$ , which are double covers of del Pezzo surfaces of degree 1, where the 240  $(-2)$ -curves of  $Y$  come from lines, conics, cubics, quartics, quintics and sextics going through 8 points in the plane with various multiplicities. In particular, the 112  $(-2)$ -curves  $A_J, B_J$  with  $J \subset \{1, \dots, 8\}$  of order 3 has the same configurations as the 112  $(-2)$ -curves on a K3 surface  $Y$  which are pull-backs of conics and quartics.

*Remark 10.14.* According to Kondo [Kon89], the automorphism group of the surface  $X$  is  $(\mathbb{Z}/2\mathbb{Z})^2$ . The branch loci of the two involutions associated to  $D_2$  and  $D_8$  have genus 9 and 2, respectively. Thus these two involutions generate the automorphism group of  $X$ .

### 10.9. The lattice $U \oplus A_2 \oplus E_6$

The K3 surface  $X$  contains 11  $(-2)$ -curves  $A_1, \dots, A_{11}$ ; their dual graph is



The curves  $A_1, \dots, A_{10}$  generate the Néron-Severi lattice. In that base, the divisor

$$D_{160} = (10, 21, 33, 21, 10, 25, 18, 12, 7, 3)$$

is ample, of square 160, with  $D_{160} \cdot A_j = 1$  for  $j \leq 10$  and  $D_{160} \cdot A_{11} = 10$ . The divisors

$$F_1 = A_9 + A_{10} + A_{11}, \quad F_2 = A_1 + A_7 + A_5 + 2(A_2 + A_4 + A_6) + 3A_3$$


are fibers of an elliptic fibration and  $A_8$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 10.15.** *The linear system  $|4F_1 + 2A_8|$  defines a morphism  $\varphi: X \rightarrow \mathbb{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{a}_2$  singularity  $p$  and one  $\mathbf{e}_6$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

## 11. Rank 11 lattices

### 11.1. The lattice $U \oplus E_8 \oplus A_1$

The K3 surface  $X$  contains 12  $(-2)$ -curves  $A_1, \dots, A_{12}$ ; their dual graph is



The curves  $A_1, \dots, A_{11}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{848} = (48, 97, 147, 73, 125, 104, 84, 65, 47, 30, 14)$$

is ample, of square 848, with  $D_{848} \cdot A_j = 1$  for  $j \in \{1, \dots, 10\}$ ,  $D_{848} \cdot A_{11} = 2$  and  $D_{848} \cdot A_{12} = 28$ . The divisors


$$F_1 = 2A_1 + 4A_2 + 6A_3 + 3A_4 + 5A_5 + 4A_6 + 3A_7 + 2A_8 + A_9, \quad F_2 = A_{11} + A_{12}$$

are fibers of an elliptic fibration with section  $A_{10}$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 11.1.** *The linear system  $|4F_1 + 2A_{10}|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{e}_8$  singularity  $p$  and one node  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

### 11.2. The lattice $U \oplus D_8 \oplus A_1$

The K3 surface  $X$  contains 14  $(-2)$ -curves with dual graph



The curves  $A_1, \dots, A_9, A_{11}, A_{12}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{208} = (-23, -45, -35, -24, -12, 1, 15, 15, 9, -16, -31)$$

is ample, of square 208; the degrees  $D_{208} \cdot A_j$  of the  $(-2)$ -curves  $A_j$ ,  $j = 1, \dots, 14$ , are

$$1, 1, 1, 1, 1, 1, 1, 18, 12, 2, 1, 1, 1, 18.$$

The divisor  $F_1 = A_9 + A_{10}$  is the fiber of an elliptic fibration, and there is a second fiber  $F_2$  supported on  $A_1, \dots, A_7, A_{12}, A_{13}$  of type  $\tilde{\mathbf{D}}_8$ . The curve  $A_{11}$  is a section. By Theorem 2.3, case i) a), we have the following.

**Proposition 11.2.** *The linear system  $|4F_1 + 2A_{11}|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{d}_8$  singularity  $p$  and one node  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_2, F_1$ .*

One can also find a construction using the double cover associated to the divisor

$$D_2 = A_7 + A_8 + A_9$$


which is nef, base-point free, of square 2, with intersections  $D_2 \cdot A_j$ ,  $j = 1, \dots, 14$ , equal to, respectively,

$$0, 0, 0, 0, 0, 1, 0, 2, 0, 2, 0, 0, 0, 2.$$

### 11.3. The lattice $U \oplus D_4 \oplus D_4 \oplus A_1$

The K3 surface contains 22  $(-2)$ -curves  $A_1, \dots, A_{22}$ . The configuration of the curves  $A_1, \dots, A_{15}$  is





The curves  $A_1, \dots, A_{11}$  generate the rank 11 Néron–Severi lattice. In that base, the divisor

$$D_{48} = (1, 3, 3, 3, 1, 0, 0, 0, 1, 2, 0)$$

is ample, of square 48. We have

$$\begin{aligned} A_{13} &\equiv (-2, -3, -2, -2, -2, -1, -1, 2, 4, 3, 1), \\ A_{14} &\equiv (0, 0, 0, -1, -2, -1, -1, 0, 1, 1, 0), \\ A_{15} &\equiv (0, 1, 1, 2, 2, 1, 1, -2, -3, -2, 0). \end{aligned}$$

The divisor  $D_2 = A_2 + A_3 + A_4$  is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_5 = 1$ ,  $D_2 \cdot A_j = 0$  for  $j \in \{2, 4, 6, 7, 8, 9, 12, 13, 14, 15\}$  and else  $D_2 \cdot A_j = 2$ . We have

$$\begin{aligned} D_2 &\equiv 2A_1 + (A_2 + 2A_8 + A_9 + A_{12} + A_{13} + A_{14} + 2A_{15}), \\ D_2 &\equiv 2A_5 + (A_4 + A_6 + A_7 + 2A_8 + A_9 + 2A_{14} + A_{15}), \\ D_2 &\equiv A_{10} + (2A_8 + 2A_9 + A_{14} + A_{15}) \equiv A_{11} + (A_4 + A_{12}), \\ D_2 &\equiv A_{16} + A_4 + A_{13} \equiv A_{17} + A_2 + A_6 \equiv A_{18} + A_6 + A_{12}, \\ D_2 &\equiv A_{19} + A_{13} + A_6 \equiv A_{20} + A_2 + A_7 \equiv A_{21} + A_7 + A_{12}, \\ D_2 &\equiv A_{22} + A_7 + A_{13}. \end{aligned}$$


Therefore, the K3 surface is a double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$ , and the following holds.

**Proposition 11.3.** *The curves  $A_1, A_5$  are in the ramification locus, and their images are lines  $L, L'$  that are components of  $C_6$ . Let  $Q$  be the residual quartic curve. The lines  $L, L'$  meet in a point  $q$  which is on  $Q$ , so that the sextic has a  $\mathbf{d}_4$  singularity at  $q$ .*

The curves  $A_8, A_9, A_{14}, A_{15}$  are contracted to  $q$ . The line  $L$  meets the quartic in three other points  $p_2, p_{12}, p_{13}$ , which are the images of  $A_2, A_{12}, A_{13}$ . The line  $L'$  meet the quartic in three other points  $p_4, p_6, p_7$ , which are the images of  $A_4, A_6, A_7$ . The images of the nine curves  $A_3, A_{11}, A_{16}, \dots, A_{22}$  are lines through the points  $(p_2, p_4), (p_4, p_{12}), (p_4, p_{13}), (p_2, p_6), (p_6, p_{12}), (p_6, p_{13}), (p_2, p_7), (p_7, p_{12}), (p_7, p_{13})$ , respectively. Finally, the image of  $A_{10}$  is a line tangent to the third branch of the  $\mathbf{d}_4$  singularity.

#### 11.4. The lattice $U \oplus D_4 \oplus A_1^{\oplus 5}$

**11.4.1. First involution.**— The K3 surface contains 90  $(-2)$ -curves. The first 20 curves  $A_1, \dots, A_{20}$  have the following configuration:



The curves  $A_1, \dots, A_{11}$  generate the Néron-Severi lattice. The divisor

$$D_{18} = A_1 + 5A_2 + 2A_3 + 4A_4$$

is ample, of square 18, with  $D_{18} \cdot A_1 = 3$ ,  $D_{18} \cdot A_k = 1$  for  $k \in \{2, 3, 5, \dots, 12\}$ ,  $D_{18} \cdot A_k = 2$  for  $k \in \{4, 13, \dots, 20\}$  and  $D_{18} \cdot A_k = 8$  for  $k > 20$ .

The divisor

$$D_2 = A_1 + 2A_2 + A_3 + A_4$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_2 = D_2 \cdot A_3 = 0$ ,  $D_2 \cdot A_j = 1$  for  $j \in \{5, \dots, 20\}$  and  $D_2 \cdot A_j = 2$  for  $j = 4$  or  $j > 20$ . In the basis  $A_1, \dots, A_{11}$ , we have

$$A_{12} = (-2, 4, 2, 4, -1, -1, -1, -1, -1, -1, -1).$$

Moreover, for  $k \in \{5, \dots, 12\}$ , we have  $A_k + A_{k+8} \equiv A_2 + A_4$  (these are fibers of an elliptic fibration); thus

$$D_2 \equiv A_1 + A_2 + A_3 + A_k + A_{k+8}, \quad \forall k \in \{5, \dots, 12\},$$

and we obtain in that way the classes of  $A_1, \dots, A_{20}$ . Moreover, we see that the surface  $X$  is a double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  which has an  $a_3$  singularity  $q$ . The curves  $A_1, A_2, A_3$  are contracted to  $q$ , the curve  $A_4$  is mapped onto a line that is tangent to the branch of  $C_6$  at  $q$ , and the curves  $A_k, A_{k+8}$  with  $k \in \{5, \dots, 12\}$  are mapped to eight lines going through  $q$  which are tangent to the sextic at any other intersection point. For any subset  $J = \{i, j, k, l\}$  of  $\{5, \dots, 11\}$  of order 4 (there are 35 such choices), let us define

$$\begin{aligned} A_J &= 2A_1 - A_4 + \sum_{t \in J} A_t, \\ B_J &= 4A_2 + 2A_3 + 3A_4 - \sum_{t \in J} A_t. \end{aligned}$$

The classes  $A_J$  and  $B_J$  are the classes of the remaining 70  $(-2)$ -curves  $A_{21}, \dots, A_{90}$ . Moreover, we see that

$$2D_2 \equiv A_J + B_J, \quad \forall J = \{i, j, k, l\} \subset \{5, \dots, 11\}, \#\{i, j, k, l\} = 4,$$

and therefore there exist 35 conics that are 6-tangent to  $C_6$ .

Let  $J, J'$  be two subsets of order 4 of  $\{5, \dots, 11\}$ . The configuration of the curves  $A_J, A_{J'}, B_J, B_{J'}$  is as follows:

$$\begin{aligned} A_J \cdot A_{J'} &= B_J \cdot B_{J'} = 6 - 2\#(J \cap J'), \\ A_J \cdot B_{J'} &= -2 + 2\#(J \cap J'). \end{aligned}$$

**11.4.2. Second involution.**— The divisor  $D'_2 = 2A_2 + A_3 + 2A_4$  is nef, of square 2, with  $D'_2 \cdot A_1 = 2$ ,  $D'_2 \cdot A_j = 1$  for  $j \in \{2, 13, \dots, 20\}$ ,  $D'_2 \cdot A_j = 0$  for  $j \in \{3, \dots, 12\}$  and  $D'_2 \cdot A_j = 4$  for  $j \geq 21$ . We have

$$D'_2 = 2F + A_3,$$

where  $F = A_2 + A_4$  is a fiber of an elliptic fibration such that  $FA_3 = 1$ ; thus the linear system  $|D'_2|$  has base points. Let  $D_8 = 2D'_2$ . The linear system  $|D_8|$  is base-point free, and it is hyperelliptic since  $D_8F = 2$ . One

can check easily that

$$(11.1) \quad \begin{aligned} D_8 &\equiv 2A_1 + \sum_{j=5}^{12} A_j, \\ D_8 &\equiv 4(A_k + A_{k+8}) + 2A_3, \quad k \in \{5, \dots, 12\}, \\ D_8 &\equiv B_J + A_4 + \sum_{t \in J} A_t, \\ D_8 &\equiv A_J + A_4 + A_{12} + \sum_{t \in J^c} A_t, \end{aligned}$$

where  $J = \{i, j, k, l\}$  is a subset of order 4 of  $\{5, \dots, 11\}$  and  $J^c$  is its complement.

By [Sai74, Equation (5.9.1)], the associated map  $\varphi_{D_8} : X \rightarrow \mathbb{P}^5$  has image a cone over a rational normal curve in  $\mathbb{P}^4$ ; it factorizes through a surjective map  $\varphi' : X \rightarrow \mathbb{F}_4$ , where  $\mathbb{F}_4$  is the Hirzebruch surface with a section  $s$  such that  $s^4 = -4$ . The section  $s$  is mapped to the vertex of the cone by the map  $\mathbb{F}_4 \rightarrow \mathbb{P}^5$ . Let  $f$  denote a fiber of the unique fibration  $\mathbb{F}_4 \rightarrow \mathbb{P}^1$ . By [Sai74, Equation (5.9.1)], the branch locus of  $\varphi'$  is the union of  $s$  and a curve  $C$  such that  $Cs = 0$  and  $C \in |3(s + 4f)|$  (so that  $b + C \in |-2K_{\mathbb{F}_4}|$ ). We have

$$|D_8| = \varphi'^* |4f + s|,$$

and therefore from the equivalence relations in (11.1), we get the following:

- The curve  $A_3$  is in the ramification locus; the image by  $\varphi'$  of the curve  $A_3$  is the section  $s$ .
- The curve  $A_1$  is in the ramification locus. Since  $A_1 \cdot A_3 = 0$  and  $A_1$  is a section, the image  $C_1$  of  $A_1$  is in the linear system  $|s + 4f|$ . Let  $B'$  be the curve  $B' \in |2s + 8f|$  such that the branch locus of the double cover  $\varphi : X \rightarrow \mathbb{F}_4$  is

$$B = s + C_1 + B' \in |4s + 12f|.$$

The singular points of the branch locus are nodes  $p_4, \dots, p_{12}$ . The eight points  $p_5, \dots, p_{12}$  are the intersection points of  $C_1$  and  $B'$ ; the image by  $\varphi'$  of  $A_5, \dots, A_{12}$  are the eight points  $p_5, \dots, p_{12}$ . The curve  $B'$  has a node at the point  $p_4$  onto which the curve  $A_4$  is contracted by  $\varphi'$ . The curves  $A_2, A_{13}, \dots, A_{20}$  are sent, respectively, to the fibers passing through  $p_4, p_5, \dots, p_{12}$ .


- The image of the curve  $A_J$  ( $J = \{i, j, k, l\}$ ), is a curve in  $|4f + s|$  passing through the points  $p_t$ , for  $t \in \{4, m, n, o, 12\}$ , where  $\{i, j, k, l, m, n, o\} = \{5, \dots, 11\}$ .
- The image of the curve  $B_J$  ( $J = \{i, j, k, l\}$ ), is a curve in  $|4f + s|$  passing through the points  $p_t$ , for  $t \in \{4, i, j, k, l\}$ .

*Remark 11.4.* By [Kon89], the automorphism group of  $X$  is  $(\mathbb{Z}/2\mathbb{Z})^2$ . It is generated by the involutions associated to the two double covers we described.

## 12. Rank 12 lattices

### 12.1. The lattice $U \oplus E_8 \oplus A_1^{\oplus 2}$

The K3 surface  $X$  contains 14  $(-2)$ -curves  $A_1, \dots, A_{14}$ ; their dual graph is



The curves  $A_1, \dots, A_{12}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{456} = (20, 41, 63, 31, 55, 48, 42, 37, 33, 30, 14, 14)$$

is ample, of square 456, with  $D_{456} \cdot A_j = 1$  for  $j \leq 12$  and  $D_{456} \cdot A_j = 28$  for  $j \in \{13, 14\}$ .

Remark 12.1. The divisors

$$\begin{aligned} F_1 &= 2A_1 + 4A_2 + 6A_3 + 3A_4 + 5A_5 + 4A_6 + 3A_7 + 2A_8 + A_9, \\ F_2 &= A_{11} + A_{13}, \\ F_3 &= A_{12} + A_{14} \end{aligned}$$

are fibers of an elliptic fibration. The number of curves in  $F_1$  counted with multiplicities is 30; thus it is impossible to find an ample divisor  $D$  such that  $D(A_{11} + A_{13}) < 30$ .

By Theorem 2.3, case i) a), we have the following.

**Proposition 12.2.** *The linear system  $|4F_1 + 2A_{10}|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{e}_8$  singularity  $p$  and two nodes  $q, q'$ . The pull-backs of the fibers through  $p, q, q'$  are the fibers  $F_1, F_2, F_3$ .*

We can also construct this surface as follows. In the basis  $A_1, \dots, A_{12}$ , the divisor

$$D_2 = (2, 4, 6, 3, 5, 4, 3, 2, 2, 2, 1, 1)$$


is nef, base-point free, of square 2, with  $D_2 \cdot A_8 = 1$ ,  $D_2 \cdot A_{13} = D_2 \cdot A_{14} = 2$  and  $D_2 \cdot A_j = 0$  for  $j \notin \{8, 13, 14\}$ . Therefore, the K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve which has an  $\mathbf{e}_7$  singularity and a  $\mathbf{d}_4$  singularity. The map  $X \rightarrow \mathbb{P}^2$  is ramified over  $A_8$ . Since

$$D_2 \equiv A_9 + 2A_{10} + 2A_{11} + A_{12} + A_{13} \equiv A_9 + 2A_{10} + A_{11} + 2A_{12} + A_{14},$$

we see that the images of  $A_{13}$  and  $A_{14}$  are the two tangent lines through the two remaining branches of the  $\mathbf{d}_4$  singularities. The sextic curve is the union of a quintic and the line  $L$ . The quintic has a nodal and a cusp singularity, so that with the line  $L$ , they become an  $\mathbf{e}_7$  singularity and a  $\mathbf{d}_4$  singularity.

### 12.2. The lattice $U \oplus D_8 \oplus A_1^{\oplus 2}$

The K3 surface  $X$  contains 19  $(-2)$ -curves  $A_1, \dots, A_{19}$ ; their dual graph is



The curves  $A_1, \dots, A_{12}$  generate the Néron–Severi lattice, and in that basis the divisor

$$D_{136} = (2, 5, 9, 4, 10, 12, 15, 19, 9, 9, 6, 0)$$

is ample, of square 136, with  $D_{136} \cdot A_j = 1$  for  $j \leq 10$ ,  $D_{136} \cdot A_j = 7, 2, 2, 12, 18, 18, 12, 18, 18$  for  $j = 11, \dots, 19$ .

The divisor

$$D_2 = (0, 1, 2, 1, 2, 2, 2, 2, 1, 1, 1, 0)$$

is base-point free, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_8 = 1$ ,  $D_2 \cdot A_j = 0$  for  $j \in \{2, \dots, 7, 9, \dots, 13\}$  and  $D_2 \cdot A_j = 2$  for  $j \in \{14, \dots, 19\}$ . Moreover,

$$\begin{aligned} D_2 &\equiv A_{11} + A_{12} + A_{17} \equiv A_{11} + A_{13} + A_{14}, \\ D_2 &\equiv A_{10} + A_{12} + A_{19} \equiv A_{10} + A_{13} + A_{16}, \\ D_2 &\equiv A_9 + A_{12} + A_{15} \equiv A_9 + A_{13} + A_{18}, \\ D_2 &\equiv 2A_1 + 3A_2 + 4A_3 + 2A_4 + 3A_5 + 2A_6 + A_7 + A_{12} + A_{13}. \end{aligned}$$

By using the linear system  $|D_2|$ , we obtain the following.


**Proposition 12.3.** *The K3 surface  $X$  is a double cover of  $\mathbb{P}^2$  ramified over a sextic curve  $C_6$ ; the curves  $A_1$  and  $A_8$  are in the ramification locus, and their images are two lines. We denote by  $Q_4$  the residual quartic. The sextic curve has a  $\mathbf{d}_6$  singularity (the curves  $A_2, \dots, A_7$  are contracted to that singularity) and five nodal singularities  $p_j$  to which the curves  $A_j$  are contracted, for  $j \in \{9, \dots, 13\}$ .*

The image of  $A_{15}$  (resp.  $A_{17}, A_{19}$ ) is a line through the node  $p_{12}$  and the node  $p_9$  (resp.  $p_{11}, p_{10}$ ) which is tangent to  $Q_4$ . The image of  $A_{14}$  (resp.  $A_{16}, A_{18}$ ) is a line through the node  $p_{13}$  and the node  $p_{11}$  (resp.  $p_{10}, p_9$ ) which is tangent to  $Q_4$ .

### 12.3. The lattice $U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$

The lattice  $U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$  is isometric to  $U \oplus \mathbf{D}_6 \oplus \mathbf{A}_1^{\oplus 4}$ ; see [Kon89].

**12.3.1. First involution.**— The K3 surface  $X$  contains 59  $(-2)$ -curves. The configuration of the first 19  $(-2)$ -curves  $A_1, \dots, A_{19}$  is as follows:



The curves  $A_1, \dots, A_{12}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{40} = (3, 3, 3, 0, 0, 2, 2, 6, 6, 3, 1, 7)$$

is ample, of square 40, with  $D_{40} \cdot A_6 = D_{40} \cdot A_7 = 2$ ,  $D_{40} \cdot A_j = 1$  for  $j \in \{1, 2, 3, 8, \dots, 12, 17, 18, 19\}$  and  $D_{40} \cdot A_j = 7$  for  $j \in \{4, 5, 13, \dots, 16\}$ .

The divisor

$$D_2 = (0, 0, 0, 0, 0, 1, 1, 3, 2, 2, 1, 1)$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_j = 0$  for  $j \in \{8, \dots, 12\}$ ,  $D_2 \cdot A_j = 1$  for  $j \in \{1, \dots, 7, 13, \dots, 19\}$  and  $D_2 \cdot A_j = 2$  for  $j \geq 20$ . The K3 surface  $X$  is a double cover of the plane branched over a sextic curve  $C_6$  which has an  $\mathbf{a}_5$  singularity  $q$ ; the curves  $A_8, \dots, A_{12}$  are contracted to  $q$ . We have

$$D_2 \equiv A_8 + A_9 + A_{10} + A_{11} + A_{12} + F,$$

where

$$\begin{aligned} F &\equiv A_6 + A_7 + 2A_8 + A_9 + A_{10} \\ &\equiv A_1 + A_{14} \equiv A_2 + A_{15} \equiv A_3 + A_{16} \\ &\equiv A_4 + A_{17} \equiv A_5 + A_{18} \equiv A_{13} + A_{19}; \end{aligned}$$

thus there exist seven lines  $L, L_1, \dots, L_6$  through the  $\mathbf{a}_5$  singularity such that  $L$  intersects  $C_6$  in that point only (so  $L$  is the tangent to the branch curve of the singularity; the strict transform of  $L$  is  $A_6 + A_7$ ) and that the lines  $L_i$  have even intersection multiplicities at their other intersection points with the sextic.

The  $(-2)$ -curves on  $X$  are  $A_1, \dots, A_{19}$  and the curves

$$\begin{aligned} B_{ij}^{(1)} &= -A_r - A_s - A_t + 3A_6 + 2A_7 + 6A_8 + 3A_9 + 4A_{10} + 2A_{11}, \\ B_{ij}^{(2)} &= -A_r - A_s - A_t + 2A_6 + 3A_7 + 6A_8 + 3A_9 + 4A_{10} + 2A_{11}, \\ C_{ij}^{(1)} &= -A_i - A_j - A_{13} + 2A_6 + 3A_7 + 6A_8 + 3A_9 + 4A_{10} + 2A_{11}, \\ C_{ij}^{(2)} &= -A_i - A_j - A_{13} + 3A_6 + 2A_7 + 6A_8 + 3A_9 + 4A_{10} + 2A_{11}, \end{aligned}$$

where  $\{i, j, r, s, t\} = \{1, 2, 3, 4, 5\}$ . Using the relation

$$A_{13} = (-1, -1, -1, -1, -1, 3, 3, 6, 2, 4, 2, -2),$$

one can check that  $B_{ij}^{(a)} + C_{ij}^{(a)} = 2D_2$ ; thus the curves  $B_{ij}^{(a)}, C_{ij}^{(a)}$  are pull-backs of conics which are 6-tangent to the sextic curve  $C_6$ .

**12.3.2. Second involution.**— The divisor  $F' = A_6 + A_7 + 2A_8 + A_9 + A_{10}$  is a fiber of an elliptic fibration. The divisor  $D'_2 = 2F' + A_{11}$  is nef, of square 2, with

$$\begin{aligned} D_2 \cdot A_j &= 0 & \text{for } j \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 13\}, \\ D_2 \cdot A_j &= 1 & \text{for } j \in \{10, 14, 15, 16, 17, 18, 19\}, \\ D_2 \cdot A_j &= 4 & \text{for } j \geq 20 \end{aligned}$$

and  $D'_2 \cdot A_{12} = 2$ . The linear system  $|D'_2|$  has base points. The linear system  $|D_8|$  (where  $D_8 = 2D'_2$ ) is base-point free, and it is hyperelliptic since  $D_8 F' = 2$ . One can check easily that for  $i, j, r, s, t$  such that  $\{i, j, r, s, t\} = \{1, \dots, 5\}$ , one has

$$\begin{aligned} D_8 &\equiv B_{ij}^{(1)} + A_r + A_s + A_t + A_6 + 2A_7 + 2A_8 + A_9, \\ D_8 &\equiv B_{ij}^{(2)} + A_r + A_s + A_t + 2A_6 + A_7 + 2A_8 + A_9, \\ D_8 &\equiv C_{ij}^{(1)} + A_i + A_j + A_{13} + 2A_6 + A_7 + 2A_8 + A_9, \\ D_8 &\equiv C_{ij}^{(2)} + A_i + A_j + A_{13} + A_6 + 2A_7 + 2A_8 + A_9. \end{aligned}$$

Moreover,

$$\begin{aligned} D_8 &\equiv 3A_{10} + A_{19} + A_{13} + 3A_6 + 3A_7 + 6A_8 + 3A_9 + 2A_{11}, \\ 2D_8 &\equiv 2A_{10} + \sum_{k=14}^{19} A_k + \sum_{k=1}^5 A_k + 2(A_6 + A_7 + 2A_8 + A_9 + 2A_{11}) + A_{13}, \\ D_8 &\equiv 2A_{12} + \sum_{k=1}^7 A_k + 2A_8 + 2A_9 + A_{13}. \end{aligned}$$

By [Sai74, Equation (5.9.1)], the map  $\varphi_{D_8}: X \rightarrow \mathbb{P}^5$  associated to  $|D_8|$  has image a cone over a rational normal curve in  $\mathbb{P}^4$ ; it factorizes through a surjective map  $\varphi': X \rightarrow \mathbf{F}_4$ , where  $\mathbf{F}_4$  is the Hirzebruch surface with a section  $s$  such that  $s^4 = -4$ . The section  $s$  is mapped to the vertex of the cone by the map  $\mathbf{F}_4 \rightarrow \mathbb{P}^5$ . Let  $f$  denote a fiber of the unique fibration  $\mathbf{F}_4 \rightarrow \mathbb{P}^1$ . By [Sai74, Equation (5.9.1)], the branch locus of  $\varphi'$  is the union of  $s$  and a curve  $C$  such that  $Cs = 0$  and  $C \in |3(s + 4f)|$  (so that  $s + C \in |-2K_{\mathbf{F}_4}|$ ). We have

$$|D_8| = \varphi'^* |4f + s|,$$

and therefore from the above equivalence relations, we get the following:

- The image of curve  $A_{11}$  by  $\varphi'$  is the section  $s$ .
- The branch curve is the union of three components:  $s$ ,  $B'$  and  $C_{12}$ , where  $B' \in |2(s+4f)|$ ,  $C_{12} \in |s+4f|$ . The curve  $B'$  has a node  $q$ , and the curves  $C_{12}$  and  $B'$  meet at  $q$  and at six other points  $p_1, \dots, p_5, p_{13}$ . The singularity at  $q$  of  $B' + C_{12}$  has type  $\mathbf{d}_4$ ; the other singular points are nodes.
- The curves  $A_6, \dots, A_9$  are mapped by  $\varphi'$  to  $q$ .
- The curves  $A_1, \dots, A_5, A_{13}$  are sent to  $p_1, \dots, p_5, p_{13}$ .
- The curve  $A_{12}$  is part of the ramification locus; its image is  $C_{12}$ .
- The curve  $A_{11}$  is part of the ramification locus; its image is  $s$ .
- The curves  $A_{10}, A_{14}, \dots, A_{19}$  are mapped to the fibers through  $q, p_1, \dots, p_5, p_{13}$ .
- The images of the curves  $B_{ij}^{(1)}$  and  $B_{ij}^{(2)}$  are curves in  $|s+4f|$  passing through  $p_r, p_s, p_t$  and through  $q$  with certain tangency properties at the branches of the singularity  $q$ .
- The images of the curves  $C_{ij}^{(1)}$  and  $C_{ij}^{(2)}$  are curves in  $|s+4f|$  passing through  $p_i, p_j, p_{13}$  and through  $q$  with certain tangency properties at the branches.


The 10 curves  $B_{ij}^{(1)}$  (resp.  $B_{ij}^{(2)}, C_{ij}^{(1)}, C_{ij}^{(2)}$ ) for  $\{i, j\} \subset \{1, 2, 3, 4, 5\}$  have the configuration of the Petersen graph, with weight 2 on the edges. Moreover, for  $1 \leq i < j \leq 5$  and  $1 \leq s < t \leq 5$ , the intersections between the four types of curves are as follows:

$$\begin{aligned} B_{ij}^{(1)} \cdot B_{st}^{(2)} &= C_{ij}^{(1)} \cdot C_{st}^{(2)} = 2 + B_{ij}^{(1)} \cdot B_{st}^{(1)}, \\ B_{ij}^{(1)} \cdot C_{st}^{(2)} &= B_{ij}^{(2)} \cdot C_{st}^{(1)} = 2 - B_{ij}^{(1)} \cdot B_{st}^{(1)}, \\ B_{ij}^{(1)} \cdot C_{st}^{(1)} &= B_{ij}^{(2)} \cdot C_{st}^{(2)} = 4 - B_{ij}^{(1)} \cdot B_{st}^{(1)}. \end{aligned}$$

*Remark 12.4.* In [Kon02, Remark 1], Kondo constructed specific surfaces with Néron–Severi lattice isometric to  $U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$  as follows. Let  $C$  be a smooth curve of genus 2 and  $q$  be a point on  $C$ . The linear system  $|K_C + 2q|$  gives a plane quartic curve with a cusp. The minimal resolution  $Y$  of the cyclic degree 4 cover of  $\mathbb{P}^2$  branched over that curve has an elliptic fibration (obtained by blowing up the cusp) with a  $\tilde{\mathbf{D}}_4$  and six  $\tilde{\mathbf{A}}_1$  fibers. The automorphism group of such a surface is larger than the general one.

### 12.4. The lattice $U \oplus \mathbf{A}_2 \oplus \mathbf{E}_8$

The K3 surface  $X$  contains 13  $(-2)$ -curves  $A_1, \dots, A_{13}$ ; their configuration is as follows:



The curves  $A_1, \dots, A_{12}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{698} = (38, 77, 58, 117, 100, 84, 69, 55, 42, 30, 19, 9)$$

is ample, of square 698, with  $D_{698} \cdot A_j = 1$  for  $j \leq 12$  and  $D_{698} \cdot A_{13} = 28$ . The divisor

$$F_1 = A_{11} + A_{12} + A_{13}$$

is a fiber of an elliptic fibration with section  $A_{10}$ . That fibration has another singular fiber  $F_2$  of type  $\tilde{\mathbf{E}}_8$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 12.5.** *The linear system  $|4F_1 + 2A_{10}|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one cusp  $p$  and one  $\mathbf{e}_8$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

We can also give another construction as a double cover of  $\mathbb{P}^2$ : The divisor

$$D_2 = (2, 5, 4, 8, 7, 6, 5, 4, 3, 2, 1, 0)$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_1 = D_2 \cdot A_{12} = D_2 \cdot A_{13} = 1$  and  $D_2 \cdot A_j = 0$  for  $j \in \{2, \dots, 11\}$ . The K3 surface is the double cover of  $\mathbb{P}^2$  branched over a sextic curve with a  $\mathbf{d}_{10}$  singularity. The curve  $A_1$  is in the ramification locus; we denote by  $L$  its image and by  $Q$  the residual quintic curve. The quintic  $Q$  has a node, and  $L$  is tangent with multiplicity 4 at a branch of that node. We have


$$D_2 \equiv A_2 + A_3 + 2A_4 + 2A_5 + 2A_6 + 2A_7 + 2A_8 + 2A_9 + 2A_{10} + 2A_{11} + A_{12} + A_{13};$$

thus the image of  $A_{12}$  and  $A_{13}$  is the line which is the tangent to the other branch of the node, which is moreover tangent to  $Q$  at another point.

### 13. Rank 13 lattices

#### 13.1. The lattice $U \oplus E_8 \oplus A_1^{\oplus 3}$

The K3 surface  $X$  contains 17  $(-2)$ -curves  $A_1, \dots, A_{17}$ ; their configuration is as follows:



The curves  $A_1, \dots, A_{13}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{294} = (2, 5, 4, 9, 10, 12, 15, 19, 24, 30, 14, 14, 9)$$

is ample, of square 294, with  $D_{294} \cdot A_j = 1$  for  $j \leq 10$ ,  $D_{294} \cdot A_j = 2$  for  $j \in \{11, 12, 17\}$ ,  $D_{294} \cdot A_j = 28$  for  $j \in \{14, 15\}$ ,  $D_{294} \cdot A_{13} = 12$  and  $D_{294} \cdot A_{16} = 18$ . By considering the elliptic fibration  $s$  with section  $A_{10}$  and the fibers

$$F = A_{11} + A_{14}, A_{12} + A_{15}, A_{13} + A_{16}$$

plus the fiber of type  $\tilde{E}_8$  supported on  $A_1, \dots, A_9$ , we get the following.

**Proposition 13.1.** *The surface is a double cover of the Hirzebruch surface  $F_4$  branched over the negative section  $s$  and a curve  $B \in |3(s + 4f)|$  with three nodes and an  $\mathbf{e}_8$  singularity.*

The double cover  $\eta$  is given by the linear system  $|D_8|$ , where  $D_8 = 4F + 2A_{10}$ . One has

$$D_8 = A_{17} + 4A_1 + 7A_2 + 5A_3 + 10A_4 + 8A_5 + 6A_6 + 4A_7 + 2A_8 + A_{14} + A_{15} + A_{16};$$

thus the image by  $\eta$  of the curve  $A_{17}$  is a curve in  $|s + 4f|$  going through the three nodes plus the  $\mathbf{e}_8$  singularity and infinitely near points of it.

We can also construct the surface  $X$  as follows. The divisor

$$D_2 = (0, 1, 1, 2, 2, 2, 2, 2, 2, 1, 1, 1)$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_j = 0$  for  $j \in \{2, \dots, 9, 11, 12, 13, 17\}$ . We have the relations

$$D_2 \equiv A_{13} + A_{16} + A_{17} \equiv A_{12} + A_{15} + A_{17} \equiv A_{11} + A_{14} + A_{17}$$

and

$$D_2 \equiv 2A_1 + 4A_2 + 3A_3 + 6A_4 + 5A_5 + 4A_6 + 3A_7 + 2A_8 + A_9 + A_{17}.$$




The curves  $A_1$  and  $A_{10}$  are part of the ramification locus; their images are lines  $L, L'$ . Let us denote by  $Q$  the residual quartic. The images of the curves  $A_{14}, A_{15}, A_{16}$  are lines  $L_1, L_2, L_3$  going through the images of  $A_{17}$  and of  $A_{11}, A_{12}, A_{13}$ , respectively. The quartic curve is smooth and tangent to the line  $L$  (the image of  $A_1$ ) at the intersection point of  $L$  and  $L'$ , so that the singularity at  $q$  is  $\mathbf{d}_8$ . The branch locus has four nodes, three on the line  $L'$  (the image of  $A_{10}$ ) and one on  $L$  (the image of  $A_1$ ).

### 13.2. The lattice $U \oplus D_8 \oplus A_1^{\oplus 3}$

The lattice  $U \oplus D_8 \oplus A_1^{\oplus 3}$  is also isometric to  $U \oplus E_7 \oplus A_1^{\oplus 4}$  and  $U \oplus D_6 \oplus D_4 \oplus A_1$ ; see [Kon89].

**13.2.1. The first involution.**— The K3 surface  $X$  contains 39  $(-2)$ -curves  $A_1, \dots, A_{39}$ . The configuration of the curves  $A_1, \dots, A_{19}$  is as follows:



The curves  $A_1, \dots, A_{13}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{86} = (2, 1, 2, 5, 5, 11, 8, 6, 2, 3, 1, 0, 1)$$

is ample, of square 86. The divisor

$$D_2 = (-1, 1, 1, 1, 1, 3, 3, 3, 1, 2, 1, -1, -2)$$

is nef, of square 2, base-point free, with  $D_2 \cdot A_j = 0$  for  $j \in \{6, 7, 8, 10, 11, 12, 18\}$ ,  $D_2 \cdot A_j = 2$  for  $j \geq 20$  and else  $D_2 \cdot A_j = 1$ . For  $k \in \{10, \dots, 19\}$ , one has

$$2D_2 \equiv A_{2k} + A_{2k+1},$$

and using the equivalences from the elliptic fibration with fiber  $A_1 + A_{13}$ , we get

$$\begin{aligned} D_2 &\equiv A_{13} + A_1 + A_6 + A_7 + A_8 + A_{10} + A_{11} + A_{12} + A_{18}, \\ D_2 &\equiv A_{14} + A_2 + A_6 + A_7 + A_8 + A_{10} + A_{11} + A_{12} + A_{18}, \\ D_2 &\equiv A_{15} + A_3 + A_6 + A_7 + A_8 + A_{10} + A_{11} + A_{12} + A_{18}, \\ D_2 &\equiv A_{16} + A_4 + A_6 + A_7 + A_8 + A_{10} + A_{11} + A_{12} + A_{18}, \\ D_2 &\equiv A_{17} + A_5 + A_6 + A_7 + A_8 + A_{10} + A_{11} + A_{12} + A_{18}, \\ D_2 &\equiv A_9 + A_{19} + A_6 + 2A_7 + 3A_8 + 3A_{10} + 3A_{11} + A_{12} + 2A_{18}. \end{aligned}$$


The K3 surface is the double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with a singularity  $q$  of type  $\mathbf{a}_7$ . The line tangent to the branch of  $C_6$  has no other intersection points with  $C_6$ , and the curves  $A_9, A_{19}$  map onto that line. Moreover, there exist five lines through  $q$  that are bitangent to the sextic at their other intersection points. The images of  $A_j, A_{j+12}$  for  $j \in \{1, \dots, 5\}$  are these lines.

For  $J = \{i, j\} \subset \{1, \dots, 5\}$ , let

$$A_J = -A_i - A_j + \sum_{t=1}^5 A_t + 2A_6 + A_7 - A_9.$$

Then  $A_J$  is the class of a  $(-2)$ -curve, and  $B_J = 2D_2 - A_J$  is also a  $(-2)$ -curve. These are the 20 curves  $A_{20}, \dots, A_{39}$ ; thus the images by the double cover map of curves  $A_J, B_J$  are conics that are 6-tangent to the sextic branch curve.

We have  $A_J \cdot A_{J'} = 2$  if and only if  $\#J \cap J' = 0$ , and else  $A_J \cdot A_{J'} \in \{-2, 0\}$ . The dual graph of the 10 curves  $A_J$  (for  $J \subset \{1, \dots, 5\}$ ,  $\#J = 2$ ) is the Petersen graph with weight 2 on the edges:



The configuration of the curves  $B_J$  ( $J \subset \{1, \dots, 5\}$ ,  $\#J = 2$ ) is also the Petersen graph with weight 2 on the edges. Using that  $D_2 \cdot A_J = D_2 \cdot B_J = 2$ , we get that  $A_J \cdot A_{J'} = B_J \cdot B_{J'}$  and  $A_J \cdot B_{J'} = 4 - A_J \cdot A_{J'}$  for any subsets  $J, J'$  of order 2 of  $\{1, \dots, 5\}$ .

### 13.2.2. The second involution.— The divisor

$$D'_2 = 2A_1 + A_{12} + 2A_{13}$$

is nef, of square 2, with base points. One has  $D_2 \cdot A_j = 0$  for  $j \in \{1, \dots, 5, 7, \dots, 12, 19\}$ ,  $D'_2 \cdot A_6 = 2$ ,  $D'_2 \cdot A_j = 1$  for  $j \in \{13, \dots, 18\}$  and  $D'_2 \cdot A_j = 4$  for  $j \geq 20$ . Since  $D'_2 = 2F + A_{12}$ , where  $F = A_1 + A_{13}$  is a fiber of an elliptic fibration with  $FA_{12} = 1$ , the linear system  $|D'_2|$  has base points, and  $|D_8| = |2D'_2|$  is base-point free and hyperelliptic. We have

$$(13.1) \quad \begin{aligned} D_8 &\equiv A_{13} + A_{14} + A_{15} + A_{16} + A_1 + A_2 + A_3 + A_4 + 2A_{12}, \\ D_8 &\equiv A_{15} + A_{16} + A_{17} + A_{18} + A_3 + A_4 + A_5 + A_7 + 2A_8 \\ &\quad + A_9 + 2A_{10} + 2A_{11} + 2A_{12} + A_{19}, \\ D_8 &\equiv B_J + \sum_{t=1, t \notin J}^5 A_t + A_7 + 2A_8 + A_9 + 2A_{10} + 2A_{11} + 2A_{19}, \\ D_8 &\equiv A_J + \sum_{t \in J} A_t + 2A_7 + 4A_8 + 3A_9 + 3A_{10} + 2A_{11}, \\ D_8 &= 2A_6 + \sum_{t=1}^5 A_t + 3A_7 + 4A_8 + 2A_9 + 3A_{10} + 2A_{11} + A_{19}. \end{aligned}$$

The map  $\varphi_{D_8} : X \rightarrow \mathbb{P}^5$  associated to  $|D_8|$  has image a cone over a rational normal curve in  $\mathbb{P}^4$ ; it factorizes through a surjective map  $\varphi' : X \rightarrow \mathbf{F}_4$ , where  $\mathbf{F}_4$  is the Hirzebruch surface with a section  $s$  such that  $s^4 = -4$ . The section  $s$  is mapped to the vertex of the cone by the map  $\mathbf{F}_4 \rightarrow \mathbb{P}^5$ . Let  $f$  denote a fiber of the unique fibration  $\mathbf{F}_4 \rightarrow \mathbb{P}^1$ . By Theorem 2.3, the branch locus of  $\varphi'$  is the union of  $s$  and a curve  $C$  such that  $Cs = 0$  and  $C \in |3(s + 4f)|$  (so that  $b + C \in |-2K_{\mathbf{F}_4}|$ ). We have

$$|D_8| = \varphi'^* |4f + s|,$$


and therefore from the equivalence relations in (13.1), we get the following claims.

- The curve  $A_{12}$  is in the ramification locus; its image is  $s$ .
- The curves  $A_7, A_8, A_9, A_{10}, A_{11}, A_{19}$  are contracted to a  $\mathbf{d}_6$  singularity  $q$  of  $C$ .
- The curve  $C$  is the union of a curve  $s'$  in  $|s + 4f|$  and a curve  $B'$  in  $|2(s + 4f)|$  (thus  $s'B' = 8$ ). The curve  $B'$  has a node at  $q$ , and  $s'$  is tangent to one of the branches of  $B'$  at  $q$ , so that the intersection multiplicity of  $s'$  and  $B'$  at  $q$  is 3 and the singularity of  $C = B' + s'$  has type  $\mathbf{d}_6$ .

- The other intersection points of  $s'$  and  $B'$  are nodes  $p_1, \dots, p_5$  to which the curves  $A_1, \dots, A_5$  are mapped.
- The curve  $A_6$  is in the ramification locus; its image is  $s'$ .
- The curves  $A_{13}, \dots, A_{17}, A_{18}$  are mapped to the fibers through  $p_1, \dots, p_5$  and  $q$ , respectively.
- The curves  $A_j$  and  $B_j$  are mapped to curves in the linear system  $|s + 4f|$  passing through the points  $p_1, \dots, p_5$  and points infinitely near  $q$  with certain multiplicities.

### 13.3. The lattice $U \oplus E_8 \oplus A_3$

The K3 surface  $X$  contains 14  $(-2)$ -curves  $A_1, \dots, A_{14}$ ; their configuration is as follows:



The curves  $A_1, \dots, A_{13}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{506} = (46, 93, 141, 70, 120, 100, 81, 63, 46, 30, 15, 1, -12)$$

is ample, of square 506, with  $D_{506} \cdot A_j = 1$  for  $j \leq 12$ ,  $D_{506} \cdot A_{13} = 25$  and  $D_{506} \cdot A_{14} = 3$ . The surface has an elliptic fibration such that  $A_{10}$  is a section with singular fiber

$$F_1 = A_{11} + A_{12} + A_{13} + A_{14}$$

and a fiber  $F_2$  of type  $\tilde{E}_8$ . By Theorem 2.3, case i) a), we have the following.

**Proposition 13.2.** *The linear system  $|4F_1 + 2A_8|$  defines a morphism  $\varphi: X \rightarrow \mathbf{F}_4$  branched over the unique section  $s$  with  $s^2 = -4$  with  $s^2 = -4$  and a curve  $B \in |3s + 12f|$ . The curve  $B$  has one  $\mathbf{a}_3$  singularity  $p$  and one  $\mathbf{e}_8$  singularity  $q$ . The pull-backs of the fibers through  $p, q$  are the fibers  $F_1, F_2$ .*

In order to construct  $X$ , one can also use the divisor

$$D_2 = (2, 5, 8, 4, 7, 6, 5, 4, 3, 2, 1, 0, 0),$$

which is nef, of square 2, base-point free, with  $D_2 \cdot A_j = 0$  for  $j \in \{2, \dots, 11, 13\}$  and  $D_2 \cdot A_j = 1$  for  $j \in \{1, 12, 14\}$ . Since


$$D_2 \equiv A_2 + 2A_3 + A_4 + 2A_5 + 2A_6 + 2A_7 + 2A_8 + A_9 + 2A_{10} + 2A_{11} + A_{12} + A_{13} + A_{14},$$

we obtain that the K3 surface is the double cover of  $\mathbb{P}^2$  branched over a sextic curve which is the union of a line  $L$  and a quintic  $Q$  such that  $A_1$  is in the ramification locus and its image is  $L$ . The sextic curve has a singularity  $q$  of type  $\mathbf{d}_{10}$  and a node. The image of  $A_{12}$  and  $A_{14}$  is the line  $L$ . The situation is a specialization of Section 12.4.

## 14. Rank 14 lattices

### 14.1. The lattice $U \oplus E_8 \oplus D_4$

The K3 surface  $X$  contains 15  $(-2)$ -curves  $A_1, \dots, A_{15}$ ; their dual graph is



The curves  $A_1, \dots, A_{14}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{506} = (46, 93, 141, 70, 120, 100, 81, 63, 46, 30, 15, 1, -12, 0)$$

is ample, of square 506, with  $D_{506} \cdot A_{13} = 25$  and  $D_{506} \cdot A_j = 1$  for  $j \neq 13$ . The divisor

$$D_2 = (2, 5, 8, 4, 7, 6, 5, 4, 3, 2, 1, 0, 0, 0)$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_1 = D_2 \cdot A_{12} = 1$  and  $D_2 \cdot A_j = 0$  for  $j \notin \{1, 12\}$ . We have


$$D_2 \equiv A_2 + 2A_3 + A_4 + 2(A_5 + A_6 + A_7 + A_8 + A_9 + A_{10} + A_{11} + A_{12}) + A_{13} + A_{14} + A_{15};$$

thus, using the linear system  $|D_2|$ , we obtain the following.

**Proposition 14.1.** *The K3 surface is a double cover of  $\mathbb{P}^2$  which is ramified over  $A_1$  and  $A_{12}$ ; the images of these curves are lines in the sextic branch curve. The sextic curve has a  $\mathbf{d}_{10}$  singularity at a point  $q$  and three nodal singularities. The residual quartic curve is smooth at  $q$ . The tangent at  $q$  is the line  $L$ , and the order of tangency at  $q$  is 4. The other nodes are the intersections of the line  $L'$  with the quartic.*

## 14.2. The lattice $U \oplus D_8 \oplus D_4$

**14.2.1. First involution.**— The K3 surface  $X$  contains 20  $(-2)$ -curves  $A_1, \dots, A_{20}$ ; their configuration is as follows:



The curves  $A_1, \dots, A_{14}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{154} = (1, 3, 6, 10, 7, 15, 14, 14, 15, 17, 8, 8, 0, 4)$$

is ample, of square 154, with  $D_{154} \cdot A_j = 1$  for  $j \in \{1, 1, \dots, 12, 18, 20\}$ ,  $D_{154} \cdot A_j = 9$  for  $j \in \{14, 19\}$  and  $D_{154} \cdot A_j = 17$  for  $j \in \{15, 16, 17\}$ . The divisor

$$D_2 = (1, 2, 3, 4, 2, 5, 4, 3, 2, 1, 0, 0, 0, 0)$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_j = 0$  for  $j \in \{1, \dots, 4, 6, \dots, 10\}$  and else  $D_2 \cdot A_j = 1$ . Using the equivalences from the elliptic fibration with fiber  $A_{11} + A_{16}$ , we obtain that

$$D_2 \equiv A_{10+k} + A_{15+k} + \sum_{j=1, j \neq 5}^{10} A_j, \forall k \in \{1, 2, 3, 4, 5\}.$$

The linear system  $|D_2|$  defines the K3 surface  $X$  as the double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with an  $\mathbf{a}_9$  singularity  $q$ . The cover is ramified above  $A_5$ ; the image of  $A_5$  is a line, a component of  $C_6$  which has no other intersection points with the residual quintic curve  $Q$ . The images of  $A_{10+k} + A_{15+k}$ ,  $k \in \{1, \dots, 5\}$ , are lines through  $q$  which are bitangent to  $Q$ . The quintic is smooth, thus of genus 6.

14.2.2. **Second involution.**— The divisor

$$D'_2 = (0, 1, 2, 4, 3, 6, 5, 4, 3, 2, 1, 0, 0, 0)$$

is nef, base-point free, of square 2. It satisfies  $D_2 \cdot A_j = 0$  for  $j \in \{2, 4, \dots, 11, 17, \dots, 20\}$ . Using equivalence relations obtained from the elliptic fibration with fiber  $A_{11} + A_{16}$ , we have

$$\begin{aligned} D'_2 &\equiv A_{12} + (A_{17} + A_4 + A_5 + 2 \sum_{t=6}^{10} A_t + A_{11}), \\ D'_2 &\equiv A_{13} + (A_{18} + A_4 + A_5 + 2 \sum_{t=6}^{10} A_t + A_{11}), \\ D'_2 &\equiv A_{14} + (A_{19} + A_4 + A_5 + 2 \sum_{t=6}^{10} A_t + A_{11}), \\ D'_2 &\equiv A_{15} + (A_{20} + A_4 + A_5 + 2 \sum_{t=6}^{10} A_t + A_{11}), \\ D'_2 &\equiv A_{16} + (A_{11} + A_4 + A_5 + 2 \sum_{t=6}^{10} A_t + A_{11}); \end{aligned}$$

moreover,

$$D'_2 \equiv 2A_1 + A_2 + \sum_{j=17}^{20} A_j.$$

We have the following assertions.


- The branch curve of the corresponding double cover is a sextic with a  $\mathbf{d}_8$  singularity at a point  $q$  onto which the curves  $A_4, \dots, A_{11}$  are contracted and five nodes  $p_2, p_{17}, \dots, p_{20}$  onto which  $A_2, A_{17}, \dots, A_{20}$  are contracted. The curves  $A_1, A_3$  are in the ramification locus; their images are lines  $L_1, L_3$ . Let us denote by  $Q'$  the residual quartic curve.
- The line  $L_1$  cuts  $Q'$  (resp.  $L_3$ ) at the points  $p_{17}, \dots, p_{20}$  (resp. the point  $p_2$ ).
- The quartic  $Q'$  has a node at  $q$ , and the line  $L_3$  is tangent with multiplicity 3 at one of the branches of the node.
- For  $k \in \{12, \dots, 15\}$ , the image of  $A_k$  is a line through  $q$  and  $p_{k+5}$ .
- The image of  $A_{16}$  is a line passing through  $q$  and tangent to  $Q'$  at another point.

*Remark 14.2.* We constructed the surface  $X$  as two double covers. The associated involutions fix curves of geometric genus of 6 and 2, respectively; thus these involution generate the automorphism group of  $X$ , which is  $(\mathbb{Z}/2\mathbb{Z})^2$  according to [Kon89].

14.3. **The lattice  $U \oplus E_8 \oplus A_1^{\oplus 4}$**

The lattice  $U \oplus E_8 \oplus A_1^{\oplus 4}$  is also isometric to  $U \oplus E_7 \oplus D_4 \oplus A_1$ ; see [Kon89].

14.3.1. **First involution.**— The K3 surface  $X$  contains 27  $(-2)$ -curves  $A_1, \dots, A_{27}$ . The dual graph of  $A_1, \dots, A_{19}$  and the intersections of the curves  $A_1, \dots, A_{11}$  with the curves  $A_j$  for  $j \geq 20$  is given by



The curves  $A_1, \dots, A_{14}$  generate the Néron–Severi lattice. In that base, the divisor

$$D_{132} = (2, 5, 9, 4, 10, 12, 15, 19, 9, 15, 12, 5, 5, 0)$$

is ample, of square 132, with  $D_{132} \cdot A_j = 1$  for  $j \leq 11$ ,  $D_{132} \cdot A_j = 2$  for  $j \in \{12, 13, 18, 19\}$ ,  $D_{132} \cdot A_j = 12$  for  $j \in \{15, \dots, 18\}$ ,  $D_{132} \cdot A_j = 18$  for  $j \in \{21, 23, 24, 26\}$  and  $D_{132} \cdot A_j = 28$  for  $j \in \{20, 22, 25, 27\}$ . The divisor

$$D_2 = (1, 2, 3, 1, 3, 3, 3, 3, 1, 2, 1, 0, 0, 0)$$

is nef, of square 2, base-point free. We have  $D_2 \cdot A_j = 0$  for  $j \in \{1, 2, 3, 5, 6, 7, 8, 10, 11\}$ ,  $D_2 \cdot A_j = 1$  for  $j \in \{4, 9, 12, \dots, 19\}$  and  $D_2 \cdot A_j = 2$  for  $j \in \{20, \dots, 27\}$ . Using the elliptic fibration with fiber  $A_{12} + A_{16}$ , we get

$$D_2 \equiv A_{11+k} + A_{15+k} + \left( -A_4 - A_9 + \sum_{j=1}^{11} A_j \right), k \in \{1, 2, 3, 4\}.$$

Moreover, one has


$$2D_2 \equiv A_{2k} + A_{2k+1} \quad \text{for } k \in \{10, 11, 12, 13\};$$

thus the K3 surface is the double cover branched over a sextic curve with an  $a_9$  singularity  $q$ . The image of  $A_4$  and  $A_9$  is the line tangent to the branch of that singularity. The curves  $A_{11+k} + A_{15+k}$ ,  $k \in \{1, 2, 3, 4\}$ , are mapped onto lines going through  $q$  that are tangent to the sextic curve in two other points. The curves  $A_{2k} + A_{2k+1}$  for  $k \in \{10, 11, 12, 13\}$  are mapped onto conics that are 6-tangent to the sextic curve.


The classes of the curves  $A_{20}, A_{22}, A_{24}, A_{26}$  are

$$\begin{aligned} A_{20} &= (0, 0, 0, -1, 1, 2, 3, 4, 2, 3, 2, 0, 0, 1), & A_{22} &= (2, 4, 6, 2, 6, 6, 6, 6, 3, 3, 0, -1, -1, -1) \\ A_{24} &= (0, 0, 0, -1, 1, 2, 3, 4, 2, 3, 2, 1, 0, 0), & A_{26} &= (0, 0, 0, -1, 1, 2, 3, 4, 2, 3, 2, 0, 1, 0), \end{aligned}$$

and the dual graph of the curves  $A_{20}, \dots, A_{27}$  is



where a thick line has weight 6 and a thin line has weight 4. For completeness, we also give the intersections between the curves  $A_{12}, \dots, A_{19}$  and the curves  $A_{20}, \dots, A_{27}$ :



where a thin line has weight 2. In fact, one can check that the above graph is another occurrence of the Levi graph of the Möbius configuration.

**14.3.2. Second involution.**— The divisor

$$D'_2 = (1, 2, 4, 2, 4, 4, 4, 4, 2, 2, 0, 0, 0, 0)$$

is nef, of square 2, with base points and with  $D_2 \cdot A_j = 0$  for  $j \in \{1, 3, \dots, 10, 12, \dots, 15\}$  and  $D_2 \cdot A_j = 1$  for  $j \in \{2, 16, 17, 18, 19\}$ ,  $D_2 \cdot A_j = 4$  for  $j \geq 20$ . Let  $D_8 = 2D'_2$ ; it is base-point free and hyperelliptic. It defines a morphism  $\varphi': X \rightarrow \mathbb{P}^5$  onto a degree 4 surface. That morphism factors through the Hirzebruch surface  $F_4$  by a map denoted by  $\varphi$ ; the map  $F_4 \rightarrow \mathbb{P}^5$  contracts the section  $s$  of the Hirzebruch surface. The divisor

$$F_1 = A_2 + A_4 + 2(A_3 + A_5 + A_6 + A_7 + A_8) + A_9 + A_{10}$$

is the fiber of an elliptic fibration for which the curves  $A_{11+k} + A_{15+k}$ ,  $k \in \{1, 2, 3, 4\}$ , are also fibers. One has  $D'_2 = 2F_1 + A_1$ ; thus

$$D_8 \equiv \sum_{j=12}^{19} A_j + 2A_1.$$

Moreover, we have

$$\begin{aligned} D_8 &\equiv 2A_{11} + 2A_3 + A_4 + 3A_5 + 4A_6 + 5A_7 + 6A_8 + 3A_9 + 4A_{10}, \\ D_8 &\equiv A_{20} + 2\sum_{j=3}^8 A_j + A_9 + A_{10} + A_{12} + A_{13} + A_{15}, \\ D_8 &\equiv A_{22} + 2\sum_{j=3}^8 A_j + A_9 + A_{10} + A_{12} + A_{13} + A_{14}, \\ D_8 &\equiv A_{24} + 2\sum_{j=3}^8 A_j + A_9 + A_{10} + A_{13} + A_{14} + A_{15}, \\ D_8 &\equiv A_{26} + 2\sum_{j=3}^8 A_j + A_9 + A_{10} + A_{12} + A_{14} + A_{15}, \\ D_8 &\equiv A_{21} + 2A_3 + A_4 + 3A_5 + 4A_6 + 5A_7 + 6A_8 + 4A_9 + 3A_{10} + A_{14}, \\ D_8 &\equiv A_{23} + 2A_3 + A_4 + 3A_5 + 4A_6 + 5A_7 + 6A_8 + 4A_9 + 3A_{10} + A_{15}, \\ D_8 &\equiv A_{25} + 2A_3 + A_4 + 3A_5 + 4A_6 + 5A_7 + 6A_8 + 4A_9 + 3A_{10} + A_{12}, \\ D_8 &\equiv A_{27} + 2A_3 + A_4 + 3A_5 + 4A_6 + 5A_7 + 6A_8 + 4A_9 + 3A_{10} + A_{13}. \end{aligned}$$

We therefore have the following claims.

- The curves  $A_3, \dots, A_{10}$  are contracted to a  $\mathbf{d}_8$  singularity  $q$  of the branch curve  $B$  of  $\varphi$ ; the curves  $A_{12}, \dots, A_{15}$  are mapped to nodes  $p_{12}, \dots, p_{15}$  of  $B$ .
- The curve  $B$  is the union of  $s$  and a curve  $C$  such that  $Cs = 0$  and  $C \in |3(s + 4f)|$  (so that  $s + C \in |-2K_{F_4}|$ ). We have

$$|D_8| = \varphi^*|4f + s|$$


and therefore from the above equivalence relations, we get the following:

- The image of curve  $A_1$  by  $\varphi$  is the section  $s$ .
- The curve  $B$  is the union of three components:  $s$ ,  $B'$  and  $C_{11}$ , where  $B' \in |2(s + 4f)|$  and  $C_{11} \in |s + 4f|$ . The curve  $B'$  has a node  $q$ , and the curves  $C_{11}$  and  $B'$  meet at  $q$  in such a way that  $C_{11}$  intersect one of the branches with multiplicity 3 (so that the singularity at  $q$  of  $B' + C_{11}$  has type  $\mathbf{d}_8$ ). The four points  $p_{12}, \dots, p_{15}$  are nodes; they are the remaining intersection points of  $C_{11}$  and  $B'$  (so that  $B'C_{11} = 8$ ).
- The images of the curves  $A_{16}, \dots, A_{19}$  are the fibers through  $p_{12}, \dots, p_{15}$ ,
- The images of the curves  $A_{20}, A_{22}, A_{24}, A_{26}$  by  $\varphi$  are curves in the linear system  $|s + 4f|$  passing through three of the four points  $p_{12}, \dots, p_{15}$ , and through  $q$  and points infinitely near  $q$  with certain multiplicities.
- The images of the curves  $A_{21}, A_{23}, A_{25}, A_{27}$  by  $\varphi$  are curves in the linear system  $|s + 4f|$  passing through one of the four points  $p_{12}, \dots, p_{15}$  and through  $q$  and points infinitely near  $q$  with certain multiplicities.

## 15. Rank at least 15 lattices

### 15.1. The lattice $U \oplus E_8 \oplus D_4 \oplus A_1$

The K3 surface  $X$  contains 21  $(-2)$ -curves  $A_1, \dots, A_{21}$ . The dual graph of these curves is given in [Kon89, Section 4]; we reproduce it here:



The curves  $A_1, \dots, A_{15}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{242} = (2, 5, 9, 4, 10, 12, 15, 19, 24, 30, 14, 23, 17, 8, 4)$$

is ample, of square 242, with  $D_{242} \cdot A_j = 1$  for  $j \in \{1, \dots, 14\} \setminus \{11\}$ .

The divisor

$$D_2 = (1, 2, 3, 1, 3, 3, 3, 3, 3, 3, 1, 2, 1, 0, 0)$$

is nef, base-point free, with  $D_2^2 = 2$ ,  $D_2 \cdot A_{20} = D_2 \cdot A_{21} = 2$ ,  $D_2 \cdot A_j = 1$  for  $j \in \{4, 11, 14, \dots, 19\}$  and else  $D_2 \cdot A_j = 0$ . We have

$$2D_2 \equiv A_{20} + A_{21}$$

and

$$D_2 \equiv \sum_{k \in \{1, \dots, 13\} \setminus \{4, 11\}} A_k + A_{13+j} + A_{16+j}, \quad j \in \{1, 2, 3\}.$$

By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 15.1.** *The K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve  $C_6$  with an  $\mathbf{a}_{11}$  singularity  $q$ ; the curves  $A_j$  with  $j \in \{1, \dots, 13\} \setminus \{4, 11\}$  are mapped to  $q$ . The image of the curves  $A_4$  and  $A_{11}$  is the line  $L$  tangent to the branch of  $C_6$  at  $q$ . The images of  $A_{13+j}$  and  $A_{16+j}$ ,  $j \in \{1, 2, 3\}$ , are lines  $L_1, L_2, L_3$  through  $q$  that are bitangent to the sextic at other intersection points. The image of  $A_{20}, A_{21}$  is a conic which is 6-tangent to the sextic.*

*Remark 15.2.* The branch curve is irreducible, with geometric genus 4.

The divisor  $D'_2 = (0, 1, 2, 1, 2, 2, 2, 3, 4, 2, 3, 2, 1, 0)$  is nef, base-point free of square 2, with


$$D'_2 \cdot A_1 = D'_2 \cdot A_8 = 1, \quad D'_2 \cdot A_{15} = D'_2 \cdot A_{16} = D'_2 \cdot A_{17} = 2, \quad D'_2 \cdot A_{20} = D'_2 \cdot A_{21} = 4.$$

Therefore, the linear system  $|D'_2|$  induces a double cover  $\varphi': X \rightarrow \mathbb{P}^2$  which is ramified above the line  $L' = \varphi'(A_8)$ ; the curves  $A_2, A_3, A_5, A_6, A_7$  are contracted to a singularity of the branch curve, and the curves  $A_9, \dots, A_{14}$  are contracted to another singularity. In that way, we obtain a second involution acting on the K3 surface. By [Kon89], we know that the automorphism group of the general K3 is  $(\mathbb{Z}/2\mathbb{Z})^2$ .

## 15.2. The lattice $U \oplus E_8 \oplus D_6$

The K3 surface  $X$  contains 19  $(-2)$ -curves  $A_1, \dots, A_{19}$ . The dual graph of these curves is (see [Kon89, Section 4])





The curves  $A_1, \dots, A_{16}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{304} = (20, 41, 63, 31, 55, 48, 42, 37, 33, 30, 28, 27, 13, 14, 2, -9)$$

is ample, of square 304, with  $D_{304} \cdot A_j = 1$  for  $j \leq 15$ ,  $D_{304} \cdot A_{16} = D_{304} \cdot A_{19} = 20$  and  $D_{304} \cdot A_{17} = D_{304} \cdot A_{18} = 2$ . The divisor

$$D_2 = (1, 2, 3, 1, 3, 3, 3, 3, 3, 3, 3, 3, 1, 2, 1, 0)$$

is nef, base-point free, of square 2, with  $D_2 \cdot A_j = 1$  for  $j \in \{4, 13, 16, 17, 18, 19\}$  and else  $D_2 \cdot A_j = 0$ . One has

$$\begin{aligned} D_2 &\equiv \left(-A_4 - A_{13} + \sum_{j=1}^{15} A_j\right) + A_{16} + A_{17}, \\ D_2 &\equiv \left(-A_4 - A_{13} + \sum_{j=1}^{15} A_j\right) + A_{18} + A_{19}. \end{aligned}$$


By using the linear system  $|D_2|$ , we obtain the following.

**Proposition 15.3.** *The K3 surface is the double cover of  $\mathbb{P}^2$  branched over a sextic curve with an  $a_{13}$  singularity. The image of  $A_4$  and  $A_{13}$  is a line, so are the images of  $A_{16}$  and  $A_{17}$  and of  $A_{18}$  and  $A_{19}$ .*

The automorphism group of the general K3 surface is  $(\mathbb{Z}/2\mathbb{Z})^2$  (see [Kon89, Section 4]).

### 15.3. The lattice $U \oplus E_8 \oplus E_7$

The K3 surface  $X$  contains 19  $(-2)$ -curves  $A_1, \dots, A_{19}$ . The dual graph of these curves is (see [Kon89, Section 4])



The curves  $A_1, \dots, A_{17}$  generate the Néron–Severi lattice; in that basis, the divisor

$$D_{538} = (32, 66, 46, 27, 9, 55, 45, 36, 28, 21, 15, 10, 6, 3, 1, 0, 1)$$

is ample, of square 538, with  $D_{538} \cdot A_j = 1$  for  $j \neq 1, 4, 17$  and  $D_{538} \cdot A_j = 2, 11, 16$  for  $j = 1, 4, 17$ .

The automorphism group of the general K3 surface is  $(\mathbb{Z}/2\mathbb{Z})^2$  (see [Kon89, Section 4]).


The coarse moduli space  $\mathcal{M}^N$  of  $N = U \oplus E_8 \oplus E_7$ -polarized K3 surfaces is studied in [CD12], where it is proved that this moduli space can be seen naturally as the open set

$$\mathcal{M}^N = \{[\alpha, \beta, \gamma, \delta] \in \mathbb{W}\mathbb{P}^3(2, 3, 5, 6) \mid \gamma \neq 0 \text{ or } \delta \neq 0\}$$

of a weighted projective space; thus in particular that moduli is rational. In [CD12] is also given a (singular) model of the K3 surfaces in  $\mathcal{M}^N$  as a quartic surface in  $\mathbb{P}^3$ . The K3 surfaces with lattice  $U \oplus E_8 \oplus E_7$  also belong among the “famous 95” families discussed in Section 2.6.

**15.4. The lattice  $U \oplus E_8 \oplus E_8$**

The set of  $(-2)$ -curves  $A_1, \dots, A_{19}$  on the K3 surface  $X$  and their configuration had been determined in the classical work of Vinberg [Vin75] (see also [Kon89, Section 4]):



The curves  $A_1, \dots, A_{18}$  generate the Néron–Severi lattice; in that base, the divisor

$$D_{620} = (-46, -91, -135, -68, -110, -84, -57, -29, 30, 61, 93, 126, 160, 195, 231, 115, 153, 76)$$

is ample, of square 620, with  $D_{620} \cdot A_j = 1$  for  $1 \leq j \leq 19$ . The divisor

$$D_2 = (-5, -10, -15, -8, -12, -9, -6, -3, 3, 6, 9, 12, 15, 18, 21, 10, 14, 7)$$

is nef, of square 2, with  $D_2 \cdot A_4 = D_2 \cdot A_{16} = 1$  and  $D_2 \cdot A_j = 0$  for  $j \neq 4, 16$ . Using  $|D_2|$ , we see that the K3 surface  $X$  is the double cover of  $\mathbb{P}^2$  branched over a sextic curve with an  $\mathfrak{a}_{17}$  singularity. We have

$$D_2 \equiv A_1 + 2A_2 + 3A_3 + A_4 + 3\left(\sum_{j=5}^{15} A_j\right) + A_{16} + 2A_{17} + A_{18} + 3A_{19};$$


thus the image of  $A_4, A_{16}$  by the double cover map is a line through the singularity.

The automorphism group of the general K3 surface is  $(\mathbb{Z}/2\mathbb{Z})^2$  (see [Kon89, Section 4]).

The rich geometry of  $U \oplus E_8 \oplus E_8$ -polarized K3 surfaces is studied in [CD07]. These K3 surfaces also belong among the “famous 95” families discussed in Section 2.6.

**15.5. The lattice  $U \oplus E_8 \oplus E_8 \oplus A_1$**

The K3 surface  $X$  contains 24  $(-2)$ -curves  $A_1, \dots, A_{24}$ . The dual graph of these curves is (see [Kon89, Section 4])



Here the dotted thick segments indicate an intersection number equal to 6 between the curves. The automorphism group of the general K3 surface is  $\mathfrak{S}_3 \times \mathbb{Z}/2\mathbb{Z}$  (see [Kon89, Section 4]).

16. Table: Number of  $(-2)$ -curves

Lattice	$\#(-2)$		Aut		Lattice	$\#(-2)$		Aut	
<b>Rank 3</b>					<b>Rank 3</b>				
$S_1$	6	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{1,3,1}$	3	<b>u</b>	$\{1\}$	
$S_2$	6	<b>u</b>	$\{1\}$		$S_{1,4,1}$	4	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$S_3$	4	<b>u</b>	$\{1\}$		$S_{1,5,1}$	6	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$S_4$	4		$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{1,6,1}$	4	<b>u</b>	$\{1\}$	
$S_5$	4	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{1,9,1}$	9		$\{1\}$	
$S_6$	6	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{4,1,1}$	3	<b>u</b>	$\{1\}$	
$S_{1,1,1}$	3	$\aleph \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$S_{5,1,1}$	4	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$S_{1,1,2}$	3	$\aleph \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$S_{6,1,1}$	4	<b>u</b>	$\{1\}$	
$S_{1,1,3}$	4	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{7,1,1}$	6		$\{1\}$	
$S_{1,1,4}$	4	<b>u</b>	$\{1\}$		$S_{8,1,1}$	4	<b>u</b>	$\{1\}$	
$S_{1,1,6}$	6		$\mathbb{Z}/2\mathbb{Z}$	$n$	$S_{10,1,1}$	8		$\mathbb{Z}/2\mathbb{Z}$	$n$
$S_{1,1,8}$	8		$\{1\}$		$S_{12,1,1}$	6		$\{1\}$	
$S_{1,2,1}$	3	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$S'_{4,1,2}$	4		$\mathbb{Z}/2\mathbb{Z}$	$n$
<b>Rank 4</b>					<b>Rank 5</b>				
$L(24)$	6		$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_1^{\oplus 3}$	7		$\mathbb{Z}/2\mathbb{Z}$	$t$
$L(27)$	8		$\mathbb{Z}/2\mathbb{Z}$	$n$	$U(2) \oplus \mathbf{A}_1^{\oplus 3}$	10		$\mathbb{Z}/2\mathbb{Z}$	$t$
$[8] \oplus \mathbf{A}_1^{\oplus 3}$	12	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U(4) \oplus \mathbf{A}_1^{\oplus 3}$	24		$\mathbb{Z}/2\mathbb{Z}$	$n$
$U \oplus \mathbf{A}_1^{\oplus 2}$	5	$\angle \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2$	6	$\angle \star$	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(2) \oplus \mathbf{A}_1^{\oplus 2}$	6	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{A}_3$	5	$\angle \star$	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(3) \oplus \mathbf{A}_1^{\oplus 2}$	8	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$[4] \oplus \mathbf{D}_4$	5		$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(4) \oplus \mathbf{A}_1^{\oplus 2}$	8	<b>u</b>	$\{1\}$		$[8] \oplus \mathbf{D}_4$	7		$\mathbb{Z}/2\mathbb{Z}$	$n$
$U \oplus \mathbf{A}_2$	4	$\aleph \star \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$[16] \oplus \mathbf{D}_4$	8		$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(2) \oplus \mathbf{A}_2$	4	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$[6] \oplus \mathbf{A}_2^{\oplus 2}$	10	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(3) \oplus \mathbf{A}_2$	4	$\angle \mathbf{u}$	$\{1\}$						
$U(6) \oplus \mathbf{A}_2$	6	<b>u</b>	$\{1\}$						
$L_{12} = S_0 \oplus \mathbf{A}_2$	6	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$					
$[4] \oplus [-4] \oplus \mathbf{A}_2$	6		$\mathbb{Z}/2\mathbb{Z}$	$n$					
$[4] \oplus \mathbf{A}_3$	5	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$					

Lattice	#(-2)		Aut		Lattice	#(-2)		Aut	
<b>Rank 6</b>					<b>Rank 9</b>				
$U(3) \oplus \mathbf{A}_2^{\oplus 2}$	12	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{E}_7$	9	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U(4) \oplus \mathbf{D}_4$	9		$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_6 \oplus \mathbf{A}_1$	10		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_4$	6	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 3}$	15		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_1 \oplus \mathbf{A}_3$	7	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_1^{\oplus 7}$	37	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_2^{\oplus 2}$	7	$\angle \mathbf{★}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U(2) \oplus \mathbf{A}_1^{\oplus 7}$	240	<b>u</b>	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_2$	8	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_7$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(2) \oplus \mathbf{A}_1^{\oplus 4}$	16	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_3$	10	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U \oplus \mathbf{A}_1^{\oplus 4}$	9		$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_5 \oplus \mathbf{A}_2$	10	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U(2) \oplus \mathbf{D}_4$	6	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_7$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{D}_4$	6	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{A}_1 \oplus \mathbf{E}_6$	10	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$
<b>Rank 7</b>					<b>Rank 10</b>				
$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1$	8		$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{E}_8$	10	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_1^{\oplus 5}$	12	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_8$	10		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U(2) \oplus \mathbf{A}_1^{\oplus 5}$	27	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{E}_7 \oplus \mathbf{A}_1$	11	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_1 \oplus \mathbf{A}_2^{\oplus 2}$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_4^{\oplus 2}$	11	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_1^{\oplus 2} \oplus \mathbf{A}_3$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_6 \oplus \mathbf{A}_1^{\oplus 2}$	14		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_2 \oplus \mathbf{A}_3$	8	$\angle \mathbf{★}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U(2) \oplus \mathbf{D}_4^{\oplus 2}$	18	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus \mathbf{A}_1 \oplus \mathbf{A}_4$	8	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 4}$	27	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_5$	7	$\angle \mathbf{★}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_1^{\oplus 8}$	145		$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus \mathbf{D}_5$	7	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_2 \oplus \mathbf{E}_6$	11	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$
<b>Rank 8</b>					<b>Rank 11</b>				
$U \oplus \mathbf{D}_6$	8		$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{E}_8 \oplus \mathbf{A}_1$	12	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 2}$	10		$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1$	14		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_1^{\oplus 6}$	19	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1$	22		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U(2) \oplus \mathbf{A}_1^{\oplus 6}$	56	<b>u</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{D}_4 \oplus \mathbf{A}_1^{\oplus 5}$	90		$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus \mathbf{A}_2^{\oplus 3}$	10	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$					
$U \oplus \mathbf{A}_3^{\oplus 2}$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$					
$U \oplus \mathbf{A}_2 \oplus \mathbf{A}_4$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	<b>Rank 12</b>				
$U \oplus \mathbf{A}_1 \oplus \mathbf{A}_5$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus \mathbf{E}_8 \oplus \mathbf{A}_1^{\oplus 2}$	14		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_6$	8	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_8 \oplus \mathbf{A}_1^{\oplus 2}$	19		$\mathbb{Z}/2\mathbb{Z}$	$t$
$U \oplus \mathbf{A}_2 \oplus \mathbf{D}_4$	9	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{D}_4^{\oplus 2} \oplus \mathbf{A}_1^{\oplus 2}$	59		$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus \mathbf{A}_1 \oplus \mathbf{D}_5$	9	<b>★</b>	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus \mathbf{A}_2 \oplus \mathbf{E}_8$	13	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$
$U \oplus \mathbf{E}_6$	8	$\mathfrak{N} \mathbf{★} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$n$					

Lattice	#(-2)		Aut		Lattice	#(-2)		Aut	
<b>Rank 13</b>					<b>Rank <math>\geq 15</math></b>				
$U \oplus E_8 \oplus A_1^{\oplus 3}$	17	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$	$U \oplus E_8 \oplus D_4 \oplus A_1$	21		$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus D_8 \oplus A_1^{\oplus 3}$	39	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$	$U \oplus E_8 \oplus D_6$	19	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
$U \oplus E_8 \oplus A_3$	14	$\star$	$\mathbb{Z}/2\mathbb{Z}$	$n$	$U \oplus E_8 \oplus E_7$	19	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
					$U \oplus E_8 \oplus E_8$	19	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$
<b>Rank 14</b>					$U \oplus E_8 \oplus E_8 \oplus A_1$	24	$\mathfrak{N} \mathbf{u}$	$\mathfrak{S}_3 \times \mathbb{Z}/2\mathbb{Z}$	$tn$
$U \oplus E_8 \oplus D_4$	15	$\mathfrak{N} \mathbf{u}$	$\mathbb{Z}/2\mathbb{Z}$	$t$					
$U \oplus D_8 \oplus D_4$	20	$\mathfrak{N} \mathbf{u}$	$(\mathbb{Z}/2\mathbb{Z})^2$	$tn$					
$U \oplus E_8 \oplus A_1^{\oplus 4}$	27		$(\mathbb{Z}/2\mathbb{Z})^2$	$t$					

This table give the number of  $(-2)$ -curves. A  $\mathfrak{N}$  means that the lattice is among the 95 famous families; a  $\angle$  means that the lattice is a mirror of one of the 95 famous ones, but not one of the 95 (see Section 2.6). A  $\star$  means that their  $(-2)$ -curve configuration is predicted by Theorem 2.6. A  $\mathbf{u}$  means that the unirationality of the moduli space is proved (the absence of  $\mathbf{u}$  does not exclude the possibility of unirationality). The column Aut gives the automorphism group of the general K3 surface. A  $t$  means that the action on the Néron–Severi lattice of a hyperelliptic involution is trivial, an  $n$  means that the action of a hyperelliptic involution is not trivial, and a  $tn$  means that both cases exists. From that data, one can recover the kernel of the map  $\text{Aut}(X) \rightarrow O(\text{NS}(X))$ .

## References

- [AN06] V. Alexeev and V. V. Nikulin, *Del Pezzo and K3 surfaces*, MSJ Mem. 15, Math. Soc. Japan, Tokyo, 2006.
- [ACL21] M. Artebani, C. Correa Diesler and A. Laface, *Cox rings of K3 surfaces of Picard number three*, J. Algebra **565** (2021), 598–626
- [ACR20] M. Artebani, C. Correa Deisler and X. Roulleau, *Mori dream K3 surfaces of Picard number four: projective models and Cox rings*, Israel J. of Math. (to appear), [arXiv:2011.00475](https://arxiv.org/abs/2011.00475).
- [AHL10] M. Artebani, J. Hausen and A. Laface, *On Cox rings of K3 surfaces*, Compos. Math. **146** (2010), no. 4, 964–998.
- [AK11] M. Artebani and S. Kondo, *The moduli of curves of genus six and K3 surfaces*, Trans. Amer. Math. Soc. **363** (2011), no. 3, 1445–1462.
- [BHP<sup>+</sup>04] W. Barth, K. Hulek, C. Peters and A. Van De Ven, *Complex Compact Surfaces*, 2nd ed., *Ergebn. Math Grenzgeb. (2)*, Springer-Verlag, Berlin, 2004.
- [Bel02] S.-M. Belcastro, *Picard lattices of families of K3 surfaces*, Comm. Algebra **30** (2002), no. 1, 61–82.
- [BCP97] W. Bosma, J. Cannon and C. Playoust, *The Magma algebra system. I. The user language*, In: *Comput. alg. and num. th.* (London, 1993), J. Symbolic Comput. **24**, 1997, no. 3–4, 235–265.
- [CD07] A. Clingher and C. Doran, *Modular invariants for lattice polarized K3 surfaces*, Michigan Math. J. **55** (2007), no. 2, 355–393.
- [CD12] ———, *Lattice Polarized K3 Surfaces and Siegel Modular Forms*, Adv. Math. **231** (2012), no. 1, 172–212.

- [Cox50] H. S. M. Coxeter, *Self-dual configurations and regular graphs*, Bull. Amer. Math. Soc. **56** (1950), 413–455.
- [Deg19] A. Degtyarev, *Tritangents to smooth sextic curves*, Ann. Inst. Fourier (to appear), [arXiv:1909.05657](https://arxiv.org/abs/1909.05657).
- [Dem76] M. Demazure, *Surfaces de Del Pezzo : IV - Systèmes anticanoniques*, Séminaire sur les singularités des surfaces (Polytechnique) (1976-1977), exp. no. 6, p. 1–11. Available from [http://www.numdam.org/item/?id=SSS\\_1976-1977\\_\\_\\_\\_A7\\_0](http://www.numdam.org/item/?id=SSS_1976-1977____A7_0).
- [Dol96] I. Dolgachev, *Mirror symmetry for lattice polarized K3 surfaces*, In: *Algebraic geometry, 4*, J. Math. Sci. **81** (1996), no. 3, 2599–2630.
- [DS20] I. Dolgachev and I. Shimada, *15-nodal quartic surfaces. Part II: The automorphism group*, Rend. Circ. Mat. Palermo (2) **69** (2020), 1165–1191.
- [Elk09] N. Elkies, *About the cover: Rational curves on a K3 surface*, In: *Arithmetic geometry, 1–4*, Clay Math. Proc. 8, Amer. Math. Soc., Providence, RI, 2009.
- [GLP10] F. Galluzzi, G. Lombardo and C. Peters, *Automorphisms of indefinite binary quadratic forms and K3-surfaces with Picard number 2*, Rend. Semin. Mat. Univ. Politec. Torino **68** (2010), no. 1, 57–77.
- [Har92] J. Harris, *Algebraic Geometry, A First Course*, Grad. Texts in Math. 133, Springer-Verlag, New York, 1992.
- [Har77] R. Hartshorne, *Algebraic geometry*, Grad. Texts in Math. 52, Springer-Verlag, New York–Heidelberg, 1977.
- [Huy16] D. Huybrechts, *Lectures on K3 surfaces*, Cambridge Stud. Adv. Math. 158, Cambridge Univ. Press, Cambridge, 2016.
- [KM12] M. Kobayashi and M. Mase, *Isomorphism among families of weighted K3 hypersurfaces*, Tokyo J. Math. **35** (2012), no. 2, 461–467.
- [Kon86] S. Kondo, *On automorphisms of algebraic K3 surfaces which act trivially on Picard groups*, Proc. Japan Acad., Ser. A **62**, 356–359 (1986).
- [Kon89] ———, *Algebraic K3 surfaces with finite automorphism groups*, Nagoya Math. J. **116** (1989), 1–15.
- [Kon00] ———, *A complex hyperbolic structure for the moduli space of curves of genus three*, J. reine angew. Math. **525** (2000), 219–232.
- [Kon02] ———, *The Moduli Space of Curves of Genus 4 and Deligne-Mostow’s Complex Reflection Groups*, In: *Algebraic Geometry 2000* (Azumino, 2000), pp. 383–400, Adv. Stud. Pure Math. 36, Math. Soc. Japan, Tokyo, 2002.
- [Kov94] S.J. Kovács, *The cone of curves on a K3 surface*, Math. Annalen **300**, (1994), no. 4, 681–692.
- [vLu07] R. van Luijk, *K3 surfaces with Picard number one and infinitely many rational points*, Algebra Number Theory **1** (2007), no. 1, 1–15.
- [McK10] J. McKernan, *Mori dream spaces*, Japan. J. Math. **5** (2010), no. 1, 127–151.
- [Mor84] D. R. Morrison, *On K3 surfaces with large Picard number*, Invent. Math. **75** (1984), no. 1, 105–121.
- [Muk88a] S. Mukai, *Curves, K3 Surfaces and Fano 3-folds of Genus  $\leq 10$* , In: *Algebraic geometry and commutative algebra, Vol. I*, pp. 357–377, Kinokuniya, Tokyo, 1988.
- [Muk88b] ———, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, Invent. Math. **94** (1988), no. 1, 183–221.

- [Muk92] ———, *Polarized K3 surfaces of genus 18 and 20*, In: *Complex projective geometry*, pp. 264–276, London Math. Soc. Lecture Note Ser. 179, Cambridge Univ. Press, Cambridge, 1992.
- [Nik79] V. V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, *Izv. Akad. Nauk SSSR Ser. Mat.* **43** (1979), no. 1, 111–177.
- [Nik80] ———, *Finite groups of automorphisms of Kähler K3 surfaces*, *Tr. Mosk. Mat. Obs.* **38** (1980), 71–135.
- [Nik83] ———, *Factor groups of automorphisms of hyperbolic forms with respect to subgroup generated by 2-reflections, algebro-geometric applications*, *J. Soviet Math.* **22** (1983), 1401–1475.
- [Nik85] ———, *Surfaces of type K3 with a finite automorphism group and a Picard group of rank three*, *Proc. Steklov Inst. Math.* **3** (1985), 131–155.
- [Nik14] ———, *Elliptic fibrations on K3 surfaces*, *Proc. Edinb. Math. Soc. (2)* **57** (2014), no. 1, 253–267.
- [Ott13] J. C. Ottem, *Cox rings of K3 surfaces with Picard number 2*, *J. Pure Appl. Algebra* **217** (2013), no. 4, 709–715.
- [PS71] I. I. Pjateckii-Sapiro and I. R. Shafarevich, *Torelli's theorem for algebraic surfaces of type K3* (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 530–572.
- [Rat06] J. Ratcliffe, *Foundations of Hyperbolic Manifolds*, Grad. Texts Math. 149, Springer, New York, 2006.
- [Rou19] X. Roulleau, *On the geometry of K3 surfaces with finite automorphism group and no elliptic fibrations*, *Internat. J. Math.* **33** (2022), no. 6, Paper No. 2250040.
- [Shi10] I. Shimada, *Lattice Zariski  $k$ -ples of plane sextic curves and  $Z$ -splitting curves for double plane sextics*, *Michigan Math. J.* **59** (2010), 621–665.
- [Shi14] ———, *Projective models of the supersingular K3 surface with Artin invariant 1 in characteristic 5*, *J. Algebra* **403** (2014), 273–299.
- [Shi15] ———, *An algorithm to compute automorphism groups of K3 surfaces and an application to singular K3 surfaces*, *Int. Math. Res. Not. IMRN* (2015), no. 22, 11961–12014.
- [Sai74] B. Saint-Donat, *Projective models of K3 surfaces*, *Amer. J. Math.* **96** (1974), no. 4, 602–639.
- [Ste85] H. Sterk, *Finiteness results for algebraic K3 surfaces*, *Math. Z.* **189** (1985), no. 4, 507–513.
- [Vin75] E. B. Vinberg, *Some arithmetical discrete groups in Lobacevskii spaces*, In: *Discrete subgroups of Lie groups and applications to moduli* (Internat. Colloq., Bombay, 1973), pp. 323–348. Oxford Univ. Press, Bombay, 1975.
- [Vin07] ———, *Classification of 2-reflective hyperbolic lattices of rank 4*, *Trans. Moscow Math. Soc.* 2007, 39–66.
- [Yon90] T. Yonemura, *Hypersurface K3 Singularities*, *Tohoku Math J.* **42** (1990), 351–380.
- [Zha98] D.-Q. Zhang, *Quotients of K3 surfaces modulo involutions*, *Japan J. Math.* **24**, no. 2 (1998).