
Explicit equations of the Cartwright-Steger surface

Lev A. Borisov and Sai-Kee Yeung

Abstract. We construct explicit equations of Cartwright-Steger and related surfaces.

Keywords. Surfaces of general type; explicit equations; ball quotients

2010 Mathematics Subject Classification. 14J29; 14Q10

[Français]

Équations explicites de la surface de Cartwright-Steger

Résumé. Nous construisons des équations explicites pour la surface de Cartwright-Steger et pour d'autres surfaces similaires.

arXiv:1804.00737v6 [math.AG] 9 Jul 2020

Received by the Editors on August 1, 2019, and in final form on December 20, 2019.

Accepted on June 4, 2020.

Lev A. Borisov

Department of Mathematics, Rutgers University, Piscataway, NJ 08854

e-mail: borisov@math.rutgers.edu

Sai-Kee Yeung

Department of Mathematics, Purdue University, West Lafayette, IN 47906

e-mail: yeung@math.purdue.edu

L.B. was partially supported by the NSF grant DMS-1601907. S.-K. Y. was partially supported by the NSF grant DMS-1501282.

© by the author(s)

This work is licensed under <http://creativecommons.org/licenses/by-sa/4.0/>

Contents

1. Introduction	2
2. Constructing the surface $Z_3 = \widehat{\mathcal{B}^2/\text{Ker}(\rho_3)}$	3
3. On the quotient of CS surface by the cyclic group of order three.. . . .	8
4. Constructing 21-fold cover of CS surface	8
5. Constructing CS surface	11
6. Verification	12
References	13

1. Introduction

The Cartwright-Steger surface is a smooth projective algebraic surface of Euler number 3. It is a complex two ball quotient with first Betti number 2. It was discovered by Cartwright and Steger in their work [CS10] in the process of completing the scheme of classification of fake projective planes proposed in [PY07]. Basic properties of the Cartwright-Steger surface can be found in [CS10] and [CKY17].

Our goal in this paper is to give explicit defining equations for a Cartwright-Steger surface Z . This is motivated in part by the recent work of [BK18], where explicit equations were given for two fake projective planes.

There are several corollaries of theoretical interest to our main result. One corollary is that the Cartwright-Steger surface can be defined over \mathbb{R} and even over \mathbb{Q} . Hence the complex conjugate of the surface is biholomorphic to itself. This corrects a mistake in [Y17] which we explain in Remark 5.1.

To put our work in a more global perspective, it is well known that 3 is the smallest Euler number achievable by a smooth surface of general type and is achieved only if the surface is a smooth complex two ball quotient \mathcal{B}^2/Γ for some torsion free lattice Γ in $PU(2, 1)$, the automorphism group of the complex two ball \mathcal{B}^2 . The lattice has to be arithmetic following the work of [Y14], [Y17] or [EG18], where the recent paper [EG18] gives a general result about integrality of lattices and is applicable to our surface. The results of [PY07] and [CS10] then imply that such a surface is either a fake projective plane or a Cartwright-Steger surface. Together with the result in this paper, there are exactly 101 such surfaces. Cartwright-Steger surface is the only one with positive first Betti number.

Another consequence of this paper is that we show explicitly that the bicanonical divisor $2K_Z$ is very ample and the Cartwright-Steger surface Z is defined by a quadratic relation and cubic relations among global sections of $2K_Z$. This improves the result of [Y18] for Z and fills a gap there, see Remark 6.1.

In addition to explicit equations for Z , we have also obtained explicit equations and geometric information for a few related interesting surfaces described below.

Denote by Π the lattice associated to Z so that $Z = \mathcal{B}^2/\Pi$. Let $\bar{\Gamma}$ be the maximal arithmetic lattice in the commensurable class of Π as defined in [CKY17, § 1.1]. It was observed by Cartwright and Steger that Π can be written as the intersection

$$\Pi = \Pi_2 \cap \Pi_3$$

where

- Π_2 is an index 3 normal subgroup of $\bar{\Gamma}$;
- Π_3 contains the kernel of $\rho_3 : \bar{\Gamma} \rightarrow PGU(3, 3)$ and is the preimage of a subgroup G_{21} of $PGU(3, 3)$;

see [CKY17, § 1.1]. We consider the following surfaces.

- the Cartwright-Steger surface $Z = \mathcal{B}^2/\Pi$;
- the 21 : 1 cover of Z given by $Z_2 = \mathcal{B}^2/\text{Ker}(\rho_3) \cap \Pi_2$;
- the quotient of Z_2 by C_3 given by $\mathcal{B}^2/\text{Ker}(\rho_3)$ and its minimal resolution $Z_3 = \widehat{\mathcal{B}^2/\text{Ker}(\rho_3)}$;
- the quotient $Z/C_3 = \mathcal{B}^2/\Pi_3$ and its partial resolution $Z_1 = \widehat{Z/C_3}$.

These surfaces fit in the commutative diagram

$$(1.1) \quad \begin{array}{ccccc} Z_2 & \longrightarrow & Z_2/C_3 & \longleftarrow & Z_3 \\ \downarrow & & \downarrow & & \downarrow \\ Z & \longrightarrow & Z/C_3 & \longleftarrow & Z_1 \end{array}$$

where the vertical maps are quotients by a group of order 21, the maps \longrightarrow are quotients by a cyclic group C_3 , and the maps \longleftarrow are birational morphisms.

The structure of the paper is as follows. In Section 2 we consider the surface Z_3 which we describe explicitly by 35 quadratic equations in $\mathbb{C}P^{12}$. In Section 3 we use the results of Section 2 to construct explicit equations of the surface Z_1 . In Section 4 we describe a procedure that we used to *guess* equations of the surface Z_2 and specifically construct a number of points on it with high accuracy. While we do not prove that they are correct, we expect them to be. In Section 5 we describe the method we used to calculate the putative equations of Z . Finally, in Section 6 we explain the collection of computer-based checks that ensure that the equations we found indeed describe the Cartwright-Steger surface. The ancillary files for computations are available on arXiv:1804.00737.

Acknowledgements. L.B. thanks Igor Dolgachev, JongHae Keum and Carlos Rito for stimulating questions and useful comments. S.-K. Y. is grateful to Donald Cartwright for helpful discussions, and to Vincent Koziarz and Gopal Prasad for their interest. We used software packages GAP [GAP], Mathematica [Math16], Magma [Mag97] and Macaulay2 [Mac].

2. Constructing the surface $Z_3 = \widehat{\mathcal{B}^2/\text{Ker}(\rho_3)}$

In this section we describe the equations of the surface Z_3 in (1.1), which is the key starting point of our method.

Let $\rho_3 : \bar{\Gamma} \rightarrow PGU(3, 3)$ be the reduction homomorphism modulo three, described in [CKY17]. The action of $\Gamma_3 = \text{Ker}(\rho_3)$ on \mathcal{B}^2 has fixed points, and the quotient has 63 singular points of type $(\frac{1}{3}, \frac{1}{3})$. The minimal resolution of singularities is a surface Z_3 with $K_{Z_3}^2 = 42$, $h^{1,0}(Z_3) = 0$ and $h^{2,0}(Z_3) = 13$. Importantly, Z_3 admits an effective action of the finite simple group $G = PGU(3, 3)$ of order 6048. The character table of G which is very important for our discussion can be readily obtained with GAP software package [GAP] and is presented in Table 1.¹

There is a unique up to conjugation noncommutative subgroup $G_{21} \subset G$ which normalizes a 7-Sylow subgroup of G . As we pointed out in (1.1), the quotient of Z_3 by this subgroup G_{21} is the blowup Z_1 of Z/C_3 in three singular points O_1, O_2 and O_3 of type $(\frac{1}{3}, \frac{1}{3})$. Observe that Z_1 has Gorenstein singularities and its canonical class has a global section. Specifically, in the notation of [CKY17] there is a (canonical) curve E_3 on Z which gives a curve $D_3 := E_3/C_3$ on Z_3/C_3 . From [CKY17], E_3 passes once through O_1 and O_3 , four

¹The table presents values of characters of irreducible representations of G at all conjugacy classes of G . Conjugacy classes are labeled by their order and a letter to distinguish different conjugacy classes of elements of the same order. Zero values are replaced by \cdot for ease of reading.

Table 1. Character table of $G = PGU(3, 3)$.

	1a	7a	7b	2a	3a	4a	4b	6a	12a	12b	8a	8b	4c	3b
χ_1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
χ_2	6	-1	-1	-2	-3	-2	-2	1	1	1	.	.	2	.
χ_3	7	.	.	-1	-2	3	3	2	.	.	-1	-1	-1	1
χ_4	7	.	.	3	-2	B	\bar{B}	.	E	\bar{E}	F	\bar{F}	1	1
χ_5	7	.	.	3	-2	\bar{B}	B	.	\bar{E}	E	\bar{F}	F	1	1
χ_6	14	.	.	-2	5	2	2	1	-1	-1	.	.	2	-1
χ_7	21	.	.	5	3	1	1	-1	1	1	-1	-1	1	.
χ_8	21	.	.	1	3	C	\bar{C}	1	F	\bar{F}	\bar{F}	F	-1	.
χ_9	21	.	.	1	3	\bar{C}	C	1	\bar{F}	F	F	\bar{F}	-1	.
χ_{10}	27	-1	-1	3	.	3	3	.	.	.	1	1	-1	.
χ_{11}	28	.	.	-4	1	D	\bar{D}	-1	\bar{F}	F	.	.	.	1
χ_{12}	28	.	.	-4	1	\bar{D}	D	-1	F	\bar{F}	.	.	.	1
χ_{13}	32	A	\bar{A}	.	-4	-1
χ_{14}	32	\bar{A}	A	.	-4	-1

$$(A, B, C, D, E, F) = \left(\frac{1 - i\sqrt{7}}{2}, -1 - 2i, -3 - 2i, -4i, -1 - i, -i \right).$$

times through O_2 and does not pass through any of the six other fixed points. Denote by \widehat{D}_3 the proper transform of D_3 and G_i for $i = 1, 2, 3$ the (-3) exceptional curve at O_i of $Z_1 \rightarrow Z/C_3$. It follows easily that $K_{Z_1} = \pi^*K_S - \frac{1}{3}\sum_{i=1}^3 G_i$ and $K_{Z_1} = \widehat{D}_3 + G_2$. Hence a global section of K_{Z_1} exists and is consisting of the proper preimage of D_3 plus one (-3) exceptional curve G_2 . This global section pulls back on Z_3 to give an element $h \in H^0(Z_3, K_{Z_3})$ which is G_{21} -invariant, but not G -invariant. We will denote the corresponding canonical divisor on Z_3 by H .

We will now be able to identify the action of G on $H^0(Z_3, K_{Z_3})$. By a slight abuse of terminology, we will denote by χ_k both the k -th irreducible representation of G and its character.

Proposition 2.1. *Up to a choice of square root of (-1) , the action of G on $H^0(Z_3, K_{Z_3})$ is given by the direct sum of irreducible representations $\chi_2 + \chi_4$. The equations on $\text{Sym}^2(\chi_2 + \chi_4)$ are given as follows. We have $\text{Sym}^2(\chi_2) = \chi_7$ and $\text{Sym}^2(\chi_4) = \chi_5 + \chi_7$. We also have $\chi_2\chi_4 = \chi_6 + \chi_{12}$. There is a 35-dimensional space of quadratic relations which is built from the 14-dimensional irreducible subspace χ_6 of $\chi_2\chi_4$ and the 21-dimensional subspace that lies diagonally in the direct sum of two copies of χ_7 .*

Proof. We know that the action of the order seven elements of G on Z_3 is fixed-point free. By the Holomorphic Lefschetz Formula (since $H^2(Z_3, K_{Z_3})$ is acted on trivially) we see that the trace of these elements on $H^0(Z_3, K_{Z_3})$ is (-1) . Together with $\dim H^0(Z_3, K_{Z_3}) = 13$ we see that the possible characters are

$$\chi_{H^0(Z_3, K_{Z_3})} \in \{\chi_1 + 2\chi_2, \chi_2 + \chi_3, \chi_2 + \chi_4, \chi_2 + \chi_5\}.$$

Case 1. $\chi_{H^0(Z_3, K_{Z_3})} = \chi_1 + 2\chi_2$. This is impossible, because then the G_{21} -invariant canonical divisor H would be G -invariant.

Case 2. $\chi_{H^0(Z_3, K_{Z_3})} = \chi_2 + \chi_3$. We have used GAP to readily calculate

$$\text{Sym}^2(\chi_2 + \chi_3) = \text{Sym}^2(\chi_2) + (\chi_2\chi_3) + \text{Sym}^2(\chi_3) = \chi_7 + (\chi_8 + \chi_9) + (\chi_1 + \chi_{10}).$$

We know that the kernel of the map

$$\text{Sym}^2(H^0(Z_3, K_{Z_3})) \rightarrow H^0(Z_3, 2K_{Z_3})$$

between spaces of dimension 91 and 56 respectively is of dimension at least 35. However, χ_7 component of $\text{Sym}^2(H^0(Z_3, K_{Z_3}))$ cannot be in the kernel. Indeed, in that case, the quadratic equations on the six-dimensional subspace of $H^0(Z_3, K_{Z_3})$ would imply that they are zero. Similarly, χ_{10} component cannot go to zero, since 27-dimensional space of quadratic equations on 7 variables implies that they are zero. This means that both χ_8 and χ_9 components must vanish, which again leads to a contradiction, since this would mean that the product of two nonzero sections of $H^0(Z_3, K_{Z_3})$ (one from χ_2 and another from χ_3) would be zero.

Case 3. $\chi_{H^0(Z_3, K_{Z_3})} = \chi_2 + \chi_4$ or $\chi_{H^0(Z_3, K_{Z_3})} = \chi_2 + \chi_5$. We may assume the former case by either taking complex conjugate of alternatively applying an outer automorphism of G . We have

$$\text{Sym}^2(\chi_2 + \chi_4) = \text{Sym}^2(\chi_2) + (\chi_2\chi_4) + \text{Sym}^2(\chi_4) = \chi_7 + (\chi_6 + \chi_{12}) + (\chi_5 + \chi_7)$$

and consider the kernel of the map

$$\text{Sym}^2(H^0(Z_3, K_{Z_3})) \rightarrow H^0(Z_3, 2K_{Z_3}).$$

As in Case 2, we see that either the kernel of the map $\text{Sym}^2(H^0(Z_3, K_{Z_3})) \rightarrow H^0(Z_3, 2K_{Z_3})$ has no χ_7 parts or at most one part. By looking at the $\chi_2\chi_4 = \chi_6 + \chi_{12}$ component we see that χ_{12} cannot go to zero. Indeed, Magma [Mag97] calculations show that the scheme cut out by this 28-dimensional space of relations has the same Hilbert polynomial as the union of $\mathbf{x} = \mathbf{0}$ and $\mathbf{y} = \mathbf{0}$. This means that these relations have no nontrivial solutions. Together with the fact that dimension of the kernel is at least 35 we see that the kernel must contain one copy of χ_7 and χ_6 . If this copy of χ_7 is not diagonal, then Magma calculations show that the corresponding sections are zero; this means that χ_7 must lie diagonally. Finally, we see that if, in addition to the diagonal copy of χ_7 we have χ_5 in the kernel, then the scheme cut out by these equations is empty (we did these calculations modulo $(10 - i)$, which is sufficient). \square

We denote the variables which correspond to the basis of χ_2 and χ_4 by x_1, \dots, x_6 and y_1, \dots, y_6 respectively. The action of the two standard generators of the group G on these variables is given by Table 2. We used the data from the Atlas of Finite Group Representations [A].

Table 2. Action of G on $(x_1, \dots, x_6, y_1, \dots, y_7)$

(x_1, \dots, y_7)	$\mapsto (x_2, x_1, x_4, x_3, -x_5, (-2 + i)x_1 - (2 - i)x_2 + ix_3 + ix_4 - ix_6,$ $y_2, y_1, y_4, y_3, (1 + i)y_1 - (1 + i)y_2 - y_3 + y_4 + y_5, y_6,$ $iy_3 - iy_4 + y_7)$
(x_1, \dots, y_7)	$\mapsto (x_3, -ix_1 - x_3, x_5, x_6, -ix_1 - ix_2 - x_3, x_1 + (1 - i)x_2 - ix_3$ $-x_4 - x_5 + x_6, y_3, y_1, y_5, y_6, y_7, -iy_2 - iy_3 - y_4 - y_6 - iy_7,$ $-y_1 + y_2 + y_3 - y_5 + y_7)$

In what follows we will use a specific choice of $G_{21} \subset G$ whose generators act on x_i and y_j by the formulas of Table 3.

Remark 2.2. For the choice of G_{21} in Table 3, the G_{21} -invariant section of K_{Z_3} which gives H is readily computed to equal

$$7y_1 + y_2 + y_3 - (2 + 3i)y_4 - (2 + 3i)y_5 + y_6 + y_7$$

up to scaling.

Remark 2.3. While it is impractical to write down all equations in the kernel of

$$\text{Sym}^2(H^0(Z_3, K_{Z_3})) \rightarrow H^0(Z_3, 2K_{Z_3}),$$

we write two of them in Table 4 that generate the diagonal χ_7 and χ_6 respectively. The rest can be obtained by applying the group action. We note that one can scale the variables x and y separately in the diagonal equation, but we used a particular choice that ensured relatively simple formulas.

Table 3. Action of G_{21} on $(x_1, \dots, x_6, y_1, \dots, y_7)$

$$\begin{aligned}
(x_1, \dots, y_7) &\mapsto (-ix_1 - x_3, x_3, x_6, x_5, ix_1 + ix_2 + x_3, 2ix_1 - (1-i)x_2 + ix_3 \\
&\quad + x_4 + (1+i)x_5 - (1-i)x_6, y_1, y_3, y_6, y_5, (-1-i)y_1 + (1+i)y_3 \\
&\quad - y_5 + y_6 + y_7, -iy_2 - iy_3 - y_4 - y_6 - iy_7, -y_1 + y_2 + y_3 \\
&\quad - (1-i)y_5 - iy_6 + y_7) \\
(x_1, \dots, y_7) &\mapsto ((-1-i)x_1 - x_2 - (1-i)x_3 - ix_4 + x_5 - x_6, ix_1 + (1+i)x_2 \\
&\quad + x_3 - ix_5, -x_1 - ix_2 - ix_3 - (1-i)x_4 - 2x_5, (1+i)x_1 + x_2 \\
&\quad + x_3 + ix_6, -ix_2 - (1+i)x_3 + ix_4 - (1-i)x_5 - ix_6, 2x_1 + x_2 \\
&\quad + (1-i)x_3 + x_6, y_1, iy_1 - iy_2 - 2iy_3 - y_4 + (1+i)y_5 - y_6 \\
&\quad - (1+2i)y_7, -y_3 + iy_4 + iy_6 - (1-i)y_7, -y_1 + (1-i)y_2 - 2iy_3 \\
&\quad - y_4 + iy_5 - y_6 - iy_7, y_4, (1+i)y_1 - (1+i)y_2 - y_3 + y_4 + y_5, \\
&\quad (1+i)y_2 + (2+i)y_3 + (2-i)y_4 - y_5 + (1-2i)y_6 + (2+i)y_7)
\end{aligned}$$

Table 4. Equations of $Z_3 = \mathcal{B}^2/\widehat{\text{Ker}}(\rho_3)$

$$\begin{aligned}
&x_1^2 + 2(4y_1^2 + (2i)y_1y_2 - (2-2i)y_2^2 - 2y_3^2 + (3-5i)y_3y_4 + (1-2i)y_4^2 - (2-2i)y_3y_5 \\
&\quad + (3+3i)y_4y_5 - (2-2i)y_5^2 + (1+5i)y_3y_6 - 12y_4y_6 - (1+5i)y_5y_6 + (1+2i)y_6^2 \\
&\quad + (1+5i)y_3y_7 + 4y_4y_7 - (1-3i)y_5y_7 + (2-12i)y_6y_7 + (1+2i)y_7^2 + (1-i)y_1((1+i)y_3 \\
&\quad + (1-4i)y_4 + (1-3i)y_5 + y_6 + y_7) + (1-i)y_2((5+i)y_3 + 3y_4 + (3-i)y_5 - (6+7i)y_6 \\
&\quad + (2-3i)y_7)) = 0 \\
&(1-i)((-10-6i)x_3y_1 - (4+10i)x_4y_1 + (4-3i)x_5y_1 + (4+i)x_6y_1 + (5-9i)x_3y_2 \\
&\quad + (2-4i)x_4y_2 + x_5y_2 - (2+2i)x_6y_2 - (10+9i)x_3y_3 - (1+i)x_4y_3 + (1+6i)x_5y_3 \\
&\quad + (1+10i)x_6y_3 + (2-6i)x_3y_4 - (1-5i)x_4y_4 - 8x_5y_4 + (1+7i)x_6y_4 + (2+9i)x_3y_5 \\
&\quad - (4-2i)x_4y_5 + (1+3i)x_5y_5 - (5+2i)x_6y_5 - (10-3i)x_3y_6 + (2+2i)x_4y_6 + 4x_5y_6 \\
&\quad + (13-5i)x_6y_6 + x_1((3+2i)y_1 + (3+2i)y_2 - i((13+3i)y_3 + (1+3i)y_4 + y_5 \\
&\quad + (1-12i)y_6 - (2+3i)y_7)) + x_2((13-2i)y_1 + (1+7i)y_2 + (7+7i)y_3 - (2+2i)y_4 \\
&\quad - (5+2i)y_5 + (7-2i)y_6 - (2+5i)y_7) + (5+9i)x_3y_7 - (4-2i)x_4y_7 + (1-6i)x_5y_7 \\
&\quad - (2-i)x_6y_7) = 0
\end{aligned}$$

We will eventually prove that these 35 quadratic relations cut out a surface in $\mathbb{C}\mathbb{P}^{12}$ which is isomorphic to Z_3 . However, it is a fairly delicate argument, partly because some of the computer calculations that could have simplified it are too time consuming to perform with available hardware and software.

We start by getting a rough idea of what these equations cut out.

Proposition 2.4. *Let I be the ideal of $\mathbb{C}[x_1, \dots, y_7]$ generated by the dimension 35 space of relations from χ_7 and χ_6 , which are the G -translates of Table 4. Then the dimension of the degree k component of $\mathbb{C}[x_1, \dots, y_7]/I$ is equal to $21k(k-1) + 14$ for all $k \geq 2$. The scheme $S = \text{Proj } \mathbb{C}[x_1, \dots, y_7]/I$ is a disjoint union of smooth surfaces of total degree 42 and perhaps some curves and points.*

Proof. These results are the consequence of computations we performed with Magma [Mag97]. The Hilbert polynomial calculation was quickly performed with coefficients in $\mathbb{Q}[i]$. We can also show by Magma that these relations and the 10×10 minors of the Jacobian matrix of their partial derivatives have no common solutions. For this we picked some minors only, otherwise there are too many of them. We worked modulo prime number $(10-i)$ in $\mathbb{Z}[i]$, which is sufficient from semicontinuity.

The Hilbert polynomial computation implies that S has no irreducible components of dimension more than 2, and the sum of the degrees of surface components is 42. The calculation of minors implies that surface components are smooth and do not intersect any other components. \square

Remark 2.5. The scheme S has an additional involution given by

$$(x_1, \dots, x_6, y_1, \dots, y_7) \mapsto (-x_1, \dots, -x_6, y_1, \dots, y_7)$$

which commutes with the action of G .

Remark 2.6. A Mathematica [Math16] calculation shows that scheme $S \subset \mathbb{CP}^{12}$ contains a certain set of 126 lines. These form two orbits with respect to the action of G and one orbit with respect to the action of G together with the involution of Remark 2.5. One of these lines is given by

$$\begin{aligned} (x_1, \dots, y_7) = & (v, u, iv + (1-i)u, (-1-2i)v + 2iu, (2-i)v - 2u, \\ & (-2+2i)v + 3u, (2+2i)v - (1+3i)u, (-1-i)v + \frac{1}{2}(1+3i)u, \\ & \frac{1}{2}(-3-i)v + \frac{1}{2}(3+3i)u, \frac{1}{2}(1-i)v - u, 2v - (2+i)u, \\ & \frac{1}{2}(-3+i)v + \frac{1}{2}(3+i)u, \frac{1}{2}(-1-i)v + \frac{1}{2}(1+i)u) \end{aligned}$$

for $(u : v) \in \mathbb{CP}^1$. The rest can be calculated by applying the symmetries. We also observe that these lines intersect each other and form a connected set.

Proposition 2.7. *The scheme S has a unique dimension two component S_0 of degree 42.*

Proof. Magma calculation shows that the Hilbert polynomial of the subscheme of \mathbb{P}^{12} cut out by I and the equation of Remark 2.2 is $42(k-1)$. On the other hand, we can check that 42 out of 126 lines lie in $S \cap H$. This means that these lines are the intersections of H with two-dimensional components of S . Because these are a part of a connected set of lines, this means that they come from the same component S_0 which must then have degree 42. \square

Proposition 2.8. *The 35 quadratic equations from the kernel of*

$$\mathrm{Sym}^2(H^0(Z_3, K_{Z_3})) \rightarrow H^0(Z_3, 2K_{Z_3})$$

define an embedding of Z_3 into \mathbb{CP}^{12} .

Proof. The base locus of $H^0(Z_3, K_{Z_3})$ is contained in the intersection of all G -translates of the G -invariant canonical curve H . Since we know the description of H as a union of 21 preimages of E_3/C_3 and 21 exceptional curves, we see that this base locus is empty. Thus, $H^0(Z_3, K_{Z_3})$ gives a map $Z_3 \rightarrow \mathbb{P}^{12}$. Since $K_{Z_3}^2 = 42 > 0$, we see that the image of Z_3 must be a surface. We know that it is a subscheme of the scheme $S \subset \mathbb{P}^{12}$ which is cut out by 35 explicit equations we found in Proposition 2.1. By Proposition 2.7 we see that this image must be a smooth surface S_0 .

We claim that K_{Z_3} is ample. Indeed, base point freeness and $K_{Z_3}^2 > 0$ imply that Z_3 is a minimal surface of general type. Its map to the canonical model may only contract some trees of \mathbb{CP}^1 which intersect H and its translates trivially. Such curves would embed into Z_2/C_3 , with image disjoint from the ramification points of $\mathcal{B}^2 \rightarrow Z_2/C_3$. Since \mathbb{CP}^1 is simply connected, this embedding lifts to an embedding of \mathbb{CP}^1 into \mathcal{B}^2 , which is impossible.

Since the degree of S_0 is the same as $K_{Z_3}^2$, we see that the map $Z_3 \rightarrow S_0$ is birational. Together with the ampleness, we see that it is an isomorphism. We now see that the homogeneous coordinate ring of S_0 and the pluricanonical ring of Z_3 coincide for high enough degrees. Since the former is a quotient of the homogeneous coordinate ring of S which has the same Hilbert polynomial $21(k^2 - k) + 14$ as the pluricanonical ring, we see that $S_0 = S$, i.e. the 35 quadratic equations cut out $S = S_0 \cong Z_3$ scheme-theoretically. \square

Remark 2.9. While we believe that the embedding $Z_3 \rightarrow \mathbb{CP}^{12}$ is projectively normal and the 35 equations cut it out ideal-theoretically (rather than only up to saturation) we were unable to verify it, since the required computer calculations are very time-intensive.

Remark 2.10. It is important to understand, heuristically, what factors contributed to our ability to identify Z_3 based on the group representation data, with the goal of extending these ideas to construction of related surfaces, notably Galois covers of fake projective planes. Specifically, the fact that $[\bar{\Gamma} : \Gamma]$ is large meant that the group G was relatively big compared to the square of the canonical class, so group theoretic constraints

were more severe. The fact that $\mathcal{B}^2/\text{Ker}(\rho_3)$ had non-canonical quotient singularities led to a lower value of $K_{Z_3}^2$ than what would be predicted by $[\bar{\Gamma} : \text{Ker}(\rho_3)]$ alone, which lowered the degree of Z_3 and made lower degree equations more likely. On the other hand, $h^{0,1} > 0$ could in principle be beneficial, as a larger value of $h^{0,2}$ could lead to more equations of lower degree.

3. On the quotient of CS surface by the cyclic group of order three.

In this section we describe the quotient Z_1 of $Z_3 = \widehat{\mathcal{B}^2/\text{Ker}(\rho_3)}$ by a group of order 21, which is also the blowup of the quotient of CS surface Z by its automorphism group C_3 at three singular points of type $(\frac{1}{3}, \frac{1}{3})$. The procedure was to compute G_{21} -invariant polynomials in the homogeneous coordinates of \mathbb{P}^{12} and find the ring of invariants of $\mathbb{C}[x_0, \dots, y_7]/I$. While there is only one such invariant in degree 1, up to scaling, given by the equation of Remark 2.2, there is a dimension 4 space of invariants of degree 2 and a dimension 8 space of invariants of degree 3, and choices must be made carefully to get simpler equations.

We are not particularly interested in this surface, but its equations² are (at the moment) nicer than those of Z and we list them in Table 5. It is embedded into the weighted projective space $W\mathbb{P}(1, 2, 2, 2, 3, 3, 3, 3)$ with homogeneous coordinates $(W_0 : \dots : W_7)$. The variables W_2, \dots, W_5 are odd with respect to the involution of Remark 2.5, and the other four variables are even. It took a fair bit of work to find the appropriate basis of variables and of equations to have them of palatable length. Remarkably, these equations have rational coefficients, and it was a non-trivial exercise to find the ring generators³ for which this can be achieved. We have verified by Magma that these equations cut out a ring whose degree d part has dimension $(d^2 - d + 2)$ for large d , which is consistent with the expected dimension. Indeed, we have on Z_3

$$\dim H^0(Z_3, dK_{Z_3}) = 21d(d - 1) + 14,$$

and Holomorphic Lefschetz Formula allows us to find the dimensions of the subspaces of G_{21} -invariants.

4. Constructing 21-fold cover of CS surface

To construct the Cartwright-Steger surface, we needed to understand the surface $Z_2 = \mathcal{B}^2/(\text{Ker}(\rho_3) \cap \Pi_2)$. This surface has invariants

$$h^{0,1}(Z_2) = 7, \quad h^{0,2}(Z_2) = 27, \quad K_{Z_2}^2 = 189.$$

Its automorphism group contains $G \times C_3$ where the quotient by the cyclic group of order three gives $\mathcal{B}^2/\text{Ker}(\rho_3)$. This means, in particular, that we may assume that the C_3 -invariant part of $H^0(Z_2, K_{Z_2})$ is the pullback⁴ of $H^0(Z_3, K_{Z_3})$, whose character we will call $(\chi_2 + \chi_4) \otimes 1$.

The generator of C_3 has 63 fixed points of type $(\frac{1}{3}, \frac{1}{3})$. This allows us to calculate the dimensions of the C_3 -character subspaces of $H^i(Z_2, nK_{Z_2})$ by the Holomorphic Lefschetz Formula on holomorphic vector bundles, or Woods Hole Fixed Point Theorem, cf. [AB68]. Namely, the alternating sum of traces of the action of the generator of C_3 on $H^*(Z_2, \mathcal{O}_{Z_2})$ is given by $\frac{63}{(1-w)^2}$ where $w = e^{\frac{2}{3}\pi i}$. It equals $\frac{63}{(1-w^2)^2}$ for the square of the generator, so the alternating sums of the dimensions of the character components are $\frac{1}{3}\left(21 + \frac{63}{(1-w)^2} + \frac{63}{(1-w^2)^2}\right) = 14$, $\frac{1}{3}\left(21 + w^2 \frac{63}{(1-w)^2} + w \frac{63}{(1-w^2)^2}\right) = 14$ and $\frac{1}{3}\left(21 + w \frac{63}{(1-w)^2} + w^2 \frac{63}{(1-w^2)^2}\right) = -7$ for

²These are putative equations, since we found them by constructing multiple points on Z_3 with high precision and looked for relations on values of the invariant polynomials at these points. However, it should be straightforward to derive these equations from those of Z_3 .

³Actually, W_i only generate the ring at high enough degrees; we are missing a degree four generator.

⁴While there is no map $Z_2 \rightarrow Z_3$, a rational map suffices here.

Table 5. Equations of Z_1 .

$$\begin{aligned}
0 &= -2W_1^2 + W_2W_3 + W_0W_7, \\
0 &= 16W_0^5 + 152W_0^3W_1 + 21W_0W_1^2 + 2W_0W_2W_3 - W_0W_3^2 - 16W_2W_5 + 432W_0^2W_6 + 54W_1W_6, \\
0 &= 40W_0^3W_1 - 20W_0W_1^2 + 4W_0W_2^2 + 10W_0W_2W_3 - 108W_3W_4 + 432W_0^2W_6 + 216W_1W_6 - W_1W_7, \\
0 &= 8W_0^3W_2 + 68W_0W_1W_2 - 6W_0W_1W_3 - 432W_0^2W_4 - 216W_1W_4 - 512W_0^2W_5 - 64W_1W_5 \\
&\quad + 216W_2W_6 + 108W_3W_6 - W_2W_7, \\
0 &= -1280W_0^6 - 10624W_0^4W_1 + 7776W_0^2W_1^2 + 320W_1^3 - 80W_1W_2^2 + 80W_0^2W_3^2 - 48W_1W_3^2 \\
&\quad + 3456W_0W_2W_4 - 46656W_4^2 + 512W_0W_3W_5 - 4096W_5^2 - 34560W_0^3W_6 - 1728W_0W_1W_6 \\
&\quad + 128W_0^3W_7 + W_7^2, \\
0 &= 224W_0^4W_1 + 976W_0^2W_1^2 + 48W_1^3 - 18W_1W_2^2 - 2W_1W_3^2 + 1080W_0W_2W_4 - 11664W_4^2 \\
&\quad + 128W_0W_3W_5 - 3456W_0W_1W_6 - 8W_0^3W_7 + 17W_0W_1W_7 + 54W_6W_7, \\
0 &= 64W_0^6 + 640W_0^4W_1 + 209W_0^2W_1^2 + 8W_1^3 - W_0^2W_2^2 - 4W_1W_2^2 - 4W_0^2W_3^2 + 324W_0W_2W_4 \\
&\quad - 2916W_4^2 + 1728W_0^3W_6 - 108W_0W_1W_6 + 2916W_6^2 - 8W_0^3W_7 + 2W_0W_1W_7, \\
0 &= -160W_0^6 - 1520W_0^4W_1 - 334W_0^2W_1^2 - 8W_1^3 + W_1W_2^2 + 10W_0^2W_3^2 + W_1W_3^2 + 1728W_4W_5 \\
&\quad - 4320W_0^3W_6 - 1188W_0W_1W_6 + 20W_0^3W_7 + 4W_0W_1W_7, \\
0 &= -32W_0^4W_2 - 256W_0^2W_1W_2 - 34W_1^2W_2 + 2W_2^3 + 16W_0^4W_3 + 176W_0^2W_1W_3 + 28W_1^2W_3 - W_3^3 \\
&\quad + 1728W_0^3W_4 + 1080W_0W_1W_4 + 2048W_0^3W_5 + 512W_0W_1W_5 - 648W_0W_2W_6 - W_0W_2W_7 \\
&\quad - 2W_0W_3W_7 + 54W_4W_7 + 16W_5W_7, \\
0 &= 32W_0^4W_2 + 316W_0^2W_1W_2 - 32W_1^2W_2 + 2W_2^3 + 8W_0^2W_1W_3 + 20W_1^2W_3 - 432W_0W_1W_4 \\
&\quad - 192W_0W_1W_5 + 1080W_0W_2W_6 + 3456W_5W_6 - 9W_0W_2W_7 + 2W_0W_3W_7 + 54W_4W_7, \\
0 &= -16W_0^4W_2 - 142W_0^2W_1W_2 - 8W_1^2W_2 + W_2^3 - 8W_0^2W_1W_3 + 2W_1^2W_3 + 1836W_0W_1W_4 \\
&\quad - 256W_0^3W_5 - 128W_0W_1W_5 - 324W_0W_2W_6 + 5832W_4W_6
\end{aligned}$$

the characters 1, w and w^2 respectively. We know the dimensions of the character 1 components, because they equal those of Z_3 . This allows us to prove that

$$\chi_{H^1(Z_2, K_{Z_2})} = V_7 \otimes w^2, \quad \chi_{H^0(Z_2, K_{Z_2})} = (\chi_2 + \chi_4) \otimes 1 + V_{14} \otimes w$$

where V_7 and V_{14} are some representations of G of dimension 7 and 14 respectively, and w and w^2 denote the characters of C_3 . We also recall that there is a subgroup $G_{21} \subset G \times C_3$ of order 21 that acts freely on Z_2 such that the Cartwright-Steger surface Z is given by

$$Z = Z_2/G_{21}.$$

One can see that the only option is for G_{21} to map trivially to C_3 . Moreover, the structure of G ensures that the order 3 elements of G_{21} lie in the conjugacy class $3b$ of Table 1. The fact that these elements act with no fixed points allows us to determine that $V_{14} = \chi_6$ and V_7 is one of $\chi_1 + \chi_2, \chi_3, \chi_4, \chi_5$.⁵

Remark 4.1. As will be more apparent in a moment, it is difficult to exclude all possibilities in this situation. Therefore, our approach is to find the most plausible scenario, with the eventual goal of successfully verifying the final answer to have the correct numerical invariants.

Now that we know that

$$\chi_{H^0(Z_2, K_{Z_2})} = (\chi_2 + \chi_4) \otimes 1 + \chi_6 \otimes w,$$

we will denote the corresponding 27 variables by

$$(x_1 : \dots : x_6 : y_1 : \dots : y_7 : z_1 : \dots : z_{14})$$

in a way compatible with Section 2. The action of the generators of the group G on the z -variables is given in Table 6. The action of the generators of G_{21} is given in Table 7. We are using the same generators as in Tables 2 and 3.

⁵We believe that $\Lambda^2 H^0(Z_2, \Omega_{Z_2}) \rightarrow H^0(Z_2, K_{Z_2})$ is surjective, which leads to $V_7 = \chi_3$, but we will not use it in the paper.

Table 6. Action of G on (z_1, \dots, z_{14})

(z_1, \dots, z_{14})	$\mapsto (z_1, z_4 - z_5 + z_6, z_1 - z_3, z_2 - z_5 + z_6, z_1 - z_5, z_1 - z_6,$ $z_{12} - z_4 + z_5 - z_6, -z_{11} + z_{13}, -z_1 + z_{10} - z_2 + z_3 + z_5 + z_9,$ $-z_{10} + z_2 + z_4 - z_5 + z_6, -z_1 - z_{11}, z_2 + z_7, -z_1 - z_{11} + z_8,$ $z_{10} + z_{11} + z_{14} - z_2 + z_3)$
(z_1, \dots, z_{14})	$\mapsto (z_2 + z_3, -z_2 - z_7 + z_9, z_1 + z_2, -z_2 - z_5 - z_7 - z_8, -z_7,$ $z_2 + z_6, z_{10} + z_7, -z_{12}, z_{11} + z_7, z_{14} - z_2 - z_7, -z_{12} - z_2 + z_4,$ $-z_8, -z_{12} - z_2 + z_4 - z_5, z_{12} + z_{13} - z_4)$

Table 7. Action of G_{21} on (z_1, \dots, z_{14})

(z_1, \dots, z_{14})	$\mapsto (z_2 + z_3, -z_5 + z_6 - z_8, -z_1 + z_3, z_6 + z_9, z_2 + z_3 + z_7, z_3$ $-z_6, z_5 - z_6, -z_5, z_1 + z_{11} + z_{14} - z_3 - z_9, -z_{14} - z_5 + z_6 - z_8$ $+ z_9, z_{12} - z_3 - z_4, z_{10} - z_2 + z_9, -z_3 - z_4, z_1 + z_{13} + z_{14} - z_9)$
(z_1, \dots, z_{14})	$\mapsto (z_{11} + z_{14} - z_3 - z_4 + z_6 - z_9, -z_1 - z_{10} - z_{13} - z_{14} + z_4$ $+ z_5 - z_7 + z_9, z_{12} + z_{14} - z_3 - z_4 + z_5 + z_8 - z_9, -z_{10} - z_{13}$ $- z_{14} + z_2 + z_4 + z_9, z_{11} - z_{13} - z_3 + z_6, -z_3 - z_4 - z_9, z_1 + z_{10}$ $+ z_{13} - z_4 - z_5 - z_8, -z_1 - z_{11} + z_5 + z_8, z_{13} + z_3 - z_5 - z_8, -z_1$ $- z_{10} - z_{11} - z_{13} - z_{14} + z_4 + z_5 - z_7 + z_8 + z_9, -z_{11} + z_4 + z_8,$ $-z_{11} - z_6 + z_8, -z_{11} - z_{14} + z_4 + z_9, z_{11} + z_3 - z_5 - 2z_8)$

We now study the quadratic relations on (x_1, \dots, z_{14}) . The relations among (x_1, \dots, y_7) are well understood from Section 2. As before, the Holomorphic Lefschetz Formula and Kodaira Vanishing Theorem show that the dimension of w and w^2 eigenspaces of $H^0(Z_2, 2K_{Z_2})$ are 77 each. For example, we have

$$\frac{1}{3} \left(210 + w^2 \frac{63w}{(1-w)^2} + w \frac{63w^2}{(1-w^2)^2} \right) = 77.$$

We have

$$\text{Sym}^2(\chi_6) = \chi_1 + \chi_6 + \chi_7 + \chi_8 + \chi_9 + \chi_{10}$$

and we make a *guess* that the space of relations has the expected dimension $105 - 77 = 28$. This leads to $\chi_1 + \chi_{10}$ as relations. Magma calculation shows that the corresponding equations on $(z_1 : \dots : z_{14})$ cut out a dimension 5 subscheme in $\mathbb{C}\mathbb{P}^{13}$.

The next key idea comes from looking at the relations on $\left((\chi_2 + \chi_4) \otimes 1 \right) \otimes \left(\chi_6 \otimes w \right)$ which are linear combinations of $x_i z_j$ and $y_i z_j$. These are harder to guess but we only needed a partial guess. We have

$$\chi_2 \chi_6 = \chi_2 + \chi_4 + \chi_5 + \chi_{13} + \chi_{14}, \quad \chi_4 \chi_6 = \chi_2 + \chi_{11} + \chi_{13} + \chi_{14}$$

and we need to find a space of relations of dimension at least $(6+7)14 - 77 = 105$. It appeared reasonable to expect some kind of diagonal relation of the character type χ_2 . However, in contrast to Section 2 we can no longer pick this diagonal element randomly! We calculated to high precision several points of Z_3 (i.e. values of x_i and y_j) and then used Mathematica to find the diagonal relation of type χ_2 which was a linear combination of $x_i z_j$ and $y_i z_j$ such that the quadratic relations on z_k have a solution. There were two such diagonal relations, which differ by a sign. We get to pick one of them, which breaks the additional automorphism of Remark 2.5.

Note that given (x_1, \dots, y_7) , the above procedure determines (z_1, \dots, z_{14}) uniquely up to scaling. To get the desired (birational) triple cover we also impose the additional condition

$$\text{Cubic}_y = \text{Cubic}_z$$

where Cubic_y and Cubic_z are the G -invariant cubic equations on y_i and z_j respectively. They are unique up to scaling and we are free to make a choice. The two are proportional because they represent non-zero invariant sections of $3K$ on Z_3 ; that space can be seen to be one-dimensional by a character calculation.

As the result, we were able to construct, with very high precision, multiple points in \mathbb{CP}^{26} which come from Z_2 . We don't claim that Z_2 is embedded into \mathbb{CP}^{26} by the global sections of the canonical divisor, although we consider it likely.

Remark 4.2. We can then use these points to determine the quadratic relations on x_i, y_j, z_k . We have found the space of these relations to be of expected dimension 105. We expect that some degree three equations would also be needed to generate the homogeneous ideal of Z_2 . However, since our focus lies in constructing the CS surface, we are not particularly interested in Z_2 itself.

5. Constructing CS surface

To understand how the CS surface Z is constructed from Z_2 , recall that it is given as Z_2/G_{21} . The group G_{21} was found explicitly. Then we found G_{21} -invariant quadratic polynomials in the 27 variables x_i, y_j, z_k . These form a dimension ten subspace which splits according to the number of z_k into spaces of dimension 4, 3 and 3 for no z , one z and two z 's. This is precisely the decomposition of $H^0(Z, K_Z)$ into character subspaces of C_3 action.

We picked a basis U_0, \dots, U_9 and then we use the numerically constructed points of Z_2 to find the polynomial relations on U_i . Specifically, we solve for quadratic and then cubic equations in U_i which are numerically zero on the points of Z_2 . This gives approximate linear relations on the coefficients, which are then solved by Mathematica. Since all computations are performed with interval arithmetic, it is possible to calculate the ranks of the matrices in question correctly. At the end of the day, we are able to recognize the coefficients as algebraic numbers.

Remark 5.1. It is not at all a trivial matter to find a basis $\{U_i\}$ in which relations look nice. Remarkably, after making several informed choices, described below, we were able to find a basis in which the relations have *rational* coefficients! This means that, contrary to expectations, the Cartwright-Steger surface is unique, rather than a pair of surfaces that differ by complex conjugation. The expectation comes from the fact that for fake projective planes, complex conjugation gives rise to holomorphically different surfaces as proved in [KK02]. The paper [Y17] gives an attempt to prove such an expectation in page 1146, where an error was made stating that the oriented cover of $\mathbb{RP}^2 \# T^2$ was T^2 . Our results here disprove such an expectation.

Remark 5.2. We unexpectedly found a quadratic relation on U_i . This means that the map

$$\text{Sym}^2(H^0(Z, 2K_Z)) \rightarrow H^0(Z, 4K_Z)$$

of spaces of dimension 55 each has a non-trivial kernel. There is an 84-dimensional space of cubic relations given by the kernel of

$$\text{Sym}^3(H^0(Z, 2K_Z)) \rightarrow H^0(Z, 6K_Z),$$

as one would expect from the dimension count.

Remark 5.3. We explain how the choices were made. The action of C_3 on Z has 9 fixed points. We picked the basis elements U_7, U_8, U_9 so that the $(\frac{1}{3}, \frac{1}{3})$ fixed points are $(0 : \dots : 0 : 0 : 1)$, $(0 : \dots : 0 : 1 : 0)$, and $(0 : \dots : 1 : 0 : 0)$, which determines them uniquely up to scaling. The U_4, U_5, U_6 are determined by looking at the tangent spaces at these points. The sections U_0, U_1, U_2, U_3 are W_0^2, W_1, W_2, W_3 from Section 3, up to scaling. These form the invariant subspace of the C_3 action. The other two eigenspaces of the action are $\text{Span}(U_4, U_5, U_6)$ and $\text{Span}(U_7, U_8, U_9)$.

Remark 5.4. We have not been able to reduce the equations of CS surface Z to a form suitable for a printed version, since the coefficients are often about ten decimal digits long (which is better than the initial attempts that led to 40 digits long Gaussian integers). We make the equations available at [B].

6. Verification

In this section we comment on the collection of facts that we verified by computer calculations that imply that the surface we constructed is indeed the Cartwright-Steger surface. We have to be fairly creative, since many desirable computations take too long to finish.

We begin by observing that the scheme cut out by the 84 cubic equations is connected and smooth of dimension two. Connectedness follows from a Magma calculation that the quotient ring is an integral domain, which was done with rational coefficients. To prove smoothness, we calculated minors of Jacobian matrices of appropriately chosen subsets of the equation set (a finite field calculation suffices by semicontinuity). Specifically, we looked at the fixed points of C_3 action on Z and at each of them found 7 relations whose gradients are linearly independent at that point. Then we considered the size 7 minors of the Jacobian matrices of these sets of equations. Magma calculation shows that they have no common zeros on Z .

Thus we have a smooth surface Z ⁶ which is embedded into \mathbb{P}^9 by global sections of a divisor D . By Hilbert polynomial calculation and Riemann-Roch theorem, we see that

$$(6.1) \quad D^2 = 36, \quad K_Z D = 18, \quad \chi(\mathcal{O}_Z) = 1.$$

The equation $U_0 = 0$ gives a divisor $2C$ where C is the curve⁷ whose ideal in $\mathbb{C}\mathbb{P}^9$ is generated by U_0 and eighteen quadratic polynomials in Table 8 below.

Table 8. Equations of the $C \subset Z$

$ \begin{aligned} &U_0, -8U_1^2 - 2U_1U_2 + 2U_2^2 - 2U_1U_3 + 3U_4U_9 + 6U_6U_9, 16U_1^2 + 4U_2^2 - 4U_3^2 + 3U_6U_7, \\ &20U_1^2 + 12U_1U_2 + 8U_1U_3 + 2U_3^2 + U_5U_9, -4U_1^2 - 8U_1U_2 + 6U_2^2 - 4U_1U_3 + U_5U_8, \\ &4U_1^2 + 2U_1U_2 - 2U_2^2 - 2U_1U_3 + 2U_3^2 + 3U_4U_8, -2U_1^2 + U_2U_3, \\ &-6U_1U_4 - 6U_3U_4 - U_3U_5 + 12U_1U_6 - 12U_2U_6 + 6U_8U_9, \\ &-36U_1U_4 - 12U_3U_4 - 4U_1U_5 - 2U_2U_5 - 4U_3U_5 + 24U_1U_6 + 3U_7U_9, \\ &36U_1U_4 + 2U_2U_5 + 4U_3U_5 - 24U_1U_6 - 24U_2U_6 + 3U_7U_8, \\ &6U_1U_4 + 6U_3U_4 + U_3U_5 + 12U_3U_6, 6U_1U_4 + 3U_2U_4 + U_1U_5 + 12U_1U_6, \\ &18U_6^2 - 2U_1U_8 + 3U_2U_8 - 6U_1U_9 + 4U_2U_9 + 3U_3U_9, U_5U_6 + 2U_1U_8 + 2U_1U_9 + 2U_2U_9, \\ &U_5^2 + 12U_1U_7 + 3U_2U_7 + 6U_3U_7 - 16U_1U_8 + 48U_1U_9 - 4U_2U_9, \\ &9U_4U_6 + 4U_1U_8 + 6U_1U_9 - 5U_2U_9 - 3U_3U_9, \\ &3U_4U_5 - 3U_1U_7 - 3U_3U_7 - 8U_1U_8 - 36U_1U_9 + 4U_2U_9, \\ &18U_4^2 + 3U_3U_7 - 16U_1U_8 + 8U_2U_9 + 12U_3U_9, 2U_1U_8 + U_3U_8 + 2U_1U_9 - U_2U_9 \end{aligned} $
--

These equations were found using Mathematica. Once found, we can verify inclusion of ideals by Hilbert polynomial computations, which are readily performed in Magma with rational coefficients. We verified that C is an integral (but perhaps singular) curve by verifying that the quotient by its ideal is a domain.

We now consider the short exact sequence of sheaves

$$(6.2) \quad 0 \rightarrow \mathcal{O}_Z(-C)(1) \rightarrow \mathcal{O}_Z(1) \rightarrow (j_C)_* \mathcal{O}_C(1) \rightarrow 0.$$

We verified by Macaulay2 that $\dim H^1(Z, (j_C)_* \mathcal{O}_C(1)) = 1$ while working with rational coefficients. We have also verified that $H^1(Z, \mathcal{O}_Z(1)) = 0$ by a Macaulay2 calculation modulo 101 (calculations over rationals are more complicated and appear to be outside of the software and hardware capabilities),

⁶We use the same name as for the CS surface, but we are still to prove that Z is one.

⁷We expected that C is the zero locus E_3 of the section of the canonical divisor on Z . Our calculations below verify it.

which suffices by semicontinuity. The long exact sequence in cohomology coming from (6.2) then implies that $h^2(Z, \mathcal{O}(-C)(1)) > 0$. Since $2C = D$, this means $h^2(Z, \mathcal{O}(C)) > 0$, so $H^0(Z, \mathcal{O}(K_Z - C))$ is non-zero. From $2C = D$ and (6.1), we see that $CD = K_Z D = 18$. Since D is (very) ample, we see that $K_Z - C$ is a degree zero effective divisor. Therefore, we must have $K_Z - C = 0$. Then we get $K_Z^2 = 9$, $h^0(K_Z) \geq 1$, which means that Z is a Cartwright-Steger surface. We also see that $D = 2C = 2K_Z$, so the embedding is given by $2K_Z$.

Remark 6.1. It is stated that $2K_Z$ is very ample in [Y18], but the details in the elimination of a potential curve B satisfying Proposition 1-(i) in [Y18] are overlooked, as kindly informed by JongHae Keum to the second author. The current paper gives a stronger version of the statement.

References

- [A] *Atlas of finite group representations*, <http://brauer.maths.qmul.ac.uk/Atlas/v3/>.
- [AB68] M. F. Atiyah and R. Bott, *A Lefschetz fixed point formula for elliptic complexes. II. Applications.*, Ann. of Math. (2) **88** (1968), 451–491.
- [B] L. Borisov’s home page, <http://www.math.rutgers.edu/~borisov/CS/>.
- [BK18] L. Borisov and J. Keum, *Explicit equations of a fake projective plane*, preprint arXiv:1802.06333 (2018), <https://arxiv.org/abs/1802.06333>.
- [CKY17] D. Cartwright, V. Koziarz and S. K. Yeung *On the Cartwright-Steger surface*, J. Alg. Geo. **26** (2017), no. 4, 655–689.
- [CS10] D. Cartwright and T. Steger, *Enumeration of the 50 fake projective planes*, C. R. Acad. Sci. Paris, Ser. 1, **348** (2010), no. 1–2, 11–13.
- [EG18] H. Esnault and M. Groechenig, *Cohomologically rigid local system and integrality*, Selecta Math. (N.S.) **24** (2018), no. 5, 4279–4292.
- [GAP] GAP software package, <https://www.gap-system.org/>.
- [KK02] V. S. Kharlamov and V. M. Kulikov, *On real structures on rigid surfaces*, Izv. Russ. Akad. Nauk. Ser. Mat. **66** (2002), no. 1, 133–152; *translation in* Izv. Math. **66** (2002), no. 1, 133–150.
- [Mac] D. Grayson and M. Stillman, *Macaulay2, a software system for research in algebraic geometry*, <http://www.math.uiuc.edu/Macaulay2>.
- [Mag97] W. Bosma, J. Cannon and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265; v2.23-2.
- [Math16] *Mathematica 11.1*, Wolfram Research, Inc., 2016.
- [PY07] G. Prasad and S.-K. Yeung, *Fake projective planes*, Inv. Math. **168** (2007), no. 2, 321–370; Addendum, *ibid* **182** (2010), no. 1, 213–227.
- [Y14] S. K. Yeung, *Classification of surfaces of general type with Euler number 3*, J. Reine Angew. Math. **697** (2014), 1–14.
- [Y17] S. K. Yeung, *Foliations associated to harmonic maps and some complex two-ball quotients*, Sci. China Math. **60** (2017), no. 6, 1137–1148.
- [Y18] S. K. Yeung, *Very ampleness of the bicanonical line bundle on compact complex 2-ball quotients*, Forum Math. **30** (2018), no. 2, 419–432.