

Gushel-Mukai varieties: intermediate Jacobians

Olivier Debarre and Alexander Kuznetsov

To the memory of A.N. Tyurin

Abstract. Let X be a Gushel-Mukai variety of dimension 3 or 5. If $A \subset \bigwedge^3 V_6$ is the Lagrangian subspace associated with X, we prove that the intermediate Jacobian of X is isomorphic to the Albanese variety of the canonical double covering of any of the two dual Eisenbud-Popescu-Walter surfaces $Y_A^{\geq 2}$ and $Y_{A\perp}^{\geq 2}$. As an application, we describe the period maps for Gushel-Mukai varieties of dimension 3 or 5.

Keywords. Fano varieties; Gushel-Mukai varieties; intermediate Jacobians; EPW sextics; Albanese varieties; period maps; Abel-Jacobi maps; Grassmannians.

2020 Mathematics Subject Classification. 14J45; 14J30; 14J40; 14M15; 14K30

[Français]

Variétés de Gushel-Mukai : jacobiennes intermédiaires

Résumé. Soit X une variété de Gushel-Mukai de dimension 3 ou 5. Si $A \subset \bigwedge^3 V_6$ est le sous-espace lagrangien associé à X, nous démontrons que la jacobienne intermédiaire de X est isomorphe à la variété d'Albanese du revêtement double canonique de chacune des surfaces d'Eisenbud-Popescu-Walter $Y_A^{\geq 2}$ et $Y_{A^{\perp}}^{\geq 2}$. En guise d'application, nous décrivons les applications des périodes des variétés de Gushel-Mukai de dimension 3 ou 5.

Received by the Editors on May 14, 2020, and in final form on October 21, 2020. Accepted on November 5, 2020.

The Poncelet Laboratory, Independent University of Moscow

© by the author(s)

This work is licensed under http://creativecommons.org/licenses/by-sa/4.0/

Olivier Debarre

Université de Paris and Sorbonne Université, CNRS, Institut de Mathématiques de Jussieu-Paris Rive Gauche, F-75006 Paris, France *e-mail:* olivier.debarre@imj-prg.fr

Alexander Kuznetsov

Algebraic Geometry Section, Steklov Mathematical Institute, 8 Gubkin str., Moscow 119991, Russia

Laboratory of Algebraic Geometry, National Research University Higher School of Economics, Russian Federation *e-mail:* akuznet@mi-ras.ru

A.K. was partially supported by the HSE University Basic Research Program, Russian Academic Excellence Project "5-100".

Contents

1. Introduction	 •	•		•	•	•		•	•	3
2. Gushel-Mukai and Eisenbud-Popescu-Walter varieties	 •					•			•	7
2.1. Geometry of Gr(2,5)	 •	•	•	•	•	•			•	7
2.2. Eisenbud-Popescu-Walter varieties and their double coverings	 •								•	8
2.3. Gushel–Mukai varieties	 •								•	10
2.4. Linear spaces on quadrics containing GM varieties	 •								•	12
2.5. Linear spaces on ordinary GM threefolds and fivefolds \ldots .	 •	•				•			•	14
3. Topological preliminaries	 •									16
3.1. Abel-Jacobi maps	 •	•				•			•	16
3.2. Generalized blow up decomposition	 •	•				•			•	17
3.3. The Clemens-Tyurin argument	 •	•				•			•	19
3.4. Intermediate Jacobians and their endomorphisms	 •								•	20
4. Intermediate Jacobians of GM threefolds	 •		•							21
4.1. Family of curves	 •									21
4.2. The boundary of the family	 •									23
4.3. A relation between the subschemes	 •									27
4.4. Abel-Jacobi maps	 •		•		•				•	29
4.5. The line transform	 •		•		•				•	31
5. Intermediate Jacobians of GM fivefolds	 •									33
5.1. Family of surfaces	 •									34
5.2. The boundary of the family	 •									35
5.3. A relation between the subschemes	 •	•								37
5.4. Abel-Jacobi maps	 •	•								38
5.5. Simplicity argument	 •					•			•	39
6. Period maps	 •									42
References										
	 •	•	•	•	•	·	•	·	•	

1. Introduction

This article is an addition to the series [DK18, DK19, DK20b, KP18] on the geometry of Gushel-Mukai varieties. For an introduction, we recommend the survey [Deb20].

A smooth complex Gushel–Mukai (GM for short) variety of dimension $n \in \{2, 3, 4, 5, 6\}$ is a smooth dimensionally transverse intersection

(1.1)
$$X = \operatorname{CGr}(2, V_5) \cap \mathbf{P}(W) \cap Q,$$

where V_5 is a 5-dimensional complex vector space, $CGr(2, V_5) \subset \mathbf{P}(\mathbf{C} \oplus \bigwedge^2 V_5)$ is the cone over the Grassmannian $Gr(2, V_5)$ in its Plücker embedding, $W \subset \mathbf{C} \oplus \bigwedge^2 V_5$ is a linear subspace of dimension n + 5, and $Q \subset \mathbf{P}(W)$ is a quadratic hypersurface.

GM varieties of dimension 2 are Brill-Noether general K3 surfaces of genus 6. GM varieties of dimension 4 or 6 are Fano varieties but they share some properties with K3 surfaces. For instance, their derived categories have a component of K3-type ([KP18, Propositions 2.6 and 2.9]) and their vanishing cohomology of middle dimension is isomorphic to a Tate twist of the primitive second cohomology of a certain hyperkähler fourfold associated with their K3 category ([DK19, Theorem 5.1]). This allowed us to describe the period map for GM varieties of dimension 4 or 6 in [DK19, Proposition 5.27].

GM varieties of dimension 3 or 5 are also Fano varieties but they behave differently. The nontrivial components of their derived categories bear some of the features of the derived category of a curve ([KP18, Proposition 2.9]) and the Hodge structure on their middle cohomology defines their intermediate Jacobian, a 10-dimensional principally polarized abelian variety ([DK19, Proposition 3.1]). The main goal of this article is to describe the intermediate Jacobians and period maps of GM varieties of dimension 3 or 5.

The key object we use to study a GM variety X is its associated Lagrangian data set constructed in [DK18]. It is a triple $(V_6(X), V_5(X), A(X))$ (or (V_6, V_5, A) for short) that consists of a 6-dimensional vector space V_6 , a hyperplane $V_5 \subset V_6$, and a subspace $A \subset \bigwedge^3 V_6$ which is Lagrangian with respect to the symplectic form on $\bigwedge^3 V_6$ given by exterior product, and contains no decomposable vectors (this means no nonzero products $v_1 \land v_2 \land v_3$, where $v_1, v_2, v_3 \in V_6$). A GM variety can be reconstructed from its Lagrangian data set ([DK18, Theorem 3.6]). Moreover, Lagrangian data sets can be used to describe the moduli stack of smooth GM varieties and its coarse moduli space ([DK20b]).

It is not surprising then that many geometric properties of a GM variety can be described in terms of its Lagrangian data set, particularly in terms of the Eisenbud-Popescu-Walter (EPW for short) varieties

$$Y_A^{\geq 3} \subset Y_A^{\geq 2} \subset Y_A^{\geq 1} \subset Y_A^{\geq 0} = \mathbf{P}(V_6),$$

where $Y_A^{\geq 1}$ is a sextic hypersurface (called an EPW sextic) with singular locus $Y_A^{\geq 2}$, itself an integral surface with singular locus the finite set $Y_A^{\geq 3}$, and the dual EPW varieties

$$Y_{A^{\perp}}^{\geq 3} \subset Y_{A^{\perp}}^{\geq 2} \subset Y_{A^{\perp}}^{\geq 1} \subset Y_{A^{\perp}}^{\geq 0} = \mathbf{P}(V_6^{\vee})$$

associated with the Lagrangian subspace $A^{\perp} \subset \bigwedge^{3} V_{6}^{\vee}$ (see Section 2.2 for the definitions).

Let A be a Lagrangian with no decomposable vectors (such as A(X) and $A(X)^{\perp}$). O'Grady constructed a canonical double covering

$$\widetilde{Y}_A^{\geq 1} \longrightarrow Y_A^{\geq 1}$$
,

étale away from the surface $Y_A^{\geq 2}$. When $Y_A^{\geq 3}$ is empty (this holds for A general), $\widetilde{Y}_A^{\geq 1}$ is a hyperkähler fourfold called a double EPW sextic. When X is a GM variety of even dimension, the double EPW sextic $\widetilde{Y}_{A(X)}^{\geq 1}$ is the hyperkähler fourfold mentioned above whose primitive second cohomology is isomorphic to a Tate twist of the vanishing middle cohomology of X.

We also defined in [DK20a, Theorem 5.2(2)] a canonical double covering

$$\widetilde{Y}_A^{\geq 2} \longrightarrow Y_A^{\geq 2}$$
,

étale away from the finite set $Y_A^{\geq 3}$, where $\widetilde{Y}_A^{\geq 2}$ is an integral surface (called a double EPW surface) which has an ordinary double point over each point of $Y_A^{\geq 3}$ and is smooth elsewhere; in particular, $\widetilde{Y}_A^{\geq 2}$ is smooth for Ageneral. It has a 10-dimensional Albanese variety $Alb(\widetilde{Y}_A^{\geq 2})$ which, when $\widetilde{Y}_A^{\geq 2}$ is singular, can be defined as the Albanese variety of any desingularization.

The first main result of this article is the following.

Theorem 1.1. For any Lagrangian subspace $A \subset \bigwedge^3 V_6$ with no decomposable vectors, the Albanese variety $\operatorname{Alb}(\widetilde{Y}_A^{\geq 2})$ has a canonical principal polarization such that there is an isomorphism

(1.2)
$$\operatorname{Alb}(\widetilde{Y}_A^{\geq 2}) \simeq \operatorname{Alb}(\widetilde{Y}_{A^{\perp}})$$

of principally polarized abelian varieties.

If X is a smooth GM variety of dimension $n \in \{3, 5\}$, with intermediate Jacobian Jac(X) and associated Lagrangian subspace A, there is a canonical isomorphism

(1.3)
$$H_n(X, \mathbf{Z}) \simeq H_1(\widetilde{Y}_A^{\geq 2}, \mathbf{Z})$$

of polarized Hodge structures. It induces an isomorphism

(1.4)
$$\operatorname{Jac}(X) \simeq \operatorname{Alb}(Y_A^{\geq 2})$$

of principally polarized abelian varieties.

We prove this in Theorem 4.4 for GM threefolds X with $Y_{A(X)}^{\geq 3} = \emptyset$ and in Theorem 5.3 for GM fivefolds X with $Y_{A(X)}^{\geq 3} = \emptyset$. In particular, we use the natural principal polarization of the intermediate Jacobian Jac(X) to produce a principal polarization on $Alb(\widetilde{Y}_{A(X)}^{\geq 2})$ and we deduce the isomorphism (1.2) from a birational isomorphism (a line transform) between two GM threefolds X and X' such that $A(X') = A(X)^{\perp}$ (such pairs are called *period duals* in [DK18]). The extension to arbitrary GM threefolds and fivefolds is given in Section 6.

Remark 1.2. We are not aware of a direct proof of the isomorphism (1.2). It can be thought of as a Hodge-theoretic incarnation of the equivalence between the nontrivial components of derived categories of odd-dimensional GM varieties conjectured in [KP18, Conjecture 3.7] and proved in [KP19, Corollary 6.5]. It would be interesting to extract the principal polarization on $Alb(\tilde{Y}_A^{\geq 2})$ from the categorical data and to deduce the isomorphism (1.2) from the equivalence of categories.

One can also think of the principal polarization on Alb($\widetilde{Y}_{A}^{\geq 2}$) as of an element of

$$H^{2}(\mathrm{Alb}(\widetilde{Y}_{A}^{\geq 2}), \mathbf{Z}) = \bigwedge^{2} H^{1}(\mathrm{Alb}(\widetilde{Y}_{A}^{\geq 2}), \mathbf{Z}) = \bigwedge^{2} H^{1}(\widetilde{Y}_{A}^{\geq 2}, \mathbf{Z}).$$

The latter group maps, by cup-product, to $H^2(\widetilde{Y}_A^{\geq 2}, \mathbb{Z})^+$ (the invariant space for the canonical covering involution on $\widetilde{Y}_A^{\geq 2}$) with finite (but nontrivial) cokernel. The principal polarization maps to a class 3ν in NS $(\widetilde{Y}_A^{\geq 2})^+$, where $\mathscr{O}_{\widetilde{Y}_A^{\geq 2}}(1) = 2\nu$ (thus, ν is the class of one of the components of the curve (1.5) discussed below); this follows from Welters' work on the variety of lines on quartic double solids (see the proof of Proposition 2.5 and [Wel81, (3.32), Proposition (3.60), and p. 70]).

Let X be a smooth GM variety of dimension $n \in \{3, 5\}$ and assume $Y^3_{A(X)} = \emptyset$. To prove the isomorphisms (1.3) and (1.4), it is natural to construct a subscheme or a cycle

$$Z \subset X \times \widetilde{Y}_{A(X)}^{\geq 2}$$

of codimension $\frac{n+1}{2}$ and use the Abel-Jacobi map $AJ_Z: H_1(\widetilde{Y}_{A(X)}^{\geq 2}, \mathbb{Z}) \to H_n(X, \mathbb{Z})$. For this, one needs an interpretation of the double EPW surface $\widetilde{Y}_{A(X)}^{\geq 2}$ (or some other closely related surface) as a moduli space of sheaves or as a parameter space of cycles on X.

When X is a GM threefold, the most natural moduli space of sheaves to consider is the Hilbert scheme of conics on X. This scheme was thoroughly studied in [Log12] and [DIM12, Section 6]; in [DK20+], we prove

that it is isomorphic to the blow up of a point of the dual double EPW surface $\widetilde{Y}_{A(X)^{\perp}}^{\geq 2}$. Similarly, for a GM fivefold X, one could use the Hilbert scheme of quadric surfaces in X; we proved in [DK20+] that it has a connected component isomorphic to a \mathbf{P}^1 -bundle over $\widetilde{Y}_{A(X)^{\perp}}^{\geq 2}$. In the case of GM threefolds, it is claimed in [IM11, Section 5.1] and [IM07, Theorem 9] that the Clemens-Letizia degeneration method can be applied to prove that the Abel-Jacobi map given by the universal conic is an isomorphism; however, it is not clear whether this method would work in the case of GM fivefolds, so we need a different approach.

Another possible approach in the case of a GM threefold X would be to use the moduli space $\mathcal{M}_X(2;1,5)$ of Gieseker semistable rank-2 torsion-free sheaves on X with $c_1 = 1$, $c_2 = 5$, and $c_3 = 0$. This space was shown in [DIM12, Section 8] to be birational to the Hilbert scheme of conics on X', a line transform of X (see Section 4.5), hence to the double EPW surface $\widetilde{Y}_{A(X)}^{\geq 2}$; the natural correspondence is provided by the second Chern class of the universal sheaf on the product $X \times \mathcal{M}_X(2;1,5)$. We use a small modification of this construction in which the moduli space of sheaves is kept implicit. We explain it below.

If X is a GM threefold and $L_0 \subset X$ is a line, the corresponding (inverse) line transform $X' \to X$ takes a general conic $C' \subset X'$ to a rational quartic curve $C \subset X$ to which the line L_0 is bisecant (the corresponding rank-2 sheaf on X can then be obtained by Serre's construction applied to $C \cup L_0$; in particular, the curve $C \cup L_0$ represents the second Chern class of this sheaf). We consider the union $C \cup L_0$ as a quintic curve of arithmetic genus 1 on X containing the line L_0 and construct the correspondence Z as the closure of a family of such curves parameterized by an open subscheme of $\widetilde{Y}_{A(X)}^{\geq 2}$.

To prove that the Abel–Jacobi map AJ_Z associated with this family of curves is an isomorphism, we make the crucial observation that over the curve

(1.5)
$$Y_{A(X),V_5(X)}^{\geq 2} := Y_{A(X)}^{\geq 2} \cap \mathbf{P}(V_5(X)),$$

the double covering $\widetilde{Y}_{A(X)}^{\geq 2} \to Y_{A(X)}^{\geq 2}$ splits, and that over a general point y of one of the components of its preimage, there is a relation $Z_y + L_y = S_y \cap X$ in the Chow group $CH_1(X)$ of 1-cycles. Here Z_y is the fiber of the correspondence Z over y, L_y is a line on X, and S_y is a cubic surface scroll on the fourfold $M_X := CGr(2, V_5) \cap \mathbf{P}(W)$. Moreover, the curve (1.5) is birational to the Hilbert scheme $F_1(X)$ of lines on X and the line L_y comes from the universal family of lines over $F_1(X)$.

From these observations and from the vanishing of the odd cohomology of M_X , it follows that for X general, there is a morphism $\phi: F_1(X) \to \widetilde{Y}_{A(X)}^{\geq 2}$ such that the composition

$$H_1(F_1(X), \mathbf{Z}) \xrightarrow{\phi_*} H_1(\widetilde{Y}_{A(X)}^{\geq 2}, \mathbf{Z}) \xrightarrow{\mathsf{AJ}_Z} H_3(X, \mathbf{Z})$$

is the opposite of the Abel-Jacobi map defined by the universal family of lines. The latter map is surjective by an argument of Clemens-Tyurin (see Section 3.3), hence AJ_Z is surjective as well. It is not hard to check that the source and target of AJ_Z are free abelian groups of rank 20, hence AJ_Z is an isomorphism.

A similar argument works for GM fivefolds: rational quartic curves are replaced by rational quartic surface scrolls, reducible quintic curves by reducible quintic del Pezzo surfaces, the Hilbert scheme of lines by a component of the Hilbert scheme of planes, and a higher-dimensional analogue of the Clemens–Tyurin argument is applied.

For GM fivefolds X, the isomorphism (1.4) may be proved by a completely different topological argument. When X is general, we consider the double cover $\widetilde{Y}_{A(X),V_5(X)}^{\geq 2}$ of the curve (1.5) induced by the double covering $\widetilde{Y}_{A(X)}^{\geq 2} \to Y_{A(X)}^{\geq 2}$; in contrast with the case of GM threefolds, this is a smooth curve of genus 161. Using classical monodromy arguments, we prove that its Jacobian has three simple factors: the Jacobian of the curve $Y_{A(X),V_5(X)}^{\geq 2}$ (of dimension 81), the Albanese variety of the surface $\widetilde{Y}_{A(X)}^{\geq 2}$ (of dimension 10), and a simple factor of dimension 70. The curve $\widetilde{Y}_{A(X),V_5(X)}^{\geq 2}$ parameterizes planes on X (see Section 2.5.1) and the corresponding Abel-Jacobi map

$$H_1(\widetilde{Y}_{A(X),V_5(X)}^{\geq 2}, \mathbb{Z}) \longrightarrow H_5(X, \mathbb{Z})$$

is surjective by a generalization of the Clemens-Tyurin argument. The induced surjective morphism

$$\operatorname{Jac}(\widetilde{Y}_{A(X),V_5(X)}^{\geq 2}) \longrightarrow \operatorname{Jac}(X)$$

therefore has connected kernel. The description of the simple factors implies that it has to be isogeneous to the product of the 81-dimensional and 70-dimensional factors. Therefore, Jac(X) is isomorphic to the remaining 10-dimensional factor Alb($\widetilde{Y}_{A(X)}^{\geq 2}$).

To complete the proof of Theorem 1.1 and to describe the period maps for GM varieties of dimension 3 or 5, we investigate the rational map

(1.6)
$$\mathbf{M}^{\text{EPW}} = \text{LGr}(\Lambda^3 V_6) / / \text{PGL}(V_6) \rightarrow \mathbf{A}_{10}$$

from the coarse moduli space of EPW sextics (constructed in [O'Gr08a, Section 6]) to the coarse moduli space of principally polarized abelian varieties of dimension 10 defined by

$$[A] \longmapsto [\operatorname{Alb}(\widetilde{Y}_A^{\geq 2})]$$

when A has no decomposable vectors and $Y_A^{\geq 3} = \emptyset$. Let $\mathbf{M}_{ndv}^{EPW} \subset \mathbf{M}^{EPW}$ be the open subset parameterizing Lagrangian subspaces with no decomposable vectors and let \mathbf{r} be the involution of \mathbf{M}_{ndv}^{EPW} defined by $[A] \mapsto [A^{\perp}]$ (see [O'Gr08b]). We show in Proposition 6.2 that the map (1.6) extends to a regular morphism

$$\bar{\wp} \colon \mathbf{M}_{ndv}^{\text{EPW}}/\mathbf{r} \longrightarrow \mathbf{A}_{10}$$

such that $\bar{\wp}([A])$ is the Albanese variety of (any desingularization of) the double EPW surface $\widetilde{Y}_A^{\geq 2}$. Let now \mathbf{M}_n^{GM} be the coarse moduli space of GM varieties of dimension *n* (see [DK20b] and Section 2.3). We use the above result to prove the following

Theorem 1.3. For $n \in \{3, 5\}$, the period map $\wp_n \colon \mathbf{M}_n^{\mathrm{GM}} \to \mathbf{A}_{10}$ factors as the composition

$$\varphi_n\colon \mathbf{M}_n^{\mathrm{GM}} \longrightarrow \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} \longrightarrow \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}}/\mathbf{r} \stackrel{\bar{\varphi}}{\longrightarrow} \mathbf{A}_{10},$$

where the first map is given by $[X] \mapsto [A(X)]$ and the second map is the canonical projection. In particular, $\varphi_n([X]) = [\operatorname{Alb}(\widetilde{Y}_{A(X)}^{\geq 2})].$

This factorization of the period map for GM threefolds was discussed in the introduction of [DIM12] (see also [DIM12, Remark 7.5]); moreover, it was conjectured there that the map $\bar{\wp}$ is generically injective (the computation in [DIM12, Theorem 5.1] shows that it has finite fibers).

The story of GM threefolds is very similar to the story of quartic double solids. The articles [Wel81, Voi88] were an inspiration to us; in particular, we took the idea of using the Clemens-Tyurin argument from [Wel81]

The article is organized as follows. In Section 2, we review the theory of GM varieties, EPW varieties, and their double covers. In particular, we describe the Hilbert scheme of lines on GM threefolds, the Hilbert scheme of σ -planes on GM fivefolds, and we identify double EPW varieties with the canonical covers of degeneracy loci for the family of quadrics containing a GM variety.

In Section 3, we recall basic facts about Abel-Jacobi maps, prove a generalization of the Clemens-Tyurin argument, and discuss the endomorphism ring of intermediate Jacobians; in particular, we check that the intermediate Jacobian of a very general GM variety of odd dimension is simple and has Picard number 1 (this had already been proved for GM threefolds by a different argument in [DIM12, Corollary 5.3]).

In Section 4, we construct, for any GM threefold X, a cycle $Z \subset X \times \widetilde{Y}_{A(X)}^{\geq 2}$, and prove that the Abel-Jacobi map defined by Z is an isomorphism when $Y_{A(X)}^{\geq 3} = \emptyset$. We also describe how the line transform of GM threefolds acts on their coarse moduli spaces. In Section 5, we prove analogous results for GM fivefolds. Finally, we describe in Section 6 the period map for GM threefolds and fivefolds and prove Theorems 1.1 and 1.3.

Acknowledgements. We would like to thank Kieran O'Grady, Andrey Soldatenkov, and Claire Voisin for useful discussions. We also thank the anonymous referee for her or his careful reading of our manuscript and useful comments.

2. Gushel-Mukai and Eisenbud-Popescu-Walter varieties

We work over the field of complex numbers. Given a subvariety X of a projective space, we denote by $F^k(X)$ the Hilbert scheme parameterizing linear spaces of dimension k in X.

2.1. Geometry of Gr(2, 5)

Let V_5 be a 5-dimensional vector space. A subspace of V_5 of dimension k will usually be denoted by V_k or U_k . We denote by

$$\operatorname{Gr}(2, V_5) \subset \mathbf{P}(\wedge^2 V_5)$$

the Grassmannian of 2-dimensional vector subspaces in V_5 in its Plücker embedding.

We recall some standard facts about its geometry. It has codimension 3 and degree 5 and is the intersection, for $v \in V_5 \setminus \{0\}$, of the *Plücker quadrics*

(2.1)
$$Q_{v} := \operatorname{Cone}_{\mathbf{P}(v \wedge V_{5})}(\operatorname{Gr}(2, V_{5}/\mathbf{C}v)) \subset \mathbf{P}(\wedge^{2}V_{5}).$$

In the next lemma, we describe Hilbert schemes of linear subspaces on $Gr(2, V_5)$.

Lemma 2.1. We have the following isomorphisms:

- (a) $F^1(Gr(2, V_5)) \simeq Fl(1, 3; V_5)$; the line corresponding to a flag $V_1 \subset V_3 \subset V_5$ is the set of all $[U_2] \in Gr(2, V_5)$ such that $V_1 \subset U_2 \subset V_3$.
- (b) F²(Gr(2, V₅)) = Fl(1,4; V₅) ⊔ Gr(3, V₅); the plane corresponding to a flag V₁ ⊂ V₄ ⊂ V₅ is the set of all [U₂] ∈ Gr(2, V₅) such that V₁ ⊂ U₂ ⊂ V₄, and the plane corresponding to a subspace V₃ ⊂ V₅ is the set of all [U₂] ∈ Gr(2, V₅) such that U₂ ⊂ V₃.
- (c) $F^3(Gr(2, V_5)) \simeq \mathbf{P}(V_5)$; the linear 3-space corresponding to a subspace $V_1 \subset V_5$ is the set of all 2-spaces $[U_2] \in Gr(2, V_5)$ such that $V_1 \subset U_2$.
- (d) $F^4(Gr(2, V_5)) = \emptyset$: there are no linear 4-spaces on $Gr(2, V_5)$.

Planes on Gr(2, V_5) parameterized by the two components in Lemma 2.1(b) are traditionally known as σ -planes and τ -planes. We use the notation

$$F_{\sigma}^2(\operatorname{Gr}(2, V_5)) \simeq \operatorname{Fl}(1, 4; V_5)$$

for the connected component of $F^2(Gr(2, V_5))$ parameterizing σ -planes.

If a finite morphism $\gamma: X \to Gr(2, V_5)$ is compatible with the polarizations, it induces a morphism $F^k(\gamma): F^k(X) \to F^k(Gr(2, V_5))$ between Hilbert schemes and we denote by

$$F_{\sigma}^{2}(X) \subset F^{2}(X)$$

the preimage of $F_{\sigma}^2(\text{Gr}(2, V_5))$.

We will need the following classical result.

Lemma 2.2. Let $V_2 \subset V_5$ be a 2-dimensional subspace. We have the equality

$$\operatorname{Gr}(2, V_5) \cap \mathbf{P}(V_2 \wedge V_5) = \operatorname{Cone}_{\mathbf{P}(\wedge^2 V_2)}(\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2))$$

in $\mathbf{P}(\bigwedge^2 V_5)$, where the right side is the cone over the cubic scroll.

Typically, an intersection $Gr(2, V_5) \cap \mathbf{P}^5$ is 2-dimensional (and is a quintic del Pezzo surface). In the next lemma, we discuss some pathological intersections.

Lemma 2.3. Assume $\mathbf{P}^5 \subset \mathbf{P}(\wedge^2 V_5)$ is a linear subspace such that $\dim(\operatorname{Gr}(2, V_5) \cap \mathbf{P}^5) = 3$. The only possible 3-dimensional component of $\operatorname{Gr}(2, V_5) \cap \mathbf{P}^5$ of even degree is a hyperplane section of some $\operatorname{Gr}(2, V_4) \subset \operatorname{Gr}(2, V_5)$.

Proof. Write $\mathbf{P}^5 = \mathbf{P}(W_6)$, where $W_6 \subset \bigwedge^2 V_5$. Let $W_6^{\perp} \subset \bigwedge^2 V_5^{\vee}$ be the orthogonal complement of W_6 and set $T := \mathbf{P}(W_6^{\perp}) \cap \operatorname{Gr}(2, V_5^{\vee})$. This is a subscheme of $\mathbf{P}(W_6^{\perp}) \simeq \mathbf{P}^3$, hence $\dim(T) \leq 3$. We discuss all the possibilities for $\dim(T)$ and check the claim in each case.

Assume first dim $(T) \leq 1$. We then have $\mathbf{P}(W_8^{\perp}) \cap \operatorname{Gr}(2, V_5^{\vee}) = \emptyset$ for a general subspace $W_8 \subset \bigwedge^2 V_5$ containing W_6 : hence $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_8)$ is a smooth quintic del Pezzo fourfold ([DK18, Proposition 2.24]) not contained in a hyperplane. Its Picard group is \mathbf{Z} , hence for any subspace $W_7 \subset W_8$ containing W_6 , its hyperplane section $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_7)$ is an irreducible threefold not contained in a hyperplane. Therefore, the hyperplane section $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_6)$ of this threefold has no 3-dimensional components.

Assume now dim(T) = 2. Since Gr $(2, V_5^{\vee})$ is an intersection of quadrics, so is T, hence T is either an irreducible quadric surface or contains a plane.

If *T* contains a plane, we have, by Lemma 2.1(b), either $\mathbf{P}(\bigwedge^2 V_2^{\perp}) \subset T$, in which case $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_6)$ is a hyperplane section of $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(V_2 \wedge V_5)$ hence, by Lemma 2.2, an irreducible threefold of degree 3, or $\mathbf{P}(V_1^{\perp} \wedge V_4^{\perp}) \subset T$, in which case $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_6)$ is a hyperplane section of $\operatorname{Gr}(2, V_4) \cup \mathbf{P}(V_1 \wedge V_4)$, hence is the union of a hyperplane section of $\operatorname{Gr}(2, V_4)$ and of a linear 3-space.

If *T* is an irreducible quadric, we have $\mathbf{P}(W_6^{\perp}) \subset \mathbf{P}(\bigwedge^2 V_1^{\perp})$ and $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_6)$ is the union of $\mathbf{P}(V_1 \wedge V_5)$ and two planes (or a double plane) corresponding to the intersection of $\operatorname{Gr}(2, V_5/V_1)$ with the line \mathbf{P}^1 given by the orthogonal of $W_6^{\perp} \subset \bigwedge^2 V_1^{\perp}$. Therefore, the only 3-dimensional component of $\operatorname{Gr}(2, V_5) \cap \mathbf{P}(W_6)$ has degree 1.

Finally, assume dim(T) = 3. By Lemma 2.1(c), we have $W_6^{\perp} = V_4^{\perp} \wedge V_5^{\vee}$ and $Gr(2, V_5) \cap \mathbf{P}(W_6) = Gr(2, V_4)$ has dimension 4.

Therefore, the only case when $Gr(2, V_5) \cap \mathbf{P}(W_6)$ has a 3-dimensional component of even degree is the case when T contains a σ -plane, and in this case, this component is a hyperplane section of $Gr(2, V_4)$. \Box

We will also need the following standard locally free resolution for the cone $CGr(2, V_5)$.

Lemma 2.4. There is an exact sequence

 $0 \to \mathscr{O}(-5) \to V_5^{\vee} \otimes \mathscr{O}(-3) \to V_5 \otimes \mathscr{O}(-2) \to \mathscr{O} \to \mathscr{O}_{\mathsf{CGr}(2,V_5)} \to 0$

of coherent sheaves on $\mathbf{P}(\mathbf{C} \oplus \bigwedge^2 V_5)$.

2.2. Eisenbud-Popescu-Walter varieties and their double coverings

Let V_6 be a 6-dimensional vector space. We consider subspaces $A \subset \bigwedge^3 V_6$ that are Lagrangian for the symplectic form given by exterior product. Those that contain no decomposable vectors (that is, such that $\mathbf{P}(A) \cap \operatorname{Gr}(3, V_6) = \emptyset$) are parameterized by the complement

(2.3)
$$\mathsf{LGr}_{\mathsf{ndv}}(\wedge^3 V_6) \subset \mathsf{LGr}(\wedge^3 V_6)$$

of a hypersurface in the Lagrangian Grassmannian $LGr(\bigwedge^3 V_6)$.

Given a Lagrangian subspace $A \subset \bigwedge^3 V_6$, one defines its EPW varieties; they form a chain

$$Y_A^{\geq 4} \subset Y_A^{\geq 3} \subset Y_A^{\geq 2} \subset Y_A^{\geq 1} \subset Y_A^{\geq 0} = \mathbf{P}(V_6)$$

of closed subschemes, where $Y_A^{\geq k}$ is, at least set-theoretically, the set of points [v] in $\mathbf{P}(V_6)$ such that $\dim(A \cap (v \wedge \bigwedge^2 V_6)) \geq k$ (the scheme structures were defined in [DK20a, (18)]).

The Lagrangian subspace $A \subset \bigwedge^3 V_6$ defines a Lagrangian subspace $A^{\perp} \subset \bigwedge^3 V_6^{\vee}$ hence, as above, dual EPW varieties

$$Y_{A^{\perp}}^{\geq 4} \subset Y_{A^{\perp}}^{\geq 3} \subset Y_{A^{\perp}}^{\geq 2} \subset Y_{A^{\perp}}^{\geq 1} \subset Y_{A^{\perp}}^{\geq 0} = \mathbf{P}(V_6^{\vee}).$$

Set-theoretically, the variety $Y_{A^{\perp}}^{\geq k}$ is the set of points $[V_5]$ in $Gr(5, V_6) = \mathbf{P}(V_6^{\vee})$ such that $\dim(A \cap \bigwedge^3 V_5) \geq k$. We also use the notation

$$Y_A^k = Y_A^{\geq k} \smallsetminus Y_A^{\geq k+1} \quad \text{and} \quad Y_{A^\perp}^k = Y_{A^\perp}^{\geq k} \smallsetminus Y_{A^\perp}^{\geq k+1}.$$

Assume for the rest of this section that A contains no decomposable vectors. A combination of results of O'Grady (see [DK18, Theorem B.2]) gives the following:

- Y_A := Y_A^{≥1} is a normal sextic hypersurface (called an EPW sextic);
 Y_A^{≥2} = Sing(Y_A) is a normal integral surface of degree 40 (called an EPW surface);
 Y_A^{≥3} = Sing(Y_A^{≥2}) is a finite scheme, empty when A is general;
 Y_A^{≥4} is empty.

Note that the Lagrangian subspace A^{\perp} also contains no decomposable vectors and analogous statements hold for dual EPW varieties.

EPW varieties have canonical double coverings. First, there is a double covering

$$\widetilde{Y}_A^{\geq 0} \longrightarrow Y_A^{\geq 0} = \mathbf{P}(V_6)$$

branched along the EPW sextic Y_A . Next, O'Grady constructed in [O'Gr13, Section 1.2] a canonical double covering

$$\widetilde{Y}_A \longrightarrow Y_A$$

étale away from $Y_A^{\geq 2}$. When $Y_A^{\geq 3} = \emptyset$, the scheme \widetilde{Y}_A is a smooth hyperkähler fourfold (called a double EPW sextic). Finally, in [DK20a, Theorem 5.2(2)], we constructed a canonical double covering

(2.4)
$$\pi_A \colon \widetilde{Y}_A^{\geq 2} \longrightarrow Y_A^{\geq 2}$$

étale away from $Y_A^{\geq 3}$, where $\widetilde{Y}_A^{\geq 2}$ is an integral normal surface (called a double EPW surface), and proved an isomorphism

(2.5)
$$\pi_{A*}\mathscr{O}_{\widetilde{Y}_A^{\geq 2}} \simeq \mathscr{O}_{Y_A^{\geq 2}} \oplus \omega_{Y_A^{\geq 2}}(-3)$$

The double coverings π_A are the main characters of this article. We now prove some results about the surface $\widetilde{Y}_A^{\geq 2}$ that will be needed later on.

Proposition 2.5. Let A be a Lagrangian subspace with no decomposable vectors and assume $Y_A^{\geq 3} = \emptyset$, so that $\widetilde{Y}_A^{\geq 2}$ and $Y_A^{\geq 2}$ are smooth connected projective surfaces. One has

(2.6)
$$H^1(Y_A^{\geq 2}, \mathscr{O}_{Y_A^{\geq 2}}) = 0, \qquad H^1(\widetilde{Y}_A^{\geq 2}, \mathscr{O}_{\widetilde{Y}_A^{\geq 2}}) \simeq A,$$

where the isomorphism is canonical, and the abelian group $H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z})$ is free of rank 20.

Proof. From (2.5) and Serre duality, we deduce that there are isomorphisms

$$\begin{split} H^1(\widetilde{Y}_A^{\geq 2}, \mathscr{O}_{\widetilde{Y}_A^{\geq 2}}) &\simeq H^1(Y_A^{\geq 2}, \mathscr{O}_{Y_A^{\geq 2}}) \oplus H^1(Y_A^{\geq 2}, \omega_{Y_A^{\geq 2}}(-3)) \\ &\simeq H^1(Y_A^{\geq 2}, \mathscr{O}_{Y_A^{\geq 2}}) \oplus H^1(Y_A^{\geq 2}, \mathscr{O}(3))^{\vee}. \end{split}$$

From the table in [DK19, Corollary B.5], we see that the first summand vanishes, whereas the second summand is canonically isomorphic to A. This proves the statements (2.6) of the proposition.

To prove that $H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z})$ is torsion-free, we use a degeneration argument. Let $S \subset \mathbb{P}^3$ be a smooth quartic surface containing no lines. Ferretti proved in [Fer12, Proposition 4.1 and Corollary 4.2] that there is a smooth deformation of the surface $Y_A^{\geq 2}$ to the surface Bit(S) \subset Gr(2, 4) of bitangent lines to S.

Let $X \to \mathbf{P}^3$ be the double solid branched over S. The variety $F_1(X)$ of lines on X is a connected surface ([Wel81, Remark (3.58)a)]) and the canonical map $F_1(X) \rightarrow Bit(S)$ is a double étale cover ([Wel81, Corollary (1.3)]) whose associated order-2 line bundle on Bit(S) is $\omega_{\text{Bit}(S)}(-3)$ ([Wel81, Propositions (3.1)a) and (3.35)]). It then follows from (2.5) that the surface $F_1(X)$ is a smooth deformation of the surface $\widetilde{Y}_A^{\geq 2}$. They are therefore diffeomorphic. The statement that $H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z})$ is free of rank 20 then follows from the analogous statement for $H_1(F_1(X), \mathbb{Z})$ proved in [Wel81, Section 6, Proposition, p. 71].

2.3. Gushel-Mukai varieties

Let $n \in \{3, 4, 5, 6\}$. As recalled in the introduction (see (1.1)), a Gushel-Mukai variety of dimension n is a dimensionally transverse intersection

$$X = \mathsf{CGr}(2, V_5) \cap \mathbf{P}(W) \cap Q.$$

It is the intersection in $\mathbf{P}(W)$ of the 6-dimensional space $V_6(X) \subset \text{Sym}^2(W^{\vee})$ of quadrics containing X, generated by the space

$$V_5(X) := V_5 = \bigwedge^4 V_5^{\vee} \subset \operatorname{Sym}^2(\bigwedge^2 V_5^{\vee})$$

of (the restrictions to W of) Plücker quadrics and the quadric Q. In particular, one can replace Q by any other quadric in the space $V_6(X) \setminus V_5(X)$. The intersection

$$(2.7) M_X := \operatorname{CGr}(2, V_5) \cap \mathbf{P}(W)$$

is called the Grassmannian hull of X. There are two types of GM varieties:

- if M_X does not contain the vertex of the cone CGr(2, V₅), then M_X ≃ Gr(2, V₅) ∩ P(W) is a linear section of Gr(2, V₅) and X = M_X ∩ Q is a quadratic section of M_X; these GM varieties are called ordinary;
- if M_X contains the vertex of the cone CGr(2, V₅), then M_X is a cone over M'_X = Gr(2, V₅) ∩ P(W'), a linear section of Gr(2, V₅), and X → M'_X is a double covering branched along a quadratic section X' = M'_X ∩ Q'; these GM varieties are called special.

When X is a special GM variety of dimension n, the variety X' is an ordinary GM variety of dimension n-1; the varieties X and X' are called opposite GM varieties.

With every GM variety X, we associated in [DK18, Section 3.2] a Lagrangian data set $(V_6(X), V_5(X), A(X))$ consisting of

- the 6-dimensional space $V_6(X)$ of quadrics containing X,
- the hyperplane $V_5(X) \subset V_6(X)$ of Plücker quadrics,
- a Lagrangian subspace $A(X) \subset \bigwedge^3 V_6(X)$.

The Lagrangian data sets of a GM variety and of its opposite GM variety coincide.

Many properties of X are related to properties of its Lagrangian data set. For instance, when X is smooth and dim $(X) \ge 3$, the space A(X) contains no decomposable vectors ([DK18, Theorem 3.16]) and dim $(A(X) \cap \bigwedge^3 V_5(X)) \le 3$.

Conversely, if (V_6, V_5, A) is a Lagrangian data set such that A contains no decomposable vectors and $\ell = \dim(A \cap \bigwedge^3 V_5) \leq 3$, there are exactly two smooth GM varieties X such that

$$(V_6(X), V_5(X), A(X)) = (V_6, V_5, A):$$

one ordinary GM variety of dimension $5 - \ell$ and one special GM variety of dimension $6 - \ell$ ([DK18, Theorem 3.10]); they are opposite of one another.

In [DK20b], we upgraded the above constructions to a description of the moduli stack $\mathfrak{M}_n^{\text{GM}}$ of smooth GM varieties of dimension n and its coarse moduli space \mathbf{M}_n^{GM} . In particular, we showed in [DK20b, Theorem 5.15(a)] that the coarse moduli space of smooth GM varieties of dimension $n \ge 3$ is the quasiprojective GIT quotient

(2.8)
$$\mathbf{M}_{n}^{\mathrm{GM}} = \{(A, V_{5}) \in \mathrm{LGr}_{\mathrm{ndv}}(\bigwedge^{3} V_{6}) \times \mathbf{P}(V_{6}^{\vee}) \mid \dim(A \cap \bigwedge^{3} V_{5}) \in \{5 - n, 6 - n\}\} / / \mathrm{PGL}(V_{6})$$

(see (2.3) for the notation). In particular, as explained in [DK20b, Section 6.1], there is a map

(2.9)
$$\rho_n \colon \mathbf{M}_n^{\mathrm{GM}} \longrightarrow \mathsf{LGr}_{\mathrm{ndv}}(\bigwedge^3 V_6) / / \mathsf{PGL}(V_6)$$
$$[X] \longmapsto [A(X)]$$

and

(2.10)
$$p_n^{-1}([A]) \simeq (Y_{A^{\perp}}^{\geq 5-n} \smallsetminus Y_{A^{\perp}}^{\geq 7-n}) / / PGL(V_6)_A$$

(when n = 6, the right side is $Y_{A\perp}^0 / / PGL(V_6)_A$), where $PGL(V_6)_A$ is the stabilizer of A in $PGL(V_6)$, a finite (generically trivial) group ([DK18, Proposition B.9]). When $n \in \{4, 6\}$, we showed in [DK19, Proposition 5.27] (see also [DK20b, Proposition 6.1]) that the map ρ_n can be thought of as the period map for GM n-folds.

We computed in [DK19] the integral cohomology groups of GM varieties of dimensions 3 or 5 and of their Grassmannian hulls. We denote by $F^{\bullet}H^n(X, \mathbb{C})$ the Hodge filtration on the cohomology $H^n(X, \mathbb{C})$.

Proposition 2.6. Let X be a smooth GM variety of odd dimension $n \in \{3, 5\}$.

- The even cohomology $H^{\text{even}}(X, \mathbb{Z})$ is pure Tate of ranks (1, 1, 1, 1) when n = 3, and (1, 1, 2, 2, 1, 1) when n = 5.
- For the odd cohomology, we have

$$H^{\text{odd}}(X, \mathbb{Z}) = H^n(X, \mathbb{Z}) \simeq \mathbb{Z}^{20}, \qquad F^{(n+3)/2} H^n(X, \mathbb{C}) = 0.$$

If X is moreover ordinary, so that its Grassmannian hull M_X is smooth,

- the even cohomology $H^{\text{even}}(M_X, \mathbb{Z})$ is pure Tate of ranks (1, 1, 2, 1, 1) when n = 3, and (1, 1, 2, 2, 2, 1, 1) when n = 5;
- the odd cohomology $H^{\text{odd}}(M_X, \mathbb{Z})$ vanishes.

Proof. The first part follows from [DK19, Propositions 3.1 and 3.4]. The second part is a standard consequence of the Lefschetz Theorem. \Box

We will also need the following result.

Lemma 2.7. A smooth GM fivefold contains no quadric threefold whose image in $Gr(2, V_5)$ is a hyperplane section of some $Gr(2, V_4)$.

Proof. Let $Q \subset CGr(2, V_5)$ be a quadric threefold contained in a smooth GM fivefold X. Then Q does not contain the vertex of the cone $CGr(2, V_5)$ (because X does not), hence its projection from the vertex to $Gr(2, V_5)$ is well defined. Assume it is a hyperplane section of some $Gr(2, V_4)$. Then Q is a local complete intersection and its normal bundle splits as

$$N_{O/CGr(2,V_5)} \simeq \mathscr{U}^{\vee} \oplus \mathscr{O}(1)^{\oplus 2}$$

where \mathscr{U} is the restriction of the tautological bundle of $Gr(2, V_5)$. Any GM fivefold X is the intersection of a hyperplane and a quadric in $CGr(2, V_5)$. If X contains Q, the differentials of the equations of X give a morphism

$$N_{O/CGr(2,V_5)} \rightarrow \mathscr{O}(1) \oplus \mathscr{O}(2).$$

Clearly, X is singular at any degeneracy point of that morphism. If X is smooth, this morphism is therefore surjective, hence its kernel is a vector bundle of rank 2. This is absurd, since a simple computation shows that its third Chern class is nonzero. Therefore, X cannot contain Q.

2.4. Linear spaces on quadrics containing GM varieties

If $X \subset \mathbf{P}(W)$ is a GM variety and $V_6(X)$ is the space of quadrics in $\mathbf{P}(W)$ containing X, we denote by

$$(2.11) \qquad \qquad \mathscr{Q} \subset \mathbf{P}(W) \times \mathbf{P}(V_6(X))$$

the total space of this family of quadrics and, for $v \in V_6(X)$ nonzero, by Q_v the corresponding quadric in $\mathbf{P}(W)$.

The Lagrangian data set associated with X can be used to describe the ranks of the family of quadrics (2.11): by [DK18, Proposition 3.13(b)], we have

(2.12)
$$\operatorname{Ker}(Q_{\nu}) = A(X) \cap (\nu \wedge \bigwedge^{2} V_{6}(X)) \quad \text{for all } \nu \in V_{6}(X) \setminus V_{5}(X).$$

In particular, $Y_{A(X)}^k \setminus \mathbf{P}(V_5(X))$ is the locus of non-Plücker quadrics of corank k containing X.

In fact, the family of quadrics (2.11) itself can be reconstructed from the Lagrangian data set, which allows us to relate the double covering (2.4) to the coverings associated with the family of quadrics by [DK20a, Theorem 3.1]. Note that (2.11) corresponds to an embedding

$$\mathscr{O}_{\mathbf{P}(V_6(X))}(-1) \hookrightarrow \operatorname{Sym}^2 W^{\vee} \otimes \mathscr{O}_{\mathbf{P}(V_6(X))}.$$

On $\mathbf{P}(V_6(X)) \setminus \mathbf{P}(V_5(X))$, the line bundle $\mathscr{O}_{\mathbf{P}(V_6(X))}(-1)$ is trivial, hence double coverings of any quadratic degeneracy loci are well defined over that set by [DK20a, Remark 3.2].

Lemma 2.8. Let X be a smooth GM variety of dimension n, with associated Lagrangian data set (V_6, V_5, A) and let $k \in \{0, 1, 2\}$. Over $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$, the canonical double covering of the k-th degeneracy locus $Y_A^{\geq k} \setminus \mathbf{P}(V_5)$ of the family of quadrics (2.11) coincides with the base change $\widetilde{Y}_A^{\geq k} \times_{\mathbf{P}(V_6)} (\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)) \to Y_A^{\geq k} \setminus \mathbf{P}(V_5)$ of the double covering $\widetilde{Y}_A^{\geq k} \to Y_A^{\geq k}$ defined in Section 2.2.

Proof. By [DK20a, Theorem 5.2], the double covering $\widetilde{Y}_A^{\geq k} \to Y_A^{\geq k}$ is associated with the pair of Lagrangian subbundles

$$\mathscr{A}_1 = A \otimes \mathscr{O}$$
 and $\mathscr{A}_2 = \bigwedge^2 T_{\mathbf{P}(V_6)}(-3)$

of $\bigwedge^3 V_6 \otimes \mathscr{O}$ over $\mathbf{P}(V_6)$. To identify the double coverings, we will show that the family of quadrics (2.11) can be related to this pair by the isotropic reduction procedure of [DK20a, Section 4.2]. We will use freely the notation from [DK20a].

Asssume first that X is ordinary. We restrict \mathscr{A}_1 and \mathscr{A}_2 to $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$ and consider the third Lagrangian subbundle

$$\mathscr{A}_3 := \bigwedge^3 V_5 \otimes \mathscr{O}$$

of $\bigwedge^3 V_6 \otimes \mathscr{O}$ over $\mathbf{P}(V_6) \smallsetminus \mathbf{P}(V_5)$. We apply isotropic reduction with respect to the rank-2 subbundle

$$\mathscr{I} := \mathscr{A}_1 \cap \mathscr{A}_3 = (A \cap \bigwedge^3 V_5) \otimes \mathscr{O}$$

in the sense of [DK20a, Section 4.2] and obtain three Lagrangian subbundles $\overline{\mathscr{A}}_1$, $\overline{\mathscr{A}}_2$, $\overline{\mathscr{A}}_3$ in a symplectic vector bundle $\overline{\mathscr{V}}$. We describe below all these bundles explicitly.

For each $[v] \in \mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$, there is a Lagrangian direct sum decomposition

$$\bigwedge^3 V_6 = \bigwedge^3 V_5 \oplus (v \land \bigwedge^2 V_5)$$

and the respective fibers at [v] of the bundles \mathscr{A}_1 , \mathscr{A}_2 , \mathscr{A}_3 are A, $v \wedge \bigwedge^2 V_5$ (the second summand), and $\bigwedge^3 V_5$ (the first summand). Furthermore, the fiber of \mathscr{I} is the subspace $I := A \cap \bigwedge^3 V_5$ of the first summand. By [DK18, Proposition 3.13(a)], the space I can be identified with the space of linear functions on $\bigwedge^2 V_5$ vanishing on W, hence $\bigwedge^3 V_5/I \simeq W^{\vee}$. Thus

$$\overline{\mathscr{V}}_{[v]} = W^{\vee} \oplus W$$

and the fibers at [v] of the bundles $\overline{\mathscr{A}}_1$, $\overline{\mathscr{A}}_2$, $\overline{\mathscr{A}}_3$ are A/I, W (the second summand), and W^{\vee} (the first summand). In particular, $\overline{\mathscr{A}}_2 \cap \overline{\mathscr{A}}_3 = 0$. Moreover, $\overline{\mathscr{A}}_1 \cap \overline{\mathscr{A}}_3 = 0$. Indeed, any vector in the intersection

comes from $(A + I) \cap \bigwedge^3 V_5$, and since $I \subset A$, it belongs to $A \cap \bigwedge^3 V_5 = I$, hence corresponds to the zero vector in $\overline{\mathscr{A}}_1 \cap \overline{\mathscr{A}}_3$. This implies $\overline{\mathscr{A}}_3 \simeq W^{\vee} \otimes \mathscr{O}$, and the maps [DK20a, (23)] for the triple $\overline{\mathscr{A}}_1, \overline{\mathscr{A}}_2, \overline{\mathscr{A}}_3$ and the trivial line bundle $\mathscr{L} = \mathscr{O}$ are isomorphisms over $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$. Therefore, the construction [DK20a, (24)] defines a family of quadratic forms on the trivial vector bundle $\overline{\mathscr{A}}_3^{\vee} \simeq W \otimes \mathscr{O}$ over $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$.

By [DK18, proof of Theorem 3.6 and Appendix C], this family of quadrics coincides with the restriction of \mathscr{Q} to $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$. Applying [DK20a, Propositions 4.5 and 4.7], we see that the associated double coverings coincide with $\widetilde{Y}_A^{\geq k} \times_{\mathbf{P}(V_6)} (\mathbf{P}(V_6) \setminus \mathbf{P}(V_5))$. This completes the proof of the lemma for ordinary GM varieties.

Assume now that X is a special GM variety. By [DK18, Lemma 2.33], there is a canonical direct sum decomposition $W = W_0 \oplus W_1$, with dim $(W_1) = 1$, such that for the family of quadrics $\mathbf{q} \colon V_6 \to \operatorname{Sym}^2 W^{\vee}$ defining X, we have

$$\mathbf{q} = \mathbf{q}_0 + \mathbf{q}_1, \qquad \mathbf{q}_0 \colon V_6 \to \operatorname{Sym}^2 W_0^{\vee}, \quad \mathbf{q}_1 \colon V_6 \to V_6/V_5 \xrightarrow{\sim} \operatorname{Sym}^2 W_1^{\vee},$$

and the family of quadrics \mathbf{q}_0 corresponds to the ordinary GM variety X_0 opposite to X. By [DK18, Proposition 3.14(c)], the varieties X_0 and X have the same Lagrangian data sets, hence they give rise to the same double coverings $\widetilde{Y}_A^{\geq 2} \to Y_A^{\geq 2}$. Also, the family of quadrics \mathbf{q}_1 is nondegenerate over $\mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$, hence the families of quadrics \mathbf{q} and \mathbf{q}_0 have the same degeneration loci and isomorphic cokernel sheaves. By [DK20a, Theorem 3.1], they induce the same double covers of degeneration loci. We conclude by the argument of the first part of the proof.

We will need the following consequence of the above lemma. Set

Note that $Y_{A,V_5} := Y_{A,V_5}^{\geq 1}$ is a sextic hypersurface, $Y_{A,V_5}^{\geq 2}$ a Cohen-Macaulay curve ([DK18, Lemma B.6]), and $Y_{A,V_5}^{\geq 3}$ a finite scheme. Consider the open surface

(2.14)
$$S_0 := Y_A^{\ge 2} \smallsetminus (Y_{A,V_5}^{\ge 2} \cup Y_A^3) \subset \mathbf{P}(V_6)$$

and the base change $\mathscr{Q}_0 := \mathscr{Q} \times_{\mathbb{P}(V_6)} S_0$ of the family of quadrics (2.11).

Corollary 2.9. Let X be a smooth GM variety of odd dimension n = 2s + 1 with Lagrangian data set (V_6, V_5, A) . Let $\Pi_0 \subset X$ be a linear subspace of dimension s. There is an isomorphism

$$F_{\Pi_0}^{s+3}(\mathscr{Q}_0/S_0) \simeq \widetilde{Y}_A^{\geq 2} \times_{\mathbf{P}(V_6)} S_0$$

of schemes over S_0 , where the left side is the subscheme of the relative Hilbert scheme of linear spaces of dimension s+3 contained in the fibers of $\mathscr{Q}_0 \to S_0$ and containing Π_0 .

Proof. By [DK20a, Proposition 3.10] (with $m = \dim(W) = n + 5 = 2s + 6$ and k = 2), the Stein factorization of the natural morphism $F^{s+3}(\mathscr{Q}_0/S_0) \to S_0$ takes the form

$$F^{s+3}(\mathscr{Q}_0/S_0) \to \widetilde{Y}_A^{\geq 2} \times_{\mathbf{P}(V_6)} S_0 \xrightarrow{\pi_A} Y_A^{\geq 2} \times_{\mathbf{P}(V_6)} S_0 = S_0$$

(we use [DK20a, Lemma 3.9] to check normality of $F^{s+3}(\mathcal{Q}_0/S_0)$ and Lemma 2.8 to identify the second degeneracy locus and its double covering).

As the proof of [DK20a, Proposition 3.10] shows, we have an isomorphism

(2.15)
$$F^{s+3}(\mathscr{Q}_0/S_0) \simeq F^{s+1}(\bar{\mathscr{Q}}_0/S_0),$$

where $\overline{\mathscr{Q}}_0 \to S_0$ is the family of nondegenerate (2s+2)-dimensional quadrics obtained from \mathscr{Q}_0 by passing to the quotients by the kernel spaces of quadrics. Moreover, for any $[v] \in S_0$, if $Q_v \subset \mathbf{P}(W)$ is the fiber of \mathscr{Q}_0 at [v], we have

$$X = \mathsf{CGr}(2, V_5) \cap Q_v$$

hence $X \cap \text{Sing}(Q_v) = \emptyset$ (otherwise X would be singular). Therefore, the space Π_0 intersects none of these kernel spaces hence projects isomorphically onto a space $\bar{\Pi}_0$ in the fiber \bar{Q}_v of \bar{Q}_0 at [v], so that a (s+3)-space in Q_v contains Π_0 if and only if the corresponding (s+1)-space in \bar{Q}_0 contains $\bar{\Pi}_0$. This proves that we also have an isomorphism

$$F_{\Pi_0}^{s+3}(\mathcal{Q}_0/S_0) \simeq F_{\bar{\Pi}_0}^{s+1}(\bar{\mathcal{Q}}_0/S_0)$$

of S_0 -schemes. Finally, since the family of (2s + 2)-dimensional quadrics $\bar{\mathscr{Q}}_0$ is everywhere nondegenerate, it follows from [KS18, Lemma 2.12] that $F_{\bar{\Pi}_0}^{s+1}(\bar{\mathscr{Q}}_0/S_0)$ is isomorphic to the étale double covering of S_0 obtained from the Stein factorization of the map $F^{s+1}(\bar{\mathscr{Q}}_0/S_0) \to S_0$, which, because of the isomorphism (2.15) and the observation made at the beginning of the proof, is isomorphic to $\widetilde{Y}_A^{\geq 2} \times_{\mathbf{P}(V_6)} S_0 \to S_0$.

2.5. Linear spaces on ordinary GM threefolds and fivefolds

In [DK19], we described the Hilbert schemes of linear spaces on a smooth GM variety X of dimension at least 3 in terms of its Lagrangian data set (V_6, V_5, A) , its EPW varieties, and their double covers. We focus here on the Hilbert schemes of lines on an ordinary GM threefold and σ -planes on an ordinary GM fivefold (see (2.2) for the definition). The description of [DK19] was given in terms of the first quadratic fibration

$$(2.16) \qquad \qquad \rho_1 \colon \mathbf{P}_X(\mathscr{U}_X) \longrightarrow \mathbf{P}(V_5),$$

where \mathscr{U}_X is the pullback to X of the tautological rank-2 vector bundle \mathscr{U} on $Gr(2, V_5)$ and the map ρ_1 is the pullback along the embedding $X \hookrightarrow Gr(2, V_5)$ of the projection

$$\tilde{\rho}_1 \colon \mathbf{P}_{\mathsf{Gr}(2,V_5)}(\mathscr{U}) \simeq \mathsf{Fl}(1,2;V_5) \longrightarrow \mathbf{P}(V_5).$$

For each $[v] \in \mathbf{P}(V_5)$, we have

$$\tilde{\rho}_1^{-1}([v]) = \mathbf{P}(v \wedge V_5) \simeq \mathbf{P}^3,$$

so that the fiber $\rho_1^{-1}([v])$ is a subscheme of $\mathbf{P}(v \wedge V_5)$.

The Hilbert scheme $F_1(X)$ of lines on X was identified in [DK19, Proposition 4.1] with the relative Hilbert scheme of lines of the map ρ_1 and the Hilbert scheme $F_{\sigma}^2(X)$ of σ -planes on X with the relative Hilbert schemes of planes of the map ρ_1 . This defines maps

(2.17)
$$\sigma: F_1(X) \to \mathbf{P}(V_5)$$
 and $\sigma: F_{\sigma}^2(X) \to \mathbf{P}(V_5).$

To better describe these maps, we set (see (2.13) and (2.4) for the notation)

$$\widetilde{Y}_{A,V_5}^{\geq 2} := \pi_A^{-1}(Y_{A,V_5}^{\geq 2}) \subset \widetilde{Y}_A^{\geq 2}.$$

2.5.1. σ -planes on ordinary GM fivefolds.— Let $X = \text{Gr}(2, V_5) \cap Q$ be an ordinary GM fivefold with Lagrangian data set (V_6, V_5, A) . By (2.8), we have $[V_5] \in \mathbf{P}(V_6^{\vee}) \setminus Y_{A^{\perp}}$. By [DK18, Proposition 4.5], the fibers of the first quadratic fibration ρ_1 defined in (2.16) are

(2.18)
$$\rho_{1}^{-1}([v]) \text{ is } \begin{cases} \text{a smooth quadric in } \mathbf{P}(v \wedge V_{5}) & \text{if } [v] \in \mathbf{P}(V_{5}) \setminus Y_{A,V_{5}}^{\geq 1}, \\ \text{a quadric of corank 1 in } \mathbf{P}(v \wedge V_{5}) & \text{if } [v] \in Y_{A,V_{5}}^{1}, \\ \text{the union of two planes in } \mathbf{P}(v \wedge V_{5}) & \text{if } [v] \in Y_{A,V_{5}}^{2}, \\ \text{a double plane in } \mathbf{P}(v \wedge V_{5}) & \text{if } [v] \in Y_{A,V_{5}}^{3}. \end{cases}$$

Using this, we proved in [DK19, Theorem 4.3(b)] and [DK20a, Corollary 5.5] that there is an isomorphism (2.19) $\tilde{\sigma} \colon F_{\sigma}^2(X) \xrightarrow{\sim} \widetilde{Y}_{A,V_5}^{\geq 2}$

such that $\pi_A \circ \tilde{\sigma} \colon F^2_{\sigma}(X) \to Y^{\geq 2}_{A,V_5}$ is the second map σ from (2.17). This has the following simple consequence (compare with [Nag98, Lemma 2.2]).

Lemma 2.10. Let $A \subset \bigwedge^3 V_6$ be a Lagrangian subspace with no decomposable vectors. For a general GM fivefold X such that A(X) = A, the Hilbert scheme $F_{\sigma}^2(X)$ of σ -planes contained in X is a smooth connected curve of genus 161.

Proof. According to the description of the moduli space in (2.8), a general GM fivefold X such that A(X) = A corresponds to a general point $[V_5] \in \mathbf{P}(V_6^{\vee}) \setminus Y_{A^{\perp}}$ and for such a $[V_5]$, the finite scheme $Y_{A,V_5}^{\geq 3}$ is empty, the curve $Y_{A,V_5}^{\geq 2}$ is smooth by Bertini's theorem, hence so is its étale (because $Y_{A,V_5}^{\geq 3} = \emptyset$) double cover $F_{\sigma}^2(X) \simeq \widetilde{Y}_{A,V_5}^{\geq 2}$. Since $K_{Y_A^{\geq 2}}$ is numerically equivalent to 3*H*, where *H* is the hyperplane class (see (2.5) or [O'Gr08a, (4.0.32)]), its genus is

(2.20)
$$g(Y_{A,V_5}^{\geq 2}) = 1 + \frac{1}{2}(K_{Y_A^{\geq 2}} \cdot H + H^2) = 1 + 2H^2 = 1 + 2\deg(Y_A^{\geq 2}) = 81.$$

The smooth curve $\widetilde{Y}_{A,V_5}^{\geq 2}$ is ample on the integral surface \widetilde{Y}_A^2 , hence connected. Its genus is therefore $2g(Y_{A,V_5}^{\geq 2}) - 1 = 161$.

2.5.2. Lines on ordinary GM threefolds.— Let $X = Gr(2, V_5) \cap \mathbf{P}(W) \cap Q$ be a smooth ordinary GM threefold, where $W \subset \bigwedge^2 V_5$ has codimension 2. By (2.8), we have

$$[V_5] \in Y_{A^\perp}^2$$

and the line $\mathbf{P}(W^{\perp}) \subset \mathbf{P}(\bigwedge^2 V_5^{\vee})$ is a pencil of skew-symmetric forms on V_5 . Since X is smooth, these forms all have one-dimensional kernels ([DK18, Remark 2.25]) and these kernels form a smooth conic

(2.21)
$$\Sigma_1(X) = F_{\sigma}^2(M_X) \subset \mathbf{P}(V_5)$$

(see also [DIM12, Section 3.2]). By [DK18, Proposition 4.5], we have $Y_{A,V_5}^3 \subset \Sigma_1(X) \subset Y_{A,V_5}$ and the fibers of the first quadratic fibration ρ_1 defined in (2.16) are

 $(2.22) \qquad \qquad \rho_1^{-1}([v]) = \begin{cases} \text{two reduced points in } \mathbf{P}(v \wedge V_5), & \text{if } [v] \in \mathbf{P}(V_5) \smallsetminus Y_{A,V_5}^{\geq 1}, \\ \text{a double point in } \mathbf{P}(v \wedge V_5), & \text{if } [v] \in Y_{A,V_5}^1 \smallsetminus \Sigma_1(X), \\ \text{the line } \mathbf{P}(W \cap (v \wedge V_5)), & \text{if } [v] \in Y_{A,V_5}^2 \smallsetminus \Sigma_1(X), \\ \text{a smooth conic in } \mathbf{P}(v \wedge V_5), & \text{if } [v] \in \Sigma_1(X) \cap Y_{A,V_5}^1, \\ \text{the union of two lines in } \mathbf{P}(v \wedge V_5), & \text{if } [v] \in \Sigma_1(X) \cap Y_{A,V_5}^2, \\ \text{a double line in } \mathbf{P}(v \wedge V_5), & \text{if } [v] \in Y_{A,V_5}^3. \end{cases}$

Using this, we proved the following result.

Proposition 2.11 ([DK19, Theorem 4.7]). Let X be a smooth ordinary GM threefold. The morphism defined in (2.17) factors through

(2.23)
$$\sigma: F_1(X) \longrightarrow Y_{A,V_5}^{\geq 2}.$$

It is an isomorphism over $Y_{A,V_5}^{\geq 2} \setminus \Sigma_1(X)$ and a double cover over the points of $Y_{A,V_5}^{\geq 2} \cap \Sigma_1(X)$, which is branched over $Y_{A,V_5}^3 \cap \Sigma_1(X)$.

In addition, elementary deformation theory implies that $F_1(X)$ has pure dimension 1, and local embedding dimension 2 at every singular point ([IP99, Proposition 4.2.2], [KPS18, Lemma 2.2.3]).

Lemma 2.12. Let $A \subset \bigwedge^3 V_6$ be a Lagrangian subspace with no decomposable vectors. For a general GM threefold X such that A(X) = A, the curve $Y_{A,V_5}^{\geq 2}$ has arithmetic genus 81 and the intersection $\Sigma_1(X) \cap Y_{A,V_5}^{\geq 2}$ is a finite scheme of length 10 contained in Sing $(Y_{A,V_5}^{\geq 2})$.

If also A is general, the curve $F_1(X)$ is a smooth irreducible curve of genus 71, the map $\sigma: F_1(X) \to Y_{A,V_5}^{\geq 2}$ is the normalization morphism, and $\operatorname{Sing}(Y_{A,V_5}^{\geq 2}) = \Sigma_1(X) \cap Y_{A,V_5}^{\geq 2}$.

Proof. For general $[V_5] \in Y_{A^{\perp}}^{\geq 2}$, the conic $\Sigma_1(X)$ is not contained in $Y_A^{\geq 2}$, hence not in $Y_{A,V_5}^{\geq 2}$. Indeed, $\Sigma_1(X)$ can be identified with the fiber of the partial desingularization $q: \widehat{Y}_A \to Y_{A^{\perp}}$ ([DK18, Section B.2]) over the point $[V_5] \in Y_{A^{\perp}}^2$. If a general fiber is contained in the exceptional divisor $E = p^{-1}(Y_A^{\geq 2})$ of the contraction $p: \widehat{Y}_A \to Y_A$, the exceptional divisor $E' = q^{-1}(Y_{A^{\perp}}^{\geq 2})$ of q is contained in E, hence E' = E since E is irreducible; but this was shown to be false in the proof of [DK18, Lemma B.5].

The equivalence $E \equiv 24H' - 5E'$ established in the proof of [DK18, Lemma B.5] implies that the length of the finite scheme $\Sigma_1(X) \cap Y_{A,V_5}^{\geq 2}$, which is the intersection number in \widehat{Y}_A of the curve $\Sigma_1(X)$, viewed as above as a fiber of q, with the divisor E, is

$$\Sigma_1(X) \cdot (24H' - 5E') = -5\Sigma_1(X) \cdot E'.$$

Since $Y_{A^{\perp}}$ has ordinary double points along $Y_{A^{\perp}}^2$, the exceptional divisor E' of the blow up of $Y_{A^{\perp}}^2$ has intersection -2 with the fibers, hence the length is $\Sigma_1(X) \cdot E = 10$. Furthermore, the scheme $\Sigma_1(X) \cap Y_{A,V_5}^{\geq 2}$ is contained in Sing $(Y_{A,V_5}^{\geq 2})$, because the finite birational map (2.23) is 2:1 over $\Sigma_1(X)$. The arithmetic genus of $Y_{A,V_5}^{\geq 2}$ was computed in (2.20).

When also A is general, X is a general GM threefold, hence $F_1(X)$ is a smooth irreducible curve of genus 71 ([Mar81, Proposition 6.4], [IP99, Theorem 4.2.7]), so that $Y_{A,V_5}^{\geq 2}$ is an integral curve which is smooth away from $\Sigma_1(X) \cap Y_{A,V_5}^{\geq 2}$ and $F_1(X)$ is its normalization.

Lemma 2.13. For any smooth GM threefold X, the curve $F_1(X)$ is connected.

Proof. Consider a general deformation $\varpi \colon \mathscr{X} \to B$ with central fiber X, parameterized by a smooth irreducible curve B. For any line $L \subset X$, there is an exact sequence

$$0 \to N_{L/X} \to N_{L/\mathcal{X}} \to \mathcal{O}_L \to 0.$$

It follows that $\chi(L, N_{L/\mathscr{X}}) = \chi(L, N_{L/X}) + 1 = 2$ hence, by deformation theory, every component of the Hilbert scheme $F_1(\mathscr{X})$ of lines contained in \mathscr{X} has dimension at least 2 at the point [L]. Since $F_1(X)$ has pure dimension 1 at [L], every component of $F_1(\mathscr{X})$ passing through [L] dominates B. Deformations of L in \mathscr{X} are contained in the fibers of ϖ , hence every irreducible component of $F_1(\mathscr{X}/B)$ dominates B. Since the general fiber of $F_1(\mathscr{X}/B) \to B$ is a smooth irreducible curve (Lemma 2.12), Stein factorization implies that every fiber is connected.

3. Topological preliminaries

In [Tju72, Section 4.3], Tyurin gave a beautiful argument (which he attributed to Clemens) proving the surjectivity of the Abel–Jacobi map given by the universal line on a threefold. In this section, we recall this argument and prove the generalization on which our results about GM threefolds and fivefolds are based. In Section 3.4, we use Picard–Lefschetz theory to show that intermediate Jacobians of very general GM varieties of dimensions 3 or 5 have trivial endomorphism rings.

3.1. Abel-Jacobi maps

We start by recalling a few properties of Abel-Jacobi maps.

Let X and Y be smooth proper varieties of respective dimensions d_X and d_Y , let Z be an algebraic cycle of dimension $d_Y + c$ on $X \times Y$, and let k be an integer. The Abel-Jacobi map

$$AJ_Z: H_k(Y, \mathbb{Z}) \longrightarrow H_{k+2c}(X, \mathbb{Z})$$

is defined as the composition

$$H_{k}(Y, \mathbb{Z}) \xrightarrow{\sim} H^{2d_{Y}-k}(Y, \mathbb{Z}) \xrightarrow{p_{Y}^{*}} H^{2d_{Y}-k}(X \times Y, \mathbb{Z})$$

$$\xrightarrow{\cdot \sim [Z]} H^{2d_{Y}+2d_{X}-2c-k}(X \times Y, \mathbb{Z}) \xrightarrow{\sim} H_{k+2c}(X \times Y, \mathbb{Z}) \xrightarrow{p_{X^{*}}} H_{k+2c}(X, \mathbb{Z}),$$

where p_X and p_Y are the projections from $X \times Y$ onto the factors and the isomorphisms are given by Poincaré duality and the middle map is the cup-product with the cohomology class of the cycle Z.

We will use the following functoriality properties of the Abel-Jacobi map.

Lemma 3.1. Let $i: X \to X'$ and $j: Y' \to Y$ be morphisms of smooth proper varieties.

- (a) We have $AJ_Z \circ j_* = AJ_{(Id_X \times j)^*(Z)}$ and $i_* \circ AJ_Z = AJ_{(i \times Id_Y)_*(Z)}$.
- (b) If Z' is a cycle on $X' \times Y$, one has $AJ_{(i \times Id_Y)^*(Z')} = i^* \circ AJ_{Z'}$.
- (c) If Z' is a cycle on $X \times Y'$, one has $AJ_{(Id_X \times j)_*(Z')} = AJ_{Z'} \circ j^*$.

Proof. All these statements follow from base change and the projection formula.

3.2. Generalized blow up decomposition

We will need the following (co)homological result generalizing the formula for the (co)homology of a smooth blow up (see [BFM19, Proposition 46] for another proof).

Lemma 3.2. Let S be a smooth proper variety and let E be a rank-r vector bundle on S. Let us consider $s \in H^0(S, E^{\vee}) \simeq H^0(\mathbf{P}_S(E), \mathcal{O}(1))$. We denote by $Z \subset S$ the zero-locus of s considered as a section of E^{\vee} and by $\widetilde{S} \subset \mathbf{P}_S(E)$ the zero-locus of s considered as a section of $\mathcal{O}(1)$.

Then \tilde{S} is a smooth hypersurface in $\mathbf{P}_{S}(E)$ if and only if Z is smooth of pure codimension r in S; in this case, there are direct sum decompositions

$$H^{\bullet}(\widetilde{S}, \mathbb{Z}) = H^{\bullet}(S, \mathbb{Z}) \oplus H^{\bullet-2}(S, \mathbb{Z}) \oplus \dots \oplus H^{\bullet-2(r-2)}(S, \mathbb{Z}) \oplus H^{\bullet-2(r-1)}(\mathbb{Z}, \mathbb{Z})$$

and

$$H_{\bullet}(\widetilde{S}, \mathbb{Z}) = H_{\bullet-2(r-1)}(Z, \mathbb{Z}) \oplus H_{\bullet-2(r-2)}(S, \mathbb{Z}) \oplus \cdots \oplus H_{\bullet-2}(S, \mathbb{Z}) \oplus H_{\bullet}(S, \mathbb{Z}).$$

Proof. We have a commutative diagram

$$\mathbf{P}_{Z}(E) \stackrel{\iota}{\longrightarrow} \widetilde{S} \stackrel{\iota}{\longleftrightarrow} \mathbf{P}_{S}(E)$$

$$p \downarrow \qquad \qquad \downarrow \pi$$

$$Z \stackrel{J}{\longleftrightarrow} S,$$

where π is a \mathbf{P}^{r-2} -fibration away from Z and a \mathbf{P}^{r-1} -fibration over Z. In particular, \widetilde{S} is a smooth hypersurface over $S \setminus Z$. Therefore, for the first statement, we have to check that \widetilde{S} is smooth of codimension 1 at $(x, e) \in \mathbf{P}_Z(E) \subset \widetilde{S}$ for all $e \in \mathbf{P}(E_x)$ if and only if Z is smooth of codimension r at x. There is a commutative diagram

$$0 \longrightarrow \mathbf{C} \xrightarrow{e} E_x \longrightarrow T_{\mathbf{P}_S(E),(x,e)} \longrightarrow T_{S,x} \longrightarrow 0$$

where ds is the differential of s considered as a section of $\mathscr{O}(1)$. The restriction of the map ds to E_x is zero, hence it factors through the dashed arrow, which can be identified with the differential of s considered as a section of E^{\vee} evaluated at $e \in E_x$. Thus, the vertical arrow is surjective at (x, e) for any $e \in E_x \setminus \{0\}$ if and only if $ds: T_{S,x} \to E_x^{\vee}$ is surjective, that is, if and only if Z is smooth of codimension r at x. Denote by $H \in H^2(\widetilde{S}, \mathbb{Z})$ the restriction to \widetilde{S} of the relative hyperplane class of $\mathbb{P}_S(E)$. Then $\mathbb{P}_Z(E)$ has codimension r-1 in \widetilde{S} and

$$c_{r-1}(N_{\mathbf{P}_{Z}(E)/\widetilde{S}}) = (-1)^{r-1} \Big(\iota^{*}(H)^{r-1} + p^{*} \jmath^{*} c_{1}(E) \smile \iota^{*}(H)^{r-2} + \dots + p^{*} \jmath^{*} c_{r-1}(E) \Big).$$

Indeed, this follows from the standard exact sequence

$$0 \to N_{\mathbf{P}_{Z}(E)/\widetilde{S}} \to N_{\mathbf{P}_{Z}(E)/\mathbf{P}_{S}(E)} \to \mathscr{O}(1)|_{\mathbf{P}_{Z}(E)} \to 0$$

in view of the isomorphism $N_{\mathbf{P}_Z(E)/\mathbf{P}_S(E)} \simeq p^*(N_{Z/S}) \simeq p^*j^*(E^{\vee})$ and the Whitney formula. In particular,

(3.1) $p_* c_{r-1}(N_{\mathbf{P}_Z(E)/\widetilde{S}}) = (-1)^{r-1}.$

We will now prove the direct sum decomposition of cohomology; the homological decomposition is proved analogously or follows from Poincaré duality. Consider the maps

$$\phi_k \colon H^{\bullet}(S, \mathbb{Z}) \longrightarrow H^{\bullet + 2k}(\overline{S}, \mathbb{Z})$$
$$\xi \longmapsto \pi^*(\xi) \smile H^k$$

and

$$\phi_Z \colon H^{\bullet}(Z, \mathbb{Z}) \longrightarrow H^{\bullet+2(r-1)}(\widetilde{S}, \mathbb{Z})$$
$$\xi \longmapsto \iota_*(p^*(\xi)).$$

We claim that

$$H^{\bullet}(\widetilde{S}, \mathbb{Z}) = \phi_0(H^{\bullet}(S, \mathbb{Z})) \oplus \phi_1(H^{\bullet-2}(S, \mathbb{Z})) \oplus \cdots \oplus \phi_{r-2}(H^{\bullet-2(r-2)}(S, \mathbb{Z})) \oplus \phi_Z(H^{\bullet-2(r-1)}(Z, \mathbb{Z})).$$

To prove that, we define maps

$$\psi_k \colon H^{\bullet}(\widetilde{S}, \mathbb{Z}) \longrightarrow H^{\bullet - 2k}(S, \mathbb{Z})$$
$$\eta \longmapsto \pi_*(\eta \smile H^{r-2-k})$$

and

$$\psi_Z \colon H^{\bullet}(\widetilde{S}, \mathbb{Z}) \longrightarrow H^{\bullet-2(r-1)}(\mathbb{Z}, \mathbb{Z})$$
$$\eta \longmapsto p_*(\iota^*(\eta)).$$

If $k \leq l$, we have

$$\psi_l(\phi_k(\xi)) = \pi_*(\pi^*(\xi) \smile H^{r-2-l+k}) = \xi \smile \pi_*(H^{r-2-l+k}) = \delta_{k,l}\xi$$

For $0 \le k \le r - 2$, we have

$$\psi_Z(\phi_k(\xi)) = p_*(i^*(\pi^*(\xi) \smile H^k)) = p_*(p^*(j^*(\xi)) \smile i^*(H)^k) = j^*(\xi) \smile p_*(i^*(H^k)) = 0.$$

Using (3.1), we obtain

$$\psi_{Z}(\phi_{Z}(\xi)) = p_{*}(\iota^{*}(\iota_{*}(p^{*}(\xi)))) = p_{*}(p^{*}(\xi) \smile c_{r-1}(N_{\mathbf{P}_{Z}(E)/\widetilde{S}})) = (-1)^{r-1}\xi$$

If we define maps

$$\psi := (\psi_0 \oplus \psi_1 \oplus \dots \oplus \psi_{r-2}) \oplus \psi_Z \quad \text{and} \quad \phi := (\phi_0 \oplus \phi_1 \oplus \dots \oplus \phi_{r-2}) \oplus \phi_Z$$

it follows that the map $\psi \circ \phi$ is lower triangular with ±1 on the diagonal, hence invertible, so that ϕ is injective and ψ is surjective.

The injectivity of ψ (hence the surjectivity of ϕ) follows from projective bundle formulas for the maps $\widetilde{S} \setminus \mathbf{P}_Z(E) \to S \setminus Z$ and $\mathbf{P}_Z(E) \to Z$, and excision.

3.3. The Clemens-Tyurin argument

The following result is a generalization of [Tju72, Section 4.3] (see also [Wel81, Lemma (4.6)]); the original result is the case m = 1.

Proposition 3.3. Let Y be a smooth projective variety of dimension 2m + 2 and let $X \subset Y$ be a smooth hyperplane section of Y. Let $F_Y \subset F^m(Y)$ be a smooth closed subscheme of the Hilbert scheme of m-dimensional linear projective spaces in Y, let $F_X \subset F_Y$ be the closed subscheme parameterizing projective spaces contained in X, and let $\mathscr{L}_Y \subset F_Y \times Y$ and $\mathscr{L}_X \subset F_X \times X$ be the pullbacks of the corresponding universal families of projective spaces. Assume that

- (a) F_Y is smooth;
- (b) \mathscr{L}_Y is dominant and generically finite over Y;
- (c) F_X is smooth of pure codimension m + 1 in F_Y ;
- (d) $H_{2m+1}(Y, \mathbb{Z}) = H_{2m+3}(Y, \mathbb{Z}) = 0.$

Then the Abel-Jacobi map

$$AJ_{\mathscr{L}_{Y}}: H_{1}(F_{X}, \mathbb{Z}) \longrightarrow H_{2m+1}(X, \mathbb{Z})$$

is surjective.

Proof. Consider the incidence diagram



Set $\widehat{X} := q^{-1}(X)$ and $\widehat{p} := p|_{\widehat{X}}, \ \widehat{q} := q|_{\widehat{X}}$, and consider the restricted diagram



Since $\mathscr{L}_Y \to F_Y$ is the projectivization of a rank-(m+1) vector bundle and $\widehat{X} \subset \mathscr{L}_Y$ is a relative hyperplane section, the hypotheses of Lemma 3.2 are satisfied by assumptions (a) and (c), hence \widehat{X} is smooth and there is a direct sum decomposition

$$H_{2m+1}(X, \mathbf{Z}) \simeq H_1(F_X, \mathbf{Z}) \oplus H_3(F_Y, \mathbf{Z}) \oplus \cdots \oplus H_{2m+1}(F_Y, \mathbf{Z}).$$

The Abel-Jacobi map $AJ_{\mathscr{L}_X}$ is the restriction of \hat{q}_* to the summand $H_1(F_X, \mathbb{Z})$. Let j be the inclusion $X \hookrightarrow Y$. By Lemma 3.1, the restriction of \hat{q}_* to the other summands $H_{2k+1}(F_Y, \mathbb{Z})$, for $k \in \{1, \ldots, m\}$, factors through the map $j^* \colon H_{2m+3}(Y, \mathbb{Z}) \to H_{2m+1}(X, \mathbb{Z})$, which vanishes by assumption (d). The surjectivity of $AJ_{\mathscr{L}_X}$ therefore will follow from the surjectivity of

$$\hat{q}_* \colon H_{2m+1}(X, \mathbb{Z}) \longrightarrow H_{2m+1}(X, \mathbb{Z}).$$

To prove this surjectivity, we note that by assumption (b), the map \hat{q} is dominant and generically finite hence, by [BM79, Lemma 7.15], the image of \hat{q}_* contains the vanishing cycles, that is, the kernel of the map

$$j_*: H_{2m+1}(X, \mathbb{Z}) \longrightarrow H_{2m+1}(Y, \mathbb{Z}).$$

By (d), the target is zero, so this proves the surjectivity of \hat{q}_* and of $AJ_{\mathscr{L}_X}$.

3.4. Intermediate Jacobians and their endomorphisms

Let X be a smooth projective variety of dimension 2m-1. We consider the middle cohomology $H^{2m-1}(X, \mathbb{Z})$ with its natural Hodge structure of weight 2m-1. The complex torus

$$Jac(X) = H^{2m-1}(X, \mathbf{C})/(F^m H^{2m-1}(X, \mathbf{C}) + H^{2m-1}(X, \mathbf{Z})),$$

where $F^m H^{2m-1}(X, \mathbb{C}) \subset H^{2m-1}(X, \mathbb{C})$ is part of the Hodge filtration, is called the (Griffiths) intermediate Jacobian of X (see [BL99, Chapter 4]). Poincaré duality induces a hermitian form on $H^{2m-1}(X, \mathbb{C})$ which is not necessarily positive definite but defines (in the terminology of [BL99, Chapter 2]) a canonical nondegenerate line bundle on Jac(X), making it into a nondegenerate torus. If moreover

(3.2)
$$F^{m+1}H^{2m-1}(X, \mathbf{C}) = 0,$$

the hermitian form is positive definite, the line bundle is ample, and it defines a principal polarization on the abelian variety Jac(X).

More generally, a polarized rational Hodge structure of odd weight defines an isogeny class of complex tori which, under a vanishing assumption analogous to (3.2), becomes an isogeny class of abelian varieties.

Let now M be a smooth projective variety of dimension 2m. For a smooth hypersurface $j: X \hookrightarrow M$, we denote by

$$H^{2m-1}(X, \mathbf{Q})_{\operatorname{van}} := \operatorname{Ker}\left(\mathfrak{z}_* \colon H^{2m-1}(X, \mathbf{Q}) \to H^{2m+1}(M, \mathbf{Q})\right)$$

the vanishing cohomology. By [Voi03, Proposition 2.27], there is an orthogonal direct sum decomposition

(3.3)
$$H^{2m-1}(X, \mathbf{Q}) = H^{2m-1}(X, \mathbf{Q})_{\text{van}} \oplus j^* H^{2m-1}(M, \mathbf{Q}).$$

In particular, the Hodge structure $H^{2m-1}(X, \mathbf{Q})_{\text{van}}$ acquires a polarization from Poincaré duality on X and we denote by $\text{Jac}(X)_{\text{van}}$ the corresponding isogeny class of nondegenerate complex tori.

We will say that the endomorphism ring of a complex torus T is trivial if any endomorphism of T is the multiplication by an integer. If $T \neq 0$, this means that the endomorphism ring End(T) is isomorphic to \mathbb{Z} or, equivalently, that the rational endomorphism ring $\text{End}(T) \otimes \mathbb{Q}$ is isomorphic to \mathbb{Q} ; so we can extend this terminology to isogeny classes of complex tori.

If T is a nonzero and nondegenerate complex torus with trivial endomorphism ring, it is indecomposable with Picard number 1 ([BL99, Propositions 1.7.3 and 2.3.7]).

The next result is an old statement made by Severi and proved in [CvG92, Theorem (1.1)].

Proposition 3.4. Let M be a smooth projective variety of dimension 2m. If $j: X \hookrightarrow M$ is a very general hyperplane section, the endomorphism ring of $Jac(X)_{van}$ is trivial.

Proof. Assume first that there is a rational map $f: M \to \mathbf{P}^1$, which we resolve by blowing up a smooth codimension-2 subvariety to obtain a morphism $\tilde{f}: \widetilde{M} \to M \xrightarrow{f} \mathbf{P}^1$ with critical values $t_1, \ldots, t_r \in \mathbf{P}^1$, and that the strict transform of X is the fiber over $0 \in \mathbf{P}^1 \setminus \{t_1, \ldots, t_r\}$. Let $\tilde{j}: X \hookrightarrow \widetilde{M}$ be the embedding and let

$$\rho \colon \pi_1(\mathbf{P}^1 \setminus \{t_1, \dots, t_r\}, 0) \longrightarrow \operatorname{Sp}(H^{2m-1}(X, \mathbf{Q}))$$

be the monodromy representation.

Assume moreover that the only singularities of the fibers of \tilde{f} are nodes. As explained in [Voi03, Sections 2.2.2 and 2.3.1], one can then attach to each singular point of a fiber a (noncanonically defined) vanishing cycle in $H^{2m-1}(X, \mathbf{Q})_{\text{van}}$ and the vanishing cycles span the vector space $H^{2m-1}(X, \mathbf{Q})_{\text{van}}$ ([Voi03, Lemma 2.26]; this reference deals with the case where each singular fiber has a single node but the proofs extend to the general case).

Assume that $f: M \to \mathbf{P}^1$ is a Lefschetz pencil. For each $i \in \{1, ..., r\}$, the singular fiber X_{t_i} has a single node and there exists an element γ_i of $\pi_1(\mathbf{P}^1 \setminus \{t_1, ..., t_r\}, 0)$ that acts on $H^{2m-1}(X, \mathbf{Q})$, via the monodromy representation, as the transvection

$$T_{\delta_i}: x \longmapsto x - (x \cdot \delta_i) \delta_i,$$

where $\delta_i \in H^{2m-1}(X, \mathbf{Q})_{\text{van}}$ is a vanishing cycle ([Voi03, Theorem 3.16]). The main results of Picard–Lefschetz theory are:

- the vanishing cycles $\delta_1, \ldots, \delta_r$ are in the same orbit for the monodromy representation ([Voi03, Corollary 3.24]);
- the monodromy representation is absolutely irreducible ([Voi03, Theorem 3.27] or [PS83, Lemma 3.13]).

It follows that the monodromy is "big": the Zariski closure of its image is the full symplectic group $\operatorname{Sp}(H^{2m-1}(X, \mathbb{C})_{\operatorname{van}})$ ([PS03, Lemma 4]). As in the proof of [PS03, Theorem 17], for $t \in \mathbb{P}^1$ very general, any endomorphism of $\operatorname{Jac}(X_t)_{\operatorname{van}}$ intertwines every element of the monodromy group, hence every element of the symplectic group. It must therefore be a multiple of the identity: the endomorphism ring of $\operatorname{Jac}(X_t)_{\operatorname{van}}$ is trivial.

Corollary 3.5. Let M be a smooth projective variety of dimension 2m with $H^{2m-1}(M, \mathbf{Q}) = 0$ and let $j: X \hookrightarrow M$ be a very general hyperplane section. The endomorphism ring of Jac(X) is then trivial.

These results apply to intermediate Jacobians of GM threefolds and fivefolds.

Corollary 3.6. If X is a GM variety of dimension $n \in \{3, 5\}$ the intermediate Jacobian Jac(X) is a principally polarized abelian variety of dimension 10. If, moreover, X is very general, we have

$$\operatorname{End}(\operatorname{Jac}(X)) \simeq \mathbb{Z}$$

In particular, the Picard number of Jac(X) is 1.

Proof. For any GM variety X of dimension 3 or 5, condition (3.2) holds by Proposition 2.6, hence the abelian variety Jac(X) is principally polarized; its dimension is 10 again by Proposition 2.6.

For the second statement, we may assume that X is ordinary; it is then a very ample divisor in its Grassmannian hull M_X , which is the Grassmannian $Gr(2, V_5)$ when n = 5 or the fixed smooth four-fold $Gr(2, V_5) \cap \mathbf{P}^7 \subset \mathbf{P}(\bigwedge^2 V_5)$ when n = 3. We have $H^n(M_X, \mathbf{Q}) = 0$ in both cases by Proposition 2.6, so Corollary 3.5 implies the first claim. A standard result then implies that the Picard number of Jac(X) is 1. \Box

4. Intermediate Jacobians of GM threefolds

In this section, we study the intermediate Jacobians of GM threefolds. The main result (Theorem 4.4) is stated at the end of Section 4.1 and its proof takes up the rest of Section 4.

4.1. Family of curves

Let X be an arbitrary smooth GM threefold. Its associated Lagrangian subspace $A \subset \bigwedge^3 V_6$ has no decomposable vectors (Section 2.3). Let $Y_A^{\geq 2} \subset \mathbf{P}(V_6)$ be the corresponding EPW surface and let

$$\pi_A \colon \widetilde{Y}_A^{\geq 2} \to Y_A^{\geq 2}$$

be the double covering from (2.4), which is connected and étale away from the finite set Y_A^3 .

We are going to construct a subvariety

$$Z \subset X \times \widetilde{Y}_A^{\geq 2}$$

such that the map $Z \to \widetilde{Y}_A^{\geq 2}$ is, away from a finite subset of $\widetilde{Y}_A^{\geq 2}$, a family of quintic curves of arithmetic genus 1 containing a fixed line $L_0 \subset X$. We will then check that the associated Abel-Jacobi map gives an isomorphism between $Alb(\widetilde{Y}_A^{\geq 2})$ and Jac(X).

We start by choosing a line $L_0 \subset X$. It is for the moment arbitrary, but we will impose some restrictions in Section 4.2. We consider the open surface $S_0 \subset Y_A^{\geq 2}$ defined by (2.14) and the family of quadrics $\mathscr{Q}_0 \to S_0$ obtained by base change to S_0 of the family (2.11). We denote by $F^4(\mathscr{Q}_0/S_0) \to S_0$ the relative Hilbert scheme of linear 4-spaces in the fibers of $\mathscr{Q}_0 \to S_0$ and by $F_{L_0}^4(\mathscr{Q}_0/S_0) \to S_0$ the subscheme parameterizing those 4-spaces that contain the line L_0 . Applying Corollary 2.9, we obtain an isomorphism

(4.1)
$$F_{L_0}^4(\mathscr{Q}_0/S_0) \simeq \widetilde{S}_0 := \widetilde{Y}_A^2 \times_{Y_A^2} S_0$$

of schemes over S_0 . In particular, the canonical map $F_{L_0}^4(\mathscr{Q}_0/S_0) \to S_0$ can be identified with the double étale covering $\pi \colon \widetilde{S}_0 \to S_0$ induced by the double covering π_A . Note that \widetilde{S}_0 is a smooth surface. Let

$$\widetilde{\mathscr{Q}}_0 := \mathscr{Q}_0 \times_{S_0} \widetilde{S}_0$$

be the base change of the family of quadrics $\mathscr{Q}_0 \to S_0$ along π . We have a canonical map

$$\widetilde{S}_0 \to F_{L_0}^4(\mathscr{Q}_0/S_0) \times_{S_0} \widetilde{S}_0 \hookrightarrow F^4(\mathscr{Q}_0/S_0) \times_{S_0} \widetilde{S}_0 \simeq F^4(\widetilde{\mathscr{Q}}_0/\widetilde{S}_0),$$

where the first map is the product of the isomorphism (4.1) with the identity map. By construction, it is a section of the projection $F^4(\widetilde{\mathscr{Q}}_0/\widetilde{S}_0) \to \widetilde{S}_0$.

Let $\mathscr{P}^4 \subset \widetilde{\mathscr{Q}_0} \subset \mathbf{P}(W) \times \widetilde{S}_0$ be the pullback of the universal family of projective 4-spaces over $F^4(\widetilde{\mathscr{Q}_0}/\widetilde{S}_0)$ along the section $\widetilde{S}_0 \to F^4(\widetilde{\mathscr{Q}_0}/\widetilde{S}_0)$ constructed above. Set

where the Grassmannian hull $M_X \subset \mathbf{P}(W)$ was defined in (2.7) and the intersection is taken inside $\mathbf{P}(W) \times \widetilde{S}_0$.

Proposition 4.1. The map $Z_0 \to \widetilde{S}_0$ is a flat family of local complete intersection curves in X of degree 5 and arithmetic genus 1 containing L_0 . In particular, $Z_0 \subset X \times \widetilde{S}_0$.

Proof. Let $y \in \widetilde{S}_0$ and set $[v] := \pi(y) \in \mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$. The fiber of Z_0 over y is

$$Z_{0,y} = M_X \cap \mathscr{P}_y^4 = \mathsf{CGr}(2, V_5) \cap \mathscr{P}_y^4.$$

The cone $CGr(2, V_5) \subset \mathbf{P}(\mathbf{C} \oplus \bigwedge^2 V_5)$ has codimension 3 and degree 5. Therefore, $CGr(2, V_5) \cap \mathscr{P}_y^4$ has dimension at least 1 and degree at most 5 (and if the dimension is 1, the degree is 5). Furthermore, $\mathscr{P}_y^4 \subset Q_v$, hence

$$Z_{0,v} \subset M_X \cap Q_v = X$$

Since X contains no surfaces of degrees less than 10 ([DK19, Corollary 3.5]), $Z_{0,y}$ is a local complete intersection curve in X of degree 5. This also proves the inclusion $Z_0 \subset X \times \widetilde{S}_0$.

Since the curve $Z_{0,y}$ is a dimensionally transverse linear section of CGr(2, V_5), the resolution of Lemma 2.4 restricts on $\mathscr{P}_y^4 \simeq \mathbf{P}^4$ to a resolution

$$0 \to \mathscr{O}_{\mathscr{P}^4_y}(-5) \to \mathscr{O}_{\mathscr{P}^4_y}(-3)^{\oplus 5} \to \mathscr{O}_{\mathscr{P}^4_y}(-2)^{\oplus 5} \to \mathscr{O}_{\mathscr{P}^4_y} \to \mathscr{O}_{Z_{0,y}} \to 0.$$

It follows that $h^0(Z_{0,y}, \mathscr{O}_{Z_{0,y}}) = h^1(Z_{0,y}, \mathscr{O}_{Z_{0,y}}) = 1$, hence $Z_{0,y}$ is a connected curve of arithmetic genus 1; in particular, its Hilbert polynomial is $h_{Z_{0,y}}(t) = 5t$. Since the Hilbert polynomial does not depend on y, the family of curves Z_0 is flat over \widetilde{S}_0 . Finally, $L_0 \subset M_X$ and $L_0 \subset \mathscr{P}_v^4$ by construction, hence $L_0 \subset Z_{0,y}$.

We now extend the family of curves $Z_0 \to \widetilde{S}_0$ to a family defined over the entire surface $\widetilde{Y}_A^{\geq 2}$. We will need the following construction.

Definition 4.2. Let $\mathscr{Z} \subset \mathscr{X} \times \mathscr{S}$ be an \mathscr{S} -flat family of subschemes in a projective variety \mathscr{X} , let $\varphi \colon \mathscr{S} \to \operatorname{Hilb}(\mathscr{X})$ be the induced morphism, and let $\mathscr{S} \subset \overline{\mathscr{S}}$ be a partial compactification of \mathscr{S} . Then φ can be considered as a rational map $\overline{\mathscr{S}} \dashrightarrow \operatorname{Hilb}(\mathscr{X})$. Let $\widetilde{\mathscr{F}} \subset \overline{\mathscr{S}} \times \operatorname{Hilb}(\mathscr{X})$ be the graph of φ and let $\tilde{\varphi} \colon \widetilde{\mathscr{F}} \to \operatorname{Hilb}(\mathscr{X})$ be the projection. Let $\widetilde{\mathscr{Z}} \subset \mathscr{X} \times \widetilde{\mathscr{F}}$ be the pullback of the universal subscheme in $\mathscr{X} \times \operatorname{Hilb}(\mathscr{X})$ and let

$$\overline{\mathcal{Z}} \subset \mathscr{X} \times \overline{\mathscr{S}}$$

be the scheme-theoretic image of $\widetilde{\mathcal{Z}}$ by the morphism $\mathscr{X} \times \widetilde{\mathscr{F}} \to \mathscr{X} \times \overline{\mathscr{F}}$. We will call the subscheme $\overline{\mathscr{Z}}$ the Hilbert closure of \mathscr{Z} with respect to the embedding $\mathscr{S} \subset \overline{\mathscr{S}}$.

We apply this construction to the subscheme $Z_0 \subset X \times \widetilde{S}_0$ and the embedding $\widetilde{S}_0 \subset \widetilde{Y}_A^{\geq 2}$.

Lemma 4.3. Let

be the Hilbert closure of the subscheme $Z_0 \subset X \times \widetilde{S}_0$ with respect to the embedding $\widetilde{S}_0 \subset \widetilde{Y}_A^{\geq 2}$. Away from a finite subscheme of $\widetilde{Y}_A^{\geq 2}$, the scheme Z is a flat family of curves of degree 5 and arithmetic genus 1 containing the line L_0 . Moreover, we have

$$Z \times_{\widetilde{Y}_{4}^{\geq 2}} \widetilde{S}_{0} = Z_{0}$$

as subschemes of $X \times \widetilde{S}_0$.

Proof. Since the surface $\widetilde{Y}_A^{\geq 2}$ is normal, the rational map $\widetilde{Y}_A^{\geq 2} \longrightarrow \text{Hilb}(X)$ defined by the subscheme Z_0 extends regularly to all codimension-1 points of $\widetilde{Y}_A^{\geq 2}$. The nonflat locus of the morphism $Z \to \widetilde{Y}_A^{\geq 2}$ is therefore supported in codimension 2, hence is a finite subscheme. All the remaining properties of Z are clear from the construction of the Hilbert closure.

By Proposition 4.1, every irreducible component of Z_0 has dimension 3. By definition of the Hilbert closure, the same is true for Z.

The main result of this section is the following (recall that by Propositions 2.5 and 2.6, the abelian groups $H_1(\widetilde{Y}_{A(X)}^{\geq 2}, \mathbb{Z})$ and $H_3(X, \mathbb{Z})$ are both free of rank 20).

Theorem 4.4. For any Lagrangian subspace $A \subset \bigwedge^3 V_6$ such that A has no decomposable vectors and $Y_A^{\geq 3} = \emptyset$, the abelian variety $Alb(\widetilde{Y}_A^{\geq 2})$ has a canonical principal polarization. If moreover $Y_{A^{\perp}}^{\geq 3} = \emptyset$, there is an isomorphism

(4.4)
$$\operatorname{Alb}(\widetilde{Y}_{A}^{\geq 2}) \simeq \operatorname{Alb}(\widetilde{Y}_{A\perp}^{\geq 2})$$

of principally polarized abelian varieties.

Furthermore, if X is any smooth GM threefold with associated Lagrangian subspace A such that $Y_A^{\geq 3} = \emptyset$, if $L_0 \subset X$ is any line and if $Z \subset X \times \widetilde{Y}_A^{\geq 2}$ is the subscheme defined in Lemma 4.3, the Abel-Jacobi map

(4.5)
$$\operatorname{AJ}_Z \colon H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z}) \longrightarrow H_3(X, \mathbb{Z})$$

is an isomorphism of integral Hodge structures which induces an isomorphism

(4.6)
$$\operatorname{Alb}(\widetilde{Y}_A^{\geq 2}) \xrightarrow{\sim} \operatorname{Jac}(X)$$

of principally polarized abelian varieties.

This theorem is a more precise form of Theorem 1.1 for threefolds and its proof takes up the rest of Section 4. Note that if A is very general, the principal polarization of $Alb(\widetilde{Y}_A^{\geq 2})$ is unique by Corollary 3.6.

4.2. The boundary of the family

To prove Theorem 4.4, we study, over the boundary $\widetilde{Y}_A^{\geq 2} \setminus \widetilde{S}_0$, the family of curves Z constructed in Lemma 4.3. By (2.14), this boundary consists of the curve $\widetilde{Y}_{A,V_5}^{\geq 2}$ and the finite set Y_A^3 . As we will see in the proof of Proposition 4.15, finite sets are not important for the Abel–Jacobi map, so we will concentrate on a dense open subset (denoted by S_{0,V_5} and defined in Definition 4.7) of the curve $\widetilde{Y}_{A,V_5}^{\geq 2}$. We will construct a

diagram



where $S_{0+} = S_0 \cup S_{0,V_5}$ and all squares are cartesian. The lower vertical arrows are double coverings (étale except for the right one, which is only étale away from $Y_A^{\geq 3}$). We want to emphasize that the schemes Z_{0+} and Z are *different* over the boundary $\widetilde{S}_{0+} \times \widetilde{S}_0 \subset \widetilde{Y}_A^{\geq 2} \times \widetilde{S}_0$, and this difference will be crucial for the rest of the proof. In fact, the map $Z'_{0,V_5} \to S'_{0,V_5}$ is a flat family of surfaces in M_X , while the map $Z \to \widetilde{Y}_A^{\geq 2}$ is a family of curves in X.

To construct the diagram, we need to impose some restrictions on X and L_0 . First, we will assume from now on that X is ordinary. To explain the restriction imposed on L_0 , we will need the following definition (the map $\sigma: F_1(X) \to Y_{A,V_5}^{\geq 2}$ and the conic $\Sigma_1(X) \subset \mathbf{P}(V_5)$ were defined in (2.23) and (2.21)).

Definition 4.5. A line *L* on *X* is nice if $\sigma([L]) \notin \Sigma_1(X)$.

We will use the following simple observation.

Lemma 4.6. If $L \subset X$ is a nice line and $[v] := \sigma([L])$, one has

$$\mathbf{P}(W) \cap \mathbf{P}(v \wedge V_5) = L$$

In particular, the subspace $\mathbf{P}(W)$ is transverse to $\mathbf{P}(v \wedge V_5) \subset \mathbf{P}(\wedge^2 V_5)$.

Proof. By definition of a nice line, we have $[v] \in Y^2_{A,V_5} \setminus \Sigma_1(X)$, hence (2.22) applies.

From now on, we will assume that L_0 is a nice line and set

$$[v_0] := \sigma([L_0]) \in \mathbf{P}(V_5).$$

Recall that $Y_{A,V_5}^{\geq 2} \cap \Sigma_1(X)$ is a finite scheme (Lemma 2.12).

Definition 4.7. Denote by S_{0,V_5} the dense open complement in the curve $Y_{A,V_5}^{\geq 2}$ of the finite set $Y_{A,V_5}^{\geq 2} \cap \Sigma_1(X)$ and of the finite subset of $Y_{A,V_5}^{\geq 2}$ corresponding to lines intersecting L_0 (including the line L_0 itself). Set

$$S_{0+} := Y_A^{\geq 2} \setminus (Y_{A,V_5}^{\geq 2} \setminus S_{0,V_5}) = S_0 \cup S_{0,V_5} \subset Y_A^{\geq 2}.$$

This is a smooth open subscheme of $Y_A^{\geq 2}$ containing S_0 and with finite complement.

Note that each point of the curve S_{0,V_5} corresponds to a nice line on X.

Lemma 4.8. The double covering $\pi_A : \widetilde{Y}_A^{\geq 2} \to Y_A^{\geq 2}$ splits over the curve $Y_{A,V_5}^{\geq 2}$.

Proof. As we saw in the proof of Corollary 2.9, the double covering $\pi: \widetilde{S}_0 \to S_0$ induced by π_A agrees with the relative Hilbert scheme $F_{\overline{L}_0}^2(\overline{\mathscr{Q}}_0/S_0) \to S_0$ of planes containing the projection \overline{L}_0 of the line L_0 , where the family of quadrics $\overline{\mathscr{Q}}_0 \to S_0$ is obtained from the family (2.11) by restricting to S_0 and passing to the quotients with respect to the 2-dimensional kernel spaces of quadrics. We will prove that this identification also holds over S_{0+} .

Denote by $\mathscr{Q}_{0+} \to S_{0+}$ the restriction of the family of quadrics (2.11) to S_{0+} . For $[v] \in S_{0,V_5}$, the quadric Q_v is the restriction to $\mathbf{P}(W)$ of the corresponding Plücker quadric (see (2.1)), that is

$$Q_v = \mathbf{P}(W) \cap \operatorname{Cone}_{\mathbf{P}(v \wedge V_5)}(\operatorname{Gr}(2, V_5/\mathbf{C}v)).$$

By Definition 4.7, the line L_v corresponding to the point $[v] \in S_{0,V_5} \subset Y_{A,V_5}^{\geq 2}$ is nice hence, by Lemma 4.6, the space $\mathbf{P}(W)$ intersects $\mathbf{P}(v \wedge V_5)$ transversely along the line L_v , so that

$$(4.8) Q_v = \operatorname{Cone}_{L_v}(\operatorname{Gr}(2, V_5/\mathbf{C}v)).$$

In particular, Q_v has corank 2 and its vertex L_v does not meet L_0 (by Definition 4.7). Therefore, by passing to the quotients with respect to the kernel spaces of quadrics, we obtain, as in the proof of Corollary 2.9, a family $\bar{\mathscr{Q}}_{0+} \to S_{0+}$ of nondegenerate 4-dimensional quadrics over S_{0+} and conclude that the Hilbert scheme $F_{\bar{L}_0}^2(\bar{\mathscr{Q}}_{0+}/S_{0+})$ of planes in its fibers containing \bar{L}_0 is isomorphic to $F_{L_0}^4(\mathscr{Q}_{0+}/S_{0+})$ and gives an étale double covering of S_{0+} . Over the dense open subset $S_0 \subset S_{0+}$, this covering is induced by π_A , hence the same is true over S_{0+} , that is,

(4.9)
$$F_{L_0}^4(\mathscr{Q}_{0+}/S_{0+}) \simeq F_{\bar{L}_0}^2(\bar{\mathscr{Q}}_{0+}/S_{0+}) \simeq \widetilde{S}_{0+} := \widetilde{Y}_A^2 \times_{Y_A^2} S_{0+}.$$

Therefore, to prove that the covering $\pi_A: \widetilde{Y}_A^{\geq 2} \to Y_A^{\geq 2}$ splits over $Y_{A,V_5}^{\geq 2}$, it is enough to check that the covering $F_{\tilde{L}_0}^2(\bar{\mathscr{Q}}_{0+}/S_{0+}) \to S_{0+}$ splits over S_{0,V_5} . We do that by constructing a section of this covering over S_{0,V_5} as follows: for $[v] \in S_{0,V_5}$, consider the plane

(4.10)
$$\mathbf{P}(v_0 \wedge (V_5/\mathbf{C}v)) \subset \operatorname{Gr}(2, V_5/\mathbf{C}v) = \bar{Q}_v$$

(note that $[v_0] \neq [v]$ for $[v] \in S_{0,V_5}$ by Definition 4.7). The line \overline{L}_0 is contained in this plane because $L_0 = \mathbf{P}(W) \cap \mathbf{P}(v_0 \wedge V_5)$ by Lemma 4.6. Therefore, we obtain a regular map

(4.11)
$$S_{0,V_5} \longrightarrow F^2_{\tilde{L}_0}(\bar{\mathscr{Q}}_0/S_0) = F^4_{L_0}(\mathscr{Q}_{0+}/S_{0+}) \simeq \widetilde{S}_{0+}$$

which gives the required section.

Remark 4.9. The map (4.11) is the restriction of the map

$$\mathbf{P}(V_5) \smallsetminus \{ [v_0] \} \longrightarrow F_{L_0}^4(\mathscr{Q}/\mathbf{P}(V_6))$$

$$[v] \longmapsto \mathbf{P} \Big(W \cap ((\mathbf{C}v_0 \oplus \mathbf{C}v) \land V_5) \Big)$$

hence it is well defined even if the curve $Y_{A,V_{\pi}}^{\geq 2}$ is not reduced.

We still denote by π the double covering $\widetilde{S}_{0+} \to S_{0+}$ constructed above. Note that \widetilde{S}_{0+} is a smooth irreducible open surface in $\widetilde{Y}_A^{\geq 2}$ containing \widetilde{S}_0 as an open subscheme. Let

$$\widetilde{\mathcal{Q}}_{0+} = \mathscr{Q}_{0+} \times_{S_{0+}} \widetilde{S}_{0+} \simeq \mathscr{Q} \times_{\mathbf{P}(V_6)} \widetilde{S}_{0+}$$

be the base change of the family $\mathscr{Q}_{0+} \to S_{0+}$ along π . The isomorphism (4.9) induces a section

$$\widetilde{S}_{0+} \longrightarrow F^4_{L_0}(\widetilde{\mathscr{Q}}_{0+}/\widetilde{S}_{0+})$$

of its relative Hilbert scheme of 4-spaces and we denote by

$$\mathscr{P}^4_+ \subset \widetilde{\mathscr{Q}}_{0+} \subset \mathbf{P}(W) \times \widetilde{S}_{0+}$$

the corresponding family of projective 4-spaces over \widetilde{S}_{0+} , which agrees by construction with the family \mathscr{P}^4 over $\widetilde{S}_0 \subset \widetilde{S}_{0+}$. We set

This defines the middle column of the diagram (4.7). We denote by $Z_{0+,y} = \mathscr{P}^4_{+y} \cap M_X \subset M_X$ the fiber of $Z_{0+,y}$ over a point $y \in \widetilde{S}_{0+}$.

Lemma 4.10. We have $Z_{0+} \times_{\widetilde{S}_{0+}} \widetilde{S}_0 = Z_0$ and, for general points y of every irreducible component of $\widetilde{S}_{0+} \times \widetilde{S}_0$, we have $Z_y \subset Z_{0+,y}$, where Z_y is the fiber of the scheme Z defined in (4.3).

Proof. The equality follows from the fact that the family of 4-spaces \mathscr{P}^4_+ agrees with \mathscr{P}^4 over \widetilde{S}_0 . Let y be a general point of an irreducible component of $\widetilde{S}_{0+} \setminus \widetilde{S}_0$. By continuity, we obtain $Z_y \subset \mathscr{P}^4_{+,y}$. Since $Z_y \subset X \subset M_X$, we also get $Z_y \subset Z_{0+,y}$.

We denote by

(4.13)

the image of the map (4.11) and by

$$Z'_{0,V_5} := Z_{0,+} \times_{\widetilde{S}_{0,+}} S'_{0,V_5} \subset M_X \times S'_{0,V_5}$$

 $S'_{0,V_5} \subset \widetilde{S}_{0+}$

the restriction of the family (4.12) to the curve $S'_{0,V_{\epsilon}}$.

Proposition 4.11. The map $Z'_{0,V_5} \to S'_{0,V_5}$ is a flat family of surfaces which are isomorphic to hyperplane sections of a cubic scroll $\mathbf{P}^1 \times \mathbf{P}^2$. For each point $y \in S_{0,V_5}$, the fiber Z'_y contains the corresponding nice line $L_{\pi(y)}$ and the line L_0 .

We will give a more detailed description of the fibers of Z'_{0,V_5} in Lemma 4.13.

Proof. Let $y \in \widetilde{S}'_{0,V_5}$ and set again $[v] = \pi(y) \in \mathbf{P}(V_5)$. The proof of Lemma 4.8 shows that the quadric Q_v has the form (4.8) and that the point $y \in S'_{0,V_5}$ corresponds to the plane (4.10) in \overline{Q}_v . Its preimage in Q_v is the 4-space

$$\mathscr{P}^4_{+v} = \mathbf{P}(((v \land V_5) \cap W) \oplus (v_0 \land (V_5/\mathbf{C}v))) = \mathbf{P}(V_2 \land V_5) \cap \mathbf{P}(W),$$

where $V_2 \subset V_5$ is the subspace spanned by v_0 and v (they are linearly independent by Definition 4.7). Furthermore, by Lemma 2.2, the fiber of Z'_{0,V_5} at y can be written as

(4.14)
$$Z'_{y} := M_{X} \cap \mathscr{P}^{4}_{+y} = \operatorname{Cone}_{\mathbf{P}(\wedge^{2}V_{2})} (\mathbf{P}(V_{2}) \times \mathbf{P}(V_{5}/V_{2})) \cap \mathbf{P}(W),$$

so it is a linear section of a cone over the 3-dimensional cubic scroll.

The vertex $[\bigwedge^2 V_2]$ of the cone does not belong to $\mathbf{P}(W)$. Indeed, since both L_0 and L_v are nice lines (in the sense of Definition 4.5), we have by Lemma 4.6

(4.15)
$$\mathbf{P}(W) \cap \mathbf{P}(v_0 \wedge V_5) = L_0 \quad \text{and} \quad \mathbf{P}(W) \cap \mathbf{P}(v \wedge V_5) = L_v.$$

But $[\wedge^2 V_2] \in \mathbf{P}(v_0 \wedge V_5) \cap \mathbf{P}(v \wedge V_5)$, so if it also belongs to $\mathbf{P}(W)$, we get $L_0 \cap L_v \neq \emptyset$, which contradicts Definition 4.7.

Since W has codimension 2 in $\bigwedge^2 V_5$ and is transverse to $V_2 \land V_5$ by Lemma 4.6, it follows that Z'_y is isomorphic to a hyperplane section of $\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2) \simeq \mathbf{P}^1 \times \mathbf{P}^2$. It is easy to see that its Hilbert polynomial is $h_{Z'_y}(t) = (t+1)(\frac{3}{2}t+1)$. Since it does not depend on y, the family of surfaces Z'_{0,V_5} is flat over S'_{0,V_5} .

A combination of (4.14) and (4.15) shows that $L_v, L_0 \subset Z'_v$.

$$Z'_{v} = \operatorname{Cone}_{[v_{0} \wedge v]} \big(\mathbf{P}(\mathbf{C}v_{0} \oplus \mathbf{C}v) \times \mathbf{P}(V_{5}/(\mathbf{C}v_{0} \oplus \mathbf{C}v)) \big) \cap \mathbf{P}(W)$$

over $\mathbf{P}(V_5) \setminus \{[v_0]\}$. When $[v] \in \mathbf{P}(V_5) \setminus \mathbf{P}(V_3)$, where $V_3 \subset V_5$ is defined by $L_0 = \mathbf{P}(v_0 \wedge V_3)$, these surfaces are hyperplane sections of the cubic scroll $\mathbf{P}^1 \times \mathbf{P}^2$. This proves flatness of the family Z'_{0,V_5} even when the curve S'_{0,V_5} is not reduced.

Propositions 4.1 and 4.11 show that the components of Z_0 and Z'_{0,V_5} all have dimension 3. They are components of the scheme Z_{0+} , which has other 3-dimensional components over the curve S''_{0,V_5} defined by $S''_{0,V_5} = \widetilde{S}_{0+} \setminus (\widetilde{S}_0 \cup S'_{0,V_5})$, but we will not need this fact.

Finally, to construct the left column of (4.7), recall that the curve S'_{0,V_5} is by definition isomorphic to the open subscheme $S_{0,V_5} \subset Y^{\geq 2}_{A,V_5} \setminus \Sigma_1(X)$, which via the map $\sigma : F_1(X) \to Y^{\geq 2}_{A,V_5}$ is identified with an open subscheme in the Hilbert scheme of lines $F_1(X)$ (see Proposition 2.11). Applying to Z'_{0,V_5} the construction of Hilbert closure from Definition 4.2, we obtain a subscheme

such that $Z'_F \times_{F_1(X)} S'_{0,V_5} \simeq Z'_{0,V_5}$. Note that Z'_F may be not flat over the singular locus of the curve $F_1(X)$.

4.3. A relation between the subschemes

Let X be a smooth ordinary GM threefold and let L_0 be a nice line on X. In (4.3) and (4.16), we constructed subschemes $Z \subset X \times \widetilde{Y}_A^{\geq 2}$ and $Z'_F \subset M_X \times F_1(X)$. The proof of Theorem 4.4 is based on a relation between the schemes

 $Z \cap (X \times S'_{0,V_5})$ and $Z'_F \cap (X \times S'_{0,V_5})$,

where the curve S'_{0,V_5} defined in (4.13) is considered as a subscheme of $\widetilde{Y}_A^{\geq 2}$ and $F_1(X)$.

To prove such a relation, we will assume that the Hilbert scheme of lines $F_1(X)$ is a smooth curve (by Lemma 2.13, it is then irreducible). This assumption implies that the open curves $Y^2_{A,V_5} \setminus \Sigma_1(X)$ and S'_{0,V_5} are also smooth and irreducible. The next lemma sharpens the results of Proposition 4.11 under this assumption.

Lemma 4.13. Assume that $F_1(X)$ is smooth. For a general point y in S'_{0,V_5} , the fiber Z'_y of $Z'_{0,V_5} \to S'_{0,V_5}$ is a smooth cubic surface scroll and the lines L_0 and $L_{\pi(y)}$ are distinct fibers of the ruling of this scroll.

Proof. We saw at the end of the proof of Proposition 4.11 that Z'_y is a hyperplane section of $\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2)$. These hyperplane sections come in two kinds:

- (a) smooth cubic scrolls with projection to \mathbf{P}^1 induced by $\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2) \rightarrow \mathbf{P}(V_2)$,
- (b) unions $(\mathbf{P}(V_2) \times \mathbf{P}(V'_2)) \cup (\{[v']\} \times \mathbf{P}(V_5/V_2))$, where $V'_2 \subset V_5/V_2$ and $[v'] \in \mathbf{P}(V_2)$.

In case (b), the σ -plane $\{[v']\} \times \mathbf{P}(V_5/V_2)$ is contained in $Z'_y \subset M_X$, hence $[v'] \in \Sigma_1(X)$ by (2.21) and $[v'] \neq [v_0]$. Since V_2 is the subspace of V_5 spanned by v_0 and a vector v such that $[v] = \pi(y)$, and since $[v'] \in \mathbf{P}(V_2) \setminus \{[v_0]\}$, case (b) holds only if

$$[v] \in \mathrm{pr}_{v_0}(\Sigma_1(X)) \cap \mathrm{pr}_{v_0}(Y_{A,V_5}^{\geq 2}) \subset \mathbf{P}(V_5/\mathbf{C}v_0),$$

where $\operatorname{pr}_{v_0} \colon \mathbf{P}(V_5) \to \mathbf{P}(V_5/\mathbf{C}v_0)$ is the projection from v_0 . Since $Y_{A,V_5}^{\geq 2}$ is an integral curve of degree 40, its image by pr_{v_0} is contained in the image of the conic $\Sigma_1(X)$ only if the line connecting v_0 with a general point of $\Sigma_1(X)$ intersects $Y_{A,V_5}^{\geq 2}$ in 20 points. But the surface $Y_A^{\geq 2}$ is an intersection of hypersurfaces of degree 6 by [DK19, (33)], hence the same is true for its hyperplane section $Y_{A,V_5}^{\geq 2}$, and the curve $Y_{A,V_5}^{\geq 2}$ would then contain the cone $\operatorname{Cone}_{[v_0]}(\Sigma_1(X))$, which is absurd. Therefore, for y general in \widetilde{S}'_{0,V_5} , we are in case (a).

By (4.15), the lines L_0 and L_v are contained in fibers of the map $\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2) \to \mathbf{P}(V_2)$. In case (a), they are therefore the fibers of the ruling of the scroll.

Since the curve S'_{0,V_5} is isomorphic to a dense open subscheme of the smooth curve $F_1(X)$, the locally closed embedding $S'_{0,V_5} \hookrightarrow \widetilde{Y}_A^{\geq 2}$ extends to a regular map

(4.17)
$$\phi: F_1(X) \longrightarrow \widetilde{Y}_A^{\geq 2}.$$

We combine all these maps into the commutative diagram

where $\mathscr{L}_1(X) \subset X \times F_1(X)$ is the universal family of lines, $i: X \hookrightarrow M_X$ is the embedding, and the schemes Z and Z'_F are defined by (4.3) and (4.16) respectively. We make the following key observation.

Proposition 4.14. Assume that $F_1(X)$ is smooth. There is a dense open subscheme $U \subset F_1(X)$ such that

(4.19)
$$Z'_F \cap (X \times U) = \left(\phi^{-1}(Z) \cap (X \times U)\right) + \left(\mathscr{L}_1(X) \cap (X \times U)\right)$$

as cycles.

Proof. Since we only need an equality over a dense open subset of $F_1(X)$, we may base change both sides along the open embedding $S'_{0,V_5} \hookrightarrow F_1(X)$. The left side can then be rewritten as

$$Z'_F \cap (X \times S'_{0,V_5}) = Z'_{0,V_5} \cap (X \times S'_{0,V_5}).$$

By Proposition 4.11 and Lemma 4.13, the morphism $Z'_{0,V_5} \to S'_{0,V_5}$ is a flat family of surfaces whose general fiber is a smooth cubic surface scroll. Since X contains no surfaces of degrees less than 10 ([DK19, Corollary 3.5]), it contains no components of any fiber of Z'_{0,V_5} . Therefore, the morphism

$$Z'_{0,V_5} \times_{M_X} X \to S'_{0,V_5}$$

is a flat family of curves whose fiber over a general point $y \in S'_{0,V_{5}}$ is the intersection

$$C_v := Z'_v \cap Q_0$$

of the smooth cubic surface scroll Z'_{y} with any non-Plücker quadric Q_0 containing X. Such an intersection is a connected curve of degree 6 and arithmetic genus 2. Since the lines L_0 and $L_{\pi(y)}$ are contained both in the scroll Z'_{y} and the quadric Q_0 , they are components of C_{y} .

To describe the remaining components, we denote by e the class of the exceptional section L_e of the scroll Z'_v and by f the class of a fiber of its ruling. We have

$$e^2 = -1$$
, $e \cdot f = 1$, $f^2 = 0$.

The hyperplane class is equal to e + 2f hence, the class of C_y in Z'_y is 2e + 4f. As we observed above, the lines L_0 and $L_{\pi(y)}$ are fibers of the ruling, hence

$$C_y' := C_y - L_0 - L_{\pi(y)}$$

is an effective divisor on Z'_v with class 2e + 2f.

If C'_{y} contains a line, the class of this line is either f (the class of a fiber of the ruling), or e (the class of the exceptional section L_{e} of Z'_{y}). If it is f, the residual components have class 2e + f, and since $(2e + f) \cdot e = -1$, the section L_{e} is in both cases a component of C'_{y} , hence a line on X. The line L_{e} is in the finite set of

lines on X intersecting L_0 , and $L_{\pi(y)}$ is in the finite set of lines on X intersecting a line that intersects L_0 . It follows that for y general, the curve C'_v contains no lines.

By Lemma 4.10, the curve Z_y is contained in the surface Z'_y for general $y \in S'_{0,V_5}$. Therefore, by Lemma 4.3, for general $y \in S'_{0,V_5}$, the sextic curve C_y contains the quintic curve Z_y , hence $C_y = Z_y + L'_y$, where L'_y is a line. Since Z_y contains L_0 and C'_y contains no lines, the line L'_y must be $L_{\pi(y)}$. Thus,

$$C_y = Z_y + L_{\pi(y)}$$

Since this holds for y general in S'_{0,V_5} , it follows that the equality of cycles (4.19) holds over a dense open subscheme $U \subset S'_{0,V_5} \subset F_1(X)$.

4.4. Abel-Jacobi maps

Let X be a smooth GM threefold with associated Lagrangian A. Assume that $Y_A^3 = \emptyset$, so that $\widetilde{Y}_A^{\geq 2}$ is a smooth surface, and that the Hilbert scheme of lines $F_1(X)$ is smooth. The subscheme $Z \subset X \times \widetilde{Y}_A^{\geq 2}$ was constructed in Lemma 4.3. Consider the universal line $\mathscr{L}_1(X) \subset X \times F_1(X)$, the Abel-Jacobi maps

$$\mathsf{AJ}_Z \colon H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z}) \to H_3(X, \mathbb{Z}) \quad \text{and} \quad \mathsf{AJ}_{\mathscr{L}_1(X)} \colon H_1(F_1(X), \mathbb{Z}) \to H_3(X, \mathbb{Z}),$$

and the map $\phi \colon F_1(X) \to \widetilde{Y}_A^{\geq 2}$ defined in (4.17).

Proposition 4.15. Let X be a smooth ordinary GM threefold with associated Lagrangian A satisfying $Y_A^3 = \emptyset$. Assume that $F_1(X)$ is smooth and let L_0 be a nice line on X. The composition of maps

$$H_1(F_1(X), \mathbb{Z}) \xrightarrow{\phi_*} H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z}) \xrightarrow{AJ_Z} H_3(X, \mathbb{Z})$$

is equal to the map $-AJ_{\mathcal{L}_1(X)}$.

Proof. By Lemma 3.1(a), it is enough to check that the Abel-Jacobi map given by the image of [Z] with respect to the pullback map

$$(\mathrm{Id}_X \times \phi)^* \colon H^4(X \times \widetilde{Y}_A^{\geq 2}, \mathbb{Z}) \longrightarrow H^4(X \times F_1(X), \mathbb{Z})$$

is equal to $-AJ_{\mathscr{L}_1(X)}$. Equality (4.19) implies that there is a cycle Z_D'' supported on $X \times D$, where the scheme $D = F_1(X) \setminus U \stackrel{\delta}{\longleftrightarrow} F_1(X)$ is a finite subscheme of $F_1(X)$, such that

$$(\mathrm{Id}_X \times \phi)^*([Z]) + [\mathscr{L}_1(X)] = (i \times \mathrm{Id}_{F_1(X)})^*([Z'_F]) + (\mathrm{Id}_X \times \delta)_*([Z''_D])$$

Let us show that the Abel-Jacobi map defined by the right side of this equality is zero.

By Lemma 3.1(b), the Abel-Jacobi map corresponding to the cycle $(i \times Id_{F_1(X)})^*([Z'_F])$ is equal to the composition

$$H_1(F_1(X), \mathbf{Z}) \xrightarrow{\operatorname{AJ}_{Z'_F}} H_5(M_X, \mathbf{Z}) \xrightarrow{i^*} H_3(X, \mathbf{Z}).$$

Since $H_5(M_X, \mathbb{Z}) = 0$ by Proposition 2.6, this map vanishes. Similarly, the Abel-Jacobi map corresponding to the cycle $(\mathrm{Id}_X \times \delta)_*([\mathbb{Z}''_D])$ is equal to the composition

$$H_1(F_1(X), \mathbb{Z}) \xrightarrow{\delta^*} H_{-1}(D, \mathbb{Z}) \xrightarrow{\operatorname{AJ}_{\mathbb{Z}_D''}} H_3(X, \mathbb{Z}),$$

hence vanishes as well. This completes the proof of the proposition.

The above proposition relates the Abel–Jacobi maps AJ_Z and $AJ_{\mathscr{L}_1(X)}$. The next lemma uses the Clemens– Tyurin argument (Section 3.3) to show that the latter is surjective.

Lemma 4.16. Let X be a general GM threefold and let $\mathscr{L}_1(X) \to F_1(X)$ be the universal family of lines contained in X. The Abel-Jacobi map

$$\operatorname{AJ}_{\mathscr{L}_1(X)}: H_1(F_1(X), \mathbb{Z}) \longrightarrow H_3(X, \mathbb{Z})$$

is surjective.

Proof. Let Y be a general GM fourfold and let $X \subset Y$ be a general hyperplane section, so that X is a general GM threefold. Let $F_Y = F_1(Y)$ be the Hilbert scheme of lines contained in Y. We check that the assumptions of Proposition 3.3 (with m = 1) are satisfied.

Assumption (a) holds since $F_1(Y)$ is a smooth irreducible threefold by [DK19, Proposition 5.3]. Furthermore, the map $q: \mathscr{L}_1(Y) \to Y$ is generically finite of degree 6 ([DK19, Lemma 5.6]), hence (b) holds as well. Next, $F_1(X)$ is a smooth curve by Lemma 2.12 hence (c) holds. Finally,

$$H_3(Y,\mathbf{Z}) = H_5(Y,\mathbf{Z}) = 0$$

by [DK19, Proposition 3.1], hence (d) holds.

Applying Proposition 3.3, we deduce the surjectivity of $AJ_{\mathscr{L}_1(X)}$.

Combining the above results, we can now prove Theorem 4.4.

Proof of Theorem 4.4. Let $A \subset \bigwedge^3 V_6$ be a general Lagrangian subspace. As recalled in Section 2.3, any hyperplane $V_5 \subset V_6$ corresponding to a point of $Y_{A^{\perp}}^{\geq 2} \subset \mathbf{P}(V_6^{\vee})$ defines a smooth GM threefold X with A(X) = A. We choose a general such $[V_5]$, so that X is a general GM threefold. We also choose a nice line $L_0 \subset X$. A combination of Proposition 4.15 and Lemma 4.16 proves that the map AJ_Z is surjective. By Propositions 2.5 and 2.6, its source and target are free abelian groups of rank 20. Therefore, the Abel-Jacobi map (4.5) is an isomorphism.

The Abel-Jacobi map is defined by the cohomology class of an algebraic cycle, hence it preserves the Hodge structures and induces an isomorphism (4.6) between the corresponding abelian varieties: the Albanese variety of $\widetilde{Y}_A^{\geq 2}$ and the intermediate Jacobian of X.

Since the scheme $Z \subset X \times \widetilde{Y}_A^{\geq 2}$ is defined in Section 4.1 for all smooth X and all lines $L_0 \subset X$, and since this definition works in families, the maps (4.5) and (4.6) are by continuity isomorphisms for any A such that $Y_A^3 = \emptyset$ (so that the surface $\widetilde{Y}_A^{\geq 2}$ is smooth), any smooth X such that A(X) = A, and any line $L_0 \subset X$.

The abelian variety Jac(X) carries a canonical principal polarization and the isomorphism (4.6) transports it to a principal polarization on the abelian variety $Alb(\widetilde{Y}_A^{\geq 2})$. Since $Y_{A^{\perp}}^{\geq 2}$ is connected and the isomorphism (4.6) depends continuously on X, this principal polarization is independent of the choice of $[V_5] \in Y_{A^{\perp}}^{\geq 2}$. It is therefore canonical.

It remains to construct an isomorphism (4.4). Choose a point

$$(V_1, V_5) \in (Y_A^2 \times Y_{A^{\perp}}^2) \cap Fl(1, 5; V_6)$$

such that $A \cap (V_1 \wedge \bigwedge^2 V_5) = 0$ (it exists by [DK18, Lemma B.5]). Let X and X' be the smooth ordinary GM threefolds corresponding to the Lagrangian data sets (V_6, V_5, A) and $(V_6^{\vee}, V_1^{\perp}, A^{\perp})$ respectively. We will prove in Proposition 4.17 that there is a diagram

$$\begin{array}{c} \widehat{X} - - \stackrel{\psi}{-} - \rightarrow \widehat{X}' \\ \downarrow & \qquad \downarrow \\ X & \qquad X', \end{array}$$

where both vertical morphisms are blow ups of smooth rational curves and ψ is a flop. By [FW08, Proposition 3.1], the morphism $H^3(\widehat{X}; \mathbb{Z}) \to H^3(\widehat{X}'; \mathbb{Z})$ induced by the correspondence defined by the graph of ψ is an isomorphism of polarized Hodge structures. It induces in particular an isomorphism $\operatorname{Jac}(\widehat{X}) \xrightarrow{\sim} \operatorname{Jac}(\widehat{X}')$ of principally polarized abelian varieties between intermediate Jacobians. Therefore, there is a chain of isomorphisms

$$\operatorname{Alb}(\widetilde{Y}_{A}^{\geq 2}) \simeq \operatorname{Jac}(X) \simeq \operatorname{Jac}(\widehat{X}) \xrightarrow{\sim} \operatorname{Jac}(\widehat{X}') \simeq \operatorname{Jac}(X') \simeq \operatorname{Alb}(\widetilde{Y}_{A^{\perp}}^{\geq 2})$$

of principally polarized abelian varieties. This concludes the proof of the theorem.

4.5. The line transform

In this section, we revisit the birational isomorphism of [DK18, Proposition 4.19] and identify it with an elementary transformation along a line, a birational transformation between GM threefolds defined in [IP99, Proposition 4.3.1] and [DIM12, Section 7.2] (this relation was mentioned without proof in [DK18, Section 4.6]).

Proposition 4.17. Let $A \subset \bigwedge^3 V_6$ be a Lagrangian subspace with no decomposable vectors. Consider subspaces $V_1 \subset V_5 \subset V_6$ such that

(4.20)
$$[V_1] \in Y_A^2, \quad [V_5] \in Y_{A^{\perp}}^2, \quad A \cap (V_1 \land \bigwedge^2 V_5) = 0.$$

Let X and X' be the smooth ordinary GM threefolds corresponding to the Lagrangian data sets (V_6, V_5, A) and $(V_6^{\vee}, V_1^{\perp}, A^{\perp})$, and let $L_0 \subset X$ and $L'_0 \subset X'$ be the lines corresponding to the points $[V_1]$ of Y_{A,V_5}^2 and $[V_5^{\perp}]$ of $Y_{A^{\perp},V^{\perp}}^2$ via the maps (2.23).

There is a diagram of birational maps

(4.21)
$$Bl_{L_0}(X) - - - \frac{\psi}{-} - \rightarrow Bl_{L'_0}(X')$$
$$\begin{pmatrix} \beta & \rho_X & \rho_{X'} \\ X - - - - \phi & X & \xi \\ - - - - \phi & X', \\ \end{pmatrix}$$

where β and β' are the respective blow ups of X and X' along the lines L_0 and L'_0 , the birational maps ϖ and ϖ' are the respective linear projections of X from L_0 , and of X' from L'_0 , the morphisms ρ_X and $\rho_{X'}$ are small crepant extremal birational contractions, the variety \bar{X} is a normal, Gorenstein, cubic hypersurface in $\operatorname{Gr}(2, V_5/V_1) = \operatorname{Gr}(2, V_1^{\perp}/V_5^{\perp})$ with terminal singularities, and the map $\psi = \rho_{X'}^{-1} \circ \rho_X$ is a flop.

The proposition says that the birational isomorphism $\varpi'^{-1} \circ \varpi \colon X \to X'$ is the "elementary rational map with center along the line L_0 " in the sense of [IP99, (4.1.1)] (or elementary transformation along the line L_0) if the GM threefold X and the line L_0 are sufficiently general. We use the proposition and the fact that the elementary transformation is defined for any X and L_0 to specify what the moduli point of the result of the elementary transformation is in general (we use the description of the coarse moduli space \mathbf{M}_3^{GM} for GM threefolds given in (2.8)).

Corollary 4.18. Let X be a smooth GM threefold with moduli point $([A], [V_5]) \in \mathbf{M}_3^{GM}$ and let $L \subset X$ be any line. The moduli point of the result X'_L of the elementary transformation of X along the line L is $([A^{\perp}], \sigma(L)) \in \mathbf{M}_3^{GM}$.

Proof. Let \mathfrak{M}_3^{GM} be the moduli stack of smooth GM threefolds (see [DK20b]), let $\mathscr{X} \to \mathfrak{M}_3^{GM}$ be the universal family of threefolds over it, and let $F_1(\mathscr{X}/\mathfrak{M}_3^{GM})$ be the relative Hilbert scheme of lines. As we already mentioned, by [IP99, Section 4.1], the elementary transformation is defined for any line contained in any smooth GM threefold X. Moreover, this transformation can be performed for a family of lines and will produce a family of GM threefolds. This defines a morphism

$$\begin{array}{rcl} F_1(\mathscr{U}/\mathfrak{M}_3^{\mathrm{GM}}) & \longrightarrow & \mathbf{M}_3^{\mathrm{GM}} \\ ([X], [L]) & \longmapsto & [X'_L]. \end{array}$$

By [DK20b, Theorem 5.11] and [DK19, Theorem 4.7], the left side is irreducible and birational to

$$\{(A, V_5, V_1) \in \mathsf{LGr}_{\mathsf{ndv}}(\bigwedge^3 V_6) \times \mathsf{Fl}(1, 5; V_6) \mid [V_5] \in Y_{A^{\perp}}^{\geq 2}, \ [V_1] \in Y_A^{\geq 2}\} /\!\!/ \operatorname{PGL}(V_6).$$

By Proposition 4.17, over the dense open subset of triples (A, V_5, V_1) satisfying condition (4.20), this map coincides with the projection

$$(A, V_5, V_1) \mapsto (A^{\perp}, V_1^{\perp}) \in \mathbf{M}_3^{\mathrm{GM}}$$

Since the target is separated ([DK20b, Theorem 5.15]), by continuity, the two maps coincide everywhere. \Box

To prove Proposition 4.17, we start with some preliminaries. First, subspaces V_1 and V_5 satisfying the conditions (4.20) exist: this follows from [DK18, Lemma B.5] (where \widehat{Y}_A is defined in [DK18, (B.5)]). More exactly, for $[V_5]$ general in $Y_{A^{\perp}}^2$ (so that X is a general GM threefold associated with the fixed Lagrangian A), these conditions will be satisfied for a general $[V_1] \in Y_{A,V_5}^2$, corresponding to a general line $L_0 \subset X$. As explained in the proof of [DK18, Theorem 4.20], the conditions (4.20) are equivalent to

$$[V_1] \in Y^2_{A,V_5} \smallsetminus \Sigma_1(X) \quad \text{and} \quad [V_5^{\perp}] \in Y^2_{A^{\perp},V_1^{\perp}} \smallsetminus \Sigma_1(X'),$$

hence the lines L_0 and L'_0 are nice and the assumptions of [DK18, Proposition 4.19] are satisfied. It was explained in the proof of that proposition that these lines can be written as

(4.22)
$$L_0 = \mathbf{P}(V_1 \wedge V_3), \qquad L'_0 = \mathbf{P}(V_5^{\perp} \wedge V_3^{\perp})$$

for the same subspace V_3 such that $V_1 \subset V_3 \subset V_5$.

Following [DK18, Section 4.4], we introduce the second quadratic fibration

$$p_2: \mathbf{P}_X(V_5/\mathscr{U}_X) \longrightarrow \mathrm{Gr}(3, V_5)$$

and analogously for X', and study the diagram [DK18, (4.5)]

(4.23)
$$f \xrightarrow{\widetilde{\mu}_{2}} Gr(2, V_{5}/V_{1}) \xrightarrow{\widetilde{\mu}_{2}'} \widetilde{X}' \xrightarrow{f'} X',$$

where $\tilde{\rho}_2$ is obtained from ρ_2 by restriction to $Gr(2, V_5/V_1) \subset Gr(3, V_5)$ and analogously for $\tilde{\rho}'_2$. The next lemma is a refinement of [DK18, Lemma 4.18].

Lemma 4.19. The scheme \widetilde{X} has two irreducible components, $Bl_{L_0}(X)$ and $f^{-1}(L_0)$. They are both smooth of dimension 3 and meet transversely along the exceptional divisor of $Bl_{L_0}(X)$.

Moreover, the map $\tilde{\rho}_2 \colon \widetilde{X} \to \text{Gr}(2, V_5/V_1)$ is induced by the linear projection from the line L_0 .

Proof. We defined in [DK18, Section 4.4] the EPW quartic hypersurface $Z_A \subset Gr(3, V_6)$. For any subspace $V_1 \subset V_5$, we denote by

$$Z_{A,V_5} \subset Gr(3,V_5)$$
 and $Z_{A,V_1,V_5} \subset Gr(2,V_5/V_1)$

the subschemes obtained by intersecting Z_A with the subvarieties $Gr(3, V_5)$ and $Gr(2, V_5/V_1)$ of $Gr(3, V_6)$. Consider the commutative diagram

$$\mathbf{P}_{X}(V_{5}/\mathscr{U}_{X}) \longrightarrow \mathbf{P}_{M_{X}}(V_{5}/\mathscr{U}_{M_{X}}) \longrightarrow \mathbf{P}_{\mathrm{Gr}(2,V_{5})}(V_{5}/\mathscr{U}) == \mathrm{Fl}(2,3;V_{5})$$

$$\downarrow$$

$$\downarrow$$

$$Z_{A,V_{5}} \hookrightarrow \mathrm{Gr}(3,V_{5}),$$

where M_X was defined in (2.7) and the vertical arrow is the canonical projection, which induces the diagonal arrows (the left diagonal arrow factors through Z_{A,V_5} by [DK18, Proposition 4.10]). Pulling this diagram back by the inclusion $Gr(2, V_5/V_1) \subset Gr(3, V_5)$, we obtain the diagram



Indeed, $\operatorname{Gr}(2, V_5/V_1) \subset \operatorname{Gr}(3, V_5)$ is the zero-locus of the section of the vector bundle V_5/\mathcal{U}_3 corresponding to V_1 and, by [Kuzl6, Lemma 2.1], the zero-locus of the corresponding section on $\mathbf{P}_{\operatorname{Gr}(2, V_5)}(V_5/\mathcal{U})$ is the

blow up of $Gr(2, V_5)$ along the zero-locus of induced section of V_5/\mathcal{U} , which is equal to the locus $\mathbf{P}(V_1 \wedge V_5)$ of 2-dimensional subspaces in V_5 containing V_1 . Note also that the map

$$\operatorname{Bl}_{\mathbf{P}(V_1 \wedge V_5)}(\operatorname{Gr}(2, V_5)) \to \operatorname{Gr}(2, V_5/V_1)$$

is induced by the linear projection $V_5 \to V_5/V_1$ from V_1 , or equivalently by the linear projection $\mathbf{P}(\bigwedge^2 V_5) \to \mathbf{P}(\bigwedge^2 (V_5/V_1))$ from $\mathbf{P}(V_1 \wedge V_5)$.

Furthermore, $M_X \subset Gr(2, V_5)$ is the linear section by the subspace $\mathbf{P}(W) \subset \mathbf{P}(\bigwedge^2 V_5)$ which is transverse to $\mathbf{P}(V_1 \wedge V_5)$ by Lemma 4.6, because the line L_0 is nice; the pullback of $\mathbf{P}_{M_X}(V_5/\mathscr{U}_{M_X})$ is therefore $\mathrm{Bl}_{L_0}(M_X)$. Moreover, the map $\mathrm{Bl}_{L_0}(M_X) \to \mathrm{Gr}(2, V_5/V_1)$ is induced by the linear projection from $\mathbf{P}(W \cap (V_1 \wedge V_5)) = L_0$.

To prove the lemma, it remains to invoke the equality

$$\operatorname{Bl}_{L_0}(M_X) \times_{M_X} X = \operatorname{Bl}_{L_0}(X) \cup f^{-1}(L_0),$$

which holds because the first component is the strict transform of X and the second component is the exceptional divisor of $Bl_{L_0}(M_X) \to M_X$.

It was proved in [DK18, Lemma 4.18 and Proposition 4.19] that

- $\tilde{\rho}_2$ maps $f^{-1}(L_0)$ birationally onto the Schubert hyperplane divisor $D \subset Gr(2, V_5/V_1)$ parameterizing subspaces intersecting V_3/V_1 , where V_3 was defined in (4.22);
- $\tilde{\rho}_2$ maps $\operatorname{Bl}_{L_0}(X)$ birationally onto a cubic hypersurface $\bar{X} \subset \operatorname{Gr}(2, V_5/V_1)$;
- the image of $\tilde{\rho}_2$ is the quartic hypersurface Z_{A,V_1,V_5} ; it is therefore equal to $\bar{X} \cup D$.

We denote by ρ_X : Bl_{L₀}(X) $\rightarrow \bar{X}$ the (birational) restriction of $\tilde{\rho}_2$ and we define $\rho_{X'}$ similarly.

Proof of Proposition 4.17. We have already constructed the left part of the diagram (4.21). The right part is constructed analogously. It remains to prove that ψ is a flop.

As explained in [IP99, Lemma 4.1.1 and Corollary 4.3.2], ρ_X is a flopping contraction. The same is true for $\rho_{X'}$, so ψ is either a flop or an isomorphism. If ψ is an isomorphism, the maps β and β' are the contractions of the same extremal ray, hence $X \simeq X'$. Let us show that this is impossible.

Indeed, we can perform this construction on a fixed X (that is, with A and V_5 fixed) but with $[V_1]$ varying in the curve Y_{A,V_5}^2 . Locally, the map ψ will remain an isomorphism and the threefolds X' obtained by the construction will be all isomorphic to X. But this is impossible: by definition, X' is the ordinary GM threefold corresponding to the Lagrangian data set $(V_6^{\vee}, V_1^{\perp}, A^{\perp})$, hence its moduli point describes the curve

$$Y_{A,V_5}^2 / / \operatorname{PGL}(V_6)_A \subset \mathfrak{p}_3^{-1}([A^{\perp}]) \subset \mathbf{M}_3^{\operatorname{GM}}$$

(recall that the map ρ_3 was defined in (2.9) and that the group $PGL(V_6)_A$ is finite). It follows that ψ is a flop and the proof of the proposition is complete.

Remark 4.20. One can describe the maps ρ_X and $\rho_{X'}$ in (4.21) further: in fact, the intersections with $\operatorname{Gr}(2, V_5/V_1)$ of the subscheme $\Sigma_2(X) \subset \operatorname{Gr}(3, V_5)$ defined in [DK18, (4.2)] and of the analogous subscheme $\Sigma_2(X') \subset \operatorname{Gr}(2, V_6/V_1)$ are cubic surfaces (cubic scrolls or cones) $\Sigma_{2,V_1}(X)$ and $\Sigma_{2,V_5}(X')$ such that

$$\bar{X} \cap \mathsf{D} = \Sigma_{2,V_1}(X) \cup \Sigma_{2,V_5}(X').$$

The morphisms ρ_X and $\rho_{X'}$ are the blow ups of \bar{X} along the Weil divisors $\Sigma_{2,V_1}(X)$ and $\Sigma_{2,V_5}(X')$, respectively.

5. Intermediate Jacobians of GM fivefolds

In this section, we perform, for GM fivefolds, a construction analogous to what we did in Section 4 for threefolds. The curves in the construction are replaced by surfaces: lines by planes, elliptic quintic curves by quintic del Pezzo surfaces, and rational quartic curves by rational quartic surface scrolls. In Section 5.5, we give an alternative proof of the main result.

5.1. Family of surfaces

Given an arbitrary GM fivefold X with associated Lagrangian A, we begin by choosing an arbitrary σ plane $\Pi_0 \subset X$ (that is, a point of $F^2_{\sigma}(X)$; see (2.2)). We consider the open surface $S_0 \subset Y_A^{\geq 2}$ defined by (2.14) and the family of quadrics $\mathscr{Q}_0 \to S_0$ obtained by restricting to S_0 the universal family (2.11) of 8-dimensional quadrics containing X. We denote by $F^5(\mathscr{Q}_0/S_0) \to S_0$ the relative Hilbert scheme of linear 5-spaces in the fibers of $\mathscr{Q}_0 \to S_0$ and by $F^5_{\Pi_0}(\mathscr{Q}_0/S_0) \to S_0$ the subscheme parameterizing those 5-spaces which contain the plane Π_0 . Applying Corollary 2.9, we obtain an isomorphism

(5.1)
$$F_{\Pi_0}^5(\mathscr{Q}_0/S_0) \simeq \widetilde{S}_0 := \widetilde{Y}_A^2 \times_{Y_A^2} S_0$$

of schemes over S_0 . In particular, the canonical map $F^5_{\Pi_0}(\mathscr{Q}_0/S_0) \to S_0$ is the double étale covering $\pi : \widetilde{S}_0 \to S_0$ induced by the double covering π_A .

Note that \widetilde{S}_0 is a smooth surface. Let

$$\widetilde{\mathscr{Q}}_0 := \mathscr{Q}_0 \times_{S_0} \widetilde{S}_0$$

be the base change of the family of quadrics $\mathscr{Q}_0 \to S_0$ along π . We have a canonical map

$$\widetilde{S}_0 \to F^5_{\Pi_0}(\mathscr{Q}_0/S_0) \times_{S_0} \widetilde{S}_0 \hookrightarrow F^5(\mathscr{Q}_0/S_0) \times_{S_0} \widetilde{S}_0 \simeq F^5(\widetilde{\mathscr{Q}}_0/\widetilde{S}_0),$$

where the first map is the product of the isomorphism (5.1) with the identity map. By construction, it is a section of the projection $F^5(\widetilde{\mathscr{Q}}_0/\widetilde{S}_0) \to \widetilde{S}_0$.

Let $\mathscr{P}^5 \subset \widetilde{\mathscr{Q}_0} \subset \mathbf{P}(W) \times \widetilde{S}_0$ be the pullback of the universal family of linear 5-spaces over $F^5(\widetilde{\mathscr{Q}_0}/\widetilde{S}_0)$ along this section. Set

where the Grassmannian hull $M_X \subset \mathbf{P}(W)$ was defined in (2.7).

Proposition 5.1. The map $Z_0 \to \widetilde{S}_0$ is a flat family of surfaces in X containing Π_0 with Hilbert polynomial

(5.3)
$$h(t) = \frac{5}{2}(t^2 + t) + 1.$$

In particular, $Z_0 \subset X \times \widetilde{S}_0$.

Proof. Let $y \in \widetilde{S}_0$ and set $[v] := \pi(y) \in \mathbf{P}(V_6) \smallsetminus \mathbf{P}(V_5)$. The fiber of Z_0 over y is

$$Z_{0,y} := M_X \cap \mathscr{P}_y^5 = \mathsf{CGr}(2, V_5) \cap \mathscr{P}_y^5.$$

Since the cone $\operatorname{CGr}(2, V_5) \subset \operatorname{P}(\operatorname{C} \oplus \bigwedge^2 V_5)$ has codimension 3 and degree 5, the intersection $\operatorname{CGr}(2, V_5) \cap \mathscr{P}_y^5$ has dimension at least 2 and degree at most 5 (and if the dimension is 2, the degree is 5). Furthermore, \mathscr{P}_y^5 is contained in Q_v , hence

$$Z_{0,v} \subset M_X \cap Q_v = X$$

Since X contains no divisors of degrees less than 10, we have $\dim(Z_{0,y}) \leq 3$ and, moreover, if $\dim(Z_{0,y}) = 3$, any irreducible 3-dimensional component $Z_{0,y}$ has even degree ([DK19, Corollary 3.5]). By Lemma 2.3, its image in $Gr(2, V_5)$ must be a hyperplane section of $Gr(2, V_4)$ and Lemma 2.7 gives a contradiction. Therefore $Z_{0,y}$ is a surface. This argument also proves the inclusion

$$Z_0 \subset X \times \widetilde{S}_0.$$

Since the surface $Z_{0,y}$ is a dimensionally transverse linear section of CGr(2, V_5), we obtain from Lemma 2.4 a resolution

$$0 \to \mathscr{O}_{\mathscr{P}_{y}^{5}}(-5) \to \mathscr{O}_{\mathscr{P}_{y}^{5}}(-3)^{\oplus 5} \to \mathscr{O}_{\mathscr{P}_{y}^{5}}(-2)^{\oplus 5} \to \mathscr{O}_{\mathscr{P}_{y}^{5}} \to \mathscr{O}_{Z_{0,y}} \to 0.$$

It follows that the Hilbert polynomial of $Z_{0,y}$ is given by (5.3). Since it is independent of y, the family of surfaces Z_0 is flat over \widetilde{S}_0 . Finally, since $\Pi_0 \subset M_X$ and $\Pi_0 \subset \mathscr{P}_y^5$ by construction, we obtain $\Pi_0 \subset Z_{0,y}$. \Box

Applying to the family $Z_0 \to \widetilde{S}_0$ the Hilbert closure construction from Definition 4.2, we obtain the following result.

Lemma 5.2. There is a subscheme

(5.4)

such that, away from a finite subset of the surface $\widetilde{Y}_A^{\geq 2}$, the map $Z \to \widetilde{Y}_A^{\geq 2}$ is a flat family of surfaces with Hilbert polynomial (5.3) containing the plane Π_0 . Moreover, we have

 $Z\subset X\times \widetilde{Y}_A^{\geq 2}$

$$Z \times_{\widetilde{Y}_A^{\geq 2}} \widetilde{S}_0 = Z_0$$

as subschemes of $X \times \widetilde{S}_0$.

By Proposition 5.1, the schemes Z_0 and Z have pure dimension 4.

The main result of this section is the following theorem (recall from Propositions 2.5 and 2.6 that the abelian groups $H_1(\widetilde{Y}_{A(X)}^{\geq 2}, \mathbb{Z})$ and $H_5(X, \mathbb{Z})$ are both free of rank 20 and from Theorem 4.4 that the abelian variety $\text{Alb}(\widetilde{Y}_{A(X)}^{\geq 2})$ is endowed with a canonical principal polarization when $Y_{A(X)}^{\geq 3} = \emptyset$).

Theorem 5.3. Let X be a smooth GM fivefold and let Π_0 be a σ -plane contained in X. Let $Z \subset X \times \widetilde{Y}_{A(X)}^{\geq 2}$ be the subscheme defined by (5.4). If $Y_{A(X)}^{\geq 3} = \emptyset$, the Abel-Jacobi map

$$\mathsf{AJ}_Z \colon H_1(\widetilde{Y}_{A(X)}^{\geq 2}, \mathbf{Z}) \longrightarrow H_5(X, \mathbf{Z})$$

is an isomorphism of integral Hodge structures. It induces an isomorphism

(5.5)
$$\operatorname{Alb}(Y^{\geq 2}_{A(X)}) \xrightarrow{\sim} \operatorname{Jac}(X)$$

of principally polarized abelian varieties from the Albanese variety of the surface $\widetilde{Y}_{A(X)}^{\geq 2}$ to the intermediate Jacobian of X.

The proof of this theorem takes up Sections 5.2-5.4.

5.2. The boundary of the family

Let X be a smooth GM fivefold. To prove Theorem 5.3, we study the family of surfaces Z described in Lemma 5.2 over the boundary $\widetilde{Y}_A^{\geq 2} \setminus \widetilde{S}_0$. We assume in this section that X is general (in particular it is ordinary), so that the curve $Y_{A,V_5}^{\geq 2}$ is smooth.

Consider the Hilbert scheme $F_{\sigma}^2(X)$ of σ -planes on X; we identify it with the smooth connected curve $\widetilde{Y}_{A,V_5}^{\geq 2}$ via the isomorphism (2.19). For $y \in \widetilde{Y}_{A,V_5}^{\geq 2}$, we denote by $\Pi_y \subset X$ the corresponding σ -plane.

We denote by $y_0 \in \widetilde{Y}_{A,V_5}^{\geq 2}$ the point such that $\Pi_{y_0} = \Pi_0$ is the plane chosen in Section 5.1. Set $[v_0] := \pi_A(y_0)$. By Lemma 2.1(b), we have, for an appropriate hyperplane $V_4 \subset V_5$,

(5.6)
$$\Pi_0 = \mathbf{P}(v_0 \wedge V_4).$$

We set

$$Y_{A,V_4}^{\geq 2} := Y_A^{\geq 2} \cap \mathbf{P}(V_4).$$

This is a hyperplane section of the smooth curve Y^2_{A,V_5} hence is finite; the induced double coverings of this set parameterizes σ -planes on X that intersect Π_0 .

Denote by $\widetilde{\mathscr{Q}}$ the pullback of the family of quadrics \mathscr{Q} along the map $\widetilde{Y}_A^{\geq 2} \to Y_A^{\geq 2}$. Recall that a section $\widetilde{S}_0 \to F_{\Pi_0}^5(\widetilde{\mathscr{Q}}_0/\widetilde{S}_0)$ was constructed in Section 5.1. Since the surface $\widetilde{Y}_A^{\geq 2}$ is smooth and the Hilbert scheme $F_{\Pi_0}^5(\widetilde{\mathscr{Q}}/\widetilde{Y}_A^{\geq 2})$ is proper over $\widetilde{Y}_A^{\geq 2}$, this section extends to an open subset

(5.7)
$$\widetilde{S}_{0+} \subset \pi_A^{-1}(Y_A^{\geq 2} \smallsetminus (Y_{A,V_4}^{\geq 2} \cup Y_{A,V_4}^{\geq 2})) \subset \widetilde{Y}_A^{\geq 2}$$

which contains a general point of the curve $\widetilde{S}_{0,V_5} := \widetilde{S}_{0+} \smallsetminus \widetilde{S}_0 \subset \widetilde{Y}_{A,V_5}^{\geq 2}$. We denote by

$$\mathscr{P}^5_+ \subset \mathbf{P}(\bigwedge^2 V_5) \times \widetilde{S}_0.$$

the corresponding family of 5-spaces and, for $y \in \widetilde{S}_{0+}$, by $\mathscr{P}^5_{+y} \subset \mathbf{P}(\bigwedge^2 V_5)$ the corresponding linear 5-space. By definition, we have $\Pi_0 \subset \mathscr{P}^5_{+y}$ for each $y \in \widetilde{S}_{0,V_5}$.

Lemma 5.4. For each point $y \in S_{0,V_5}$, we have

(5.8)
$$\mathscr{P}^{5}_{+y} = \langle \Pi_{y}, \Pi_{0} \rangle.$$

Proof. Set $[v] := \pi_A(v) \in Y_{A,V_5}^{\geq 2}$ and let $W_6 \subset \bigwedge^2 V_5$ be the 6-dimensional subspace corresponding to the 5-space $\mathscr{P}_v^5 \subset \mathbf{P}(\bigwedge^2 V_5)$. By definition, we have

$$\mathbf{P}(W_6) \subset Q_v = \operatorname{Cone}_{\mathbf{P}(v \wedge V_5)}(\operatorname{Gr}(2, V_5/v)).$$

Since $Gr(2, V_5/v)$ is a smooth 4-dimensional quadric, the maximal dimension of a linear subspace that it contains is 2, hence the subspace

$$W_v := W_6 \cap (v \wedge V_5)$$

is at least 3-dimensional. We claim that $\mathbf{P}(W_v)$ is contained in X.

Let $\{w_i\}$ be a basis of W_v , let q_v be an equation of Q_v , and consider a line $\langle [v], [v'] \rangle \subset \mathbf{P}(V_6)$ tangent to $Y_A^{\geq 2}$ at [v], with $[v'] \in \mathbf{P}(V_6) \setminus \mathbf{P}(V_5)$. Let $\operatorname{Spec}(\mathbf{C}[\epsilon]/\epsilon^2) \to Y_A^{\geq 2}$ be the corresponding morphism that takes the closed point to [v]. Since \widetilde{S}_{0+} is étale over $Y_{A,V_5}^{\geq 2}$, the morphism can be lifted to a morphism

$$\operatorname{Spec}(\mathbf{C}[\epsilon]/\epsilon^2) \longrightarrow \widetilde{S}_{0+}$$

that takes the closed point to y. This implies that there are vectors w'_i in $\bigwedge^2 V_5$ such that the subspace in $\bigwedge^2 V_5$ generated by $w_i + \epsilon w'_i$ is isotropic for the quadratic form $q_v + \epsilon q_{v'}$, where $q_{v'}$ is an equation of $Q_{v'}$. We have therefore

$$0 = (q_v + \epsilon q_{v'})(w_i + \epsilon w'_i, w_j + \epsilon w'_j) = \epsilon(q_v(w_i, w'_j) + q_v(w_j, w'_i) + q_{v'}(w_i, w_j)).$$

Note that $q_v(w_i, w'_j) = q_v(w_j, w'_i) = 0$, since $w_i, w_j \in v \wedge V_5 = \text{Ker}(q_v)$ for all i, j. It follows that $q_{v'}(w_i, w_j) = 0$ for all i, j, hence W_y is isotropic for $q_{v'}$, that is, $\mathbf{P}(W_y) \subset Q_{v'}$. Since $\mathbf{P}(W_y) \subset \mathbf{P}(v \wedge V_5) \subset \text{Gr}(2, V_5)$, we conclude that

$$\mathbf{P}(W_v) \subset \operatorname{Gr}(2, V_5) \cap Q_{v'} = X,$$

thus proving the claim.

Since $\mathbf{P}(W_y) \subset \mathbf{P}(v \wedge V_5) \cap X$ and X contains no linear 3-spaces ([DK19, Theorem 4.2]), it follows that $\dim(W_v) = 3$ and $\mathbf{P}(W_v)$ is a σ -plane on X. Moreover, the induced map

$$\begin{aligned} \widetilde{S}_{0,V_5} &\longrightarrow F^2_{\sigma}(X) \\ y &\longmapsto \mathbf{P}(W_y) \end{aligned}$$

is a $Y_{A,V_5}^{\geq 2}$ -morphism. But $F_{\sigma}^2(X) \simeq \widetilde{Y}_{A,V_5}^{\geq 2}$, while \widetilde{S}_{0,V_5} is an open subscheme in $\widetilde{Y}_{A,V_5}^{\geq 2}$, and $\widetilde{Y}_{A,V_5}^{\geq 2}$ is a connected étale covering of $Y_{A,V_5}^{\geq 2}$. Therefore, replacing if necessary the isomorphism (2.19) by its composition with the involution of the double covering, we may assume that the above map $\widetilde{S}_{0,V_5} \to F_{\sigma}^2(X)$ coincides with the embedding $\widetilde{S}_{0,V_5} \hookrightarrow \widetilde{Y}_{A,V_5}^{\geq 2}$, hence $\mathbf{P}(W_y) = \Pi_y$ for all $y \in \widetilde{S}_{0,V_5}$.

This means that there is an inclusion $\Pi_y \subset \mathscr{P}^5_{+y}$ for all $y \in \widetilde{S}_{0,V_5}$. Since $\Pi_0 \subset \mathscr{P}^5_{+y}$ by definition, the right side of (5.8) is contained in the left side. Finally, since $\Pi_0 \cap \Pi_y = \emptyset$ by definition of \widetilde{S}_{0+} (since planes intersecting Π_0 are parameterized by the double cover of the subscheme $Y_{A,V_4}^{\geq 2}$), the inclusion is an equality.

The family \mathscr{P}^5_+ of projective 5-spaces over \widetilde{S}_{0+} agrees by construction with the family \mathscr{P}^5 over $\widetilde{S}_0 \subset \widetilde{S}_{0+}$. We set

(5.9)
$$Z_{0+} := \mathscr{P}_{+}^{5} \cap (M_X \times \widetilde{S}_{0+})$$

Comparing this with (5.2), we obtain

$$Z_{0+} \times_{\widetilde{S}_{0+}} \widetilde{S}_0 = Z_0.$$

We denote by

$$Z_{0,V_5} := Z_{0,+} \times_{\widetilde{S}_{0+}} \widetilde{S}_{0,V_5} \subset M_X \times \widetilde{S}_{0,V_5}$$

the restriction of the family (5.9) to the curve $\widetilde{S}_{0,V_5} = \widetilde{S}_{0+} \setminus \widetilde{S}_0$.

Proposition 5.5. Let X be a general GM fivefold. The map $Z_{0,V_5} \to \widetilde{S}_{0,V_5}$ is a flat family of 3-dimensional cubic scrolls $\mathbf{P}^1 \times \mathbf{P}^2$.

Proof. Let $y \in \widetilde{S}_{0,V_5}$ and set again $[v] := \pi(y) \in \mathbf{P}(V_5)$. By (5.8), the linear 5-space \mathscr{P}_{+y}^5 is the linear span of the planes Π_0 and Π_y , that is, a hyperplane in $\mathbf{P}(V_2 \wedge V_5)$, where $V_2 \subset V_5$ is the subspace spanned by v_0 and v. Therefore, $\operatorname{Gr}(2, V_5) \cap \mathscr{P}_{+y}^5$ is a hyperplane section of the cone $\operatorname{Cone}_{\mathbf{P}(\wedge^2 V_5)}(\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2))$ (see Lemma 2.2).

The vertex $[\wedge^2 V_2] = [v_0 \wedge v]$ of the cone does not belong to \mathscr{P}^5_{+y} : if it did, Π_0 and Π_y would intersect at the point $[v_0 \wedge v]$ and this would contradict the definition of S_{0+} . Therefore, the fiber of Z_{0,V_5} over y is isomorphic to the 3-dimensional cubic scroll $\mathbf{P}(V_2) \times \mathbf{P}(V_5/V_2)$.

Propositions 5.1 and 5.5 show that all the components of Z_0 and Z_{0,V_5} have dimension 4. They are components of the scheme Z_{0+} . We also consider the Hilbert closure

of Z_{0,V_5} in $M_X \times F^2_{\sigma}(X)$, constructed as in Definition 4.2.

5.3. A relation between the subschemes

Let X be a general GM fivefold. In (5.4) and (5.10), we have constructed subschemes $Z \subset X \times \widetilde{Y}_A^{\geq 2}$ and $Z_F \subset M_X \times F_{\sigma}^2(X)$. The proof of Theorem 5.3 is based on a relation between the schemes $Z \cap (X \times \widetilde{S}_{0,V_5})$ and $Z_F \cap (X \times \widetilde{S}_{0,V_5})$, where the curve \widetilde{S}_{0,V_5} is considered as a subscheme of both the surface $\widetilde{Y}_A^{\geq 2}$ and the curve $F_{\sigma}^2(X)$.

Consider the commutative diagram



where $\mathscr{L}^2_{\sigma}(X) \subset X \times F^2_{\sigma}(X)$ is the universal family of σ -planes, $i: X \hookrightarrow M_X$ is the embedding, and $\tilde{\sigma}$ is the isomorphism (2.19).

Proposition 5.6. We have an equality

(5.12)
$$Z_F \cap (X \times \widetilde{S}_{0,V_5}) = (\tilde{\sigma}^{-1}(Z) \cap (X \times \widetilde{S}_{0,V_5})) + (\mathscr{L}^2_{\sigma}(X) \cap (X \times \widetilde{S}_{0,V_5}))$$

of cycles.

Proof. The left side of (5.12) can be rewritten as

$$Z_F \cap (X \times \widetilde{S}_{0,V_5}) = Z_{0,V_5} \cap (X \times \widetilde{S}_{0,V_5}) = Z_{0,V_5} \times_{M_X} X.$$

By Proposition 5.5, the morphism $Z_{0,V_5} \to \widetilde{S}_{0,V_5}$ is a flat family of smooth 3-dimensional cubic scrolls. Since X contains no such threefolds ([DK19, Corollary 3.5]), it contains no fibers of Z_{0,V_5} . Therefore, the morphism

$$Z_{0,V_5} \times_{M_X} X \longrightarrow \widetilde{S}_{0,V_5}$$

is a flat family of surfaces whose fiber over $y \in \widetilde{S}_{0,V_5}$ is the dimensionally transverse intersection

$$S_y := Z_{0, V_5, y} \cap Q_0$$

of the smooth 3-dimensional cubic scroll $Z_{0,V_5,y} \simeq \mathbf{P}^2 \times \mathbf{P}^1$ with any non-Plücker quadric Q_0 containing X. Such an intersection is a surface of class $2f_2 + 2f_1$ in $Z_{0,V_5,y}$, where f_i is the preimage of the hyperplane class on \mathbf{P}^i under the projection $Z_{0,V_5,y} \simeq \mathbf{P}^2 \times \mathbf{P}^1 \rightarrow \mathbf{P}^i$.

By (5.8) and (5.9), the planes Π_0 and Π_y are contained in the scroll $Z_{0,V_5,y}$. Since they are also contained in X, they are contained in the quadric Q_0 . It follows that they are components of S_y , each of class f_1 . Therefore,

$$S_v = \Pi_0 + \Pi_v + S'_v$$

where $S'_y \subset Z_{0,V_5,y}$ is a surface of class $2f_2$, that is, the product of a conic in \mathbf{P}^2 with \mathbf{P}^1 . In particular, it has degree 4 and contains no planes. Since $Z_y \subset Z_{0,V_5,y}$ is a surface of degree 5 that contains Π_0 , we have $Z_y = S'_y + \Pi_0$ for all $y \in \widetilde{S}_{0,V_5}$; this proves (5.12).

5.4. Abel-Jacobi maps

Let X be a smooth ordinary GM fivefold with associated Lagrangian A. Assume $Y_A^3 = \emptyset$, so that $\widetilde{Y}_A^{\geq 2}$ is a smooth surface and the curve $\widetilde{Y}_{A,V_5}^{\geq 2}$, hence also the Hilbert scheme $F_{\sigma}^2(X)$ of σ -planes in X, is a smooth curve. Let $\mathscr{L}_{\sigma}^2(X) \subset X \times F_{\sigma}^2(X)$ denote the universal family of σ -planes on X. Consider the Abel-Jacobi maps

$$\mathcal{A}_Z \colon H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z}) \to H_5(X, \mathbb{Z}) \quad \text{and} \quad \mathcal{A}_{\mathcal{L}_{\sigma}^2(X)} \colon H_1(F_{\sigma}^2(X), \mathbb{Z}) \to H_5(X, \mathbb{Z})$$

and recall the isomorphism $\tilde{\sigma} \colon F^2_{\sigma}(X) \xrightarrow{\sim} \widetilde{Y}^{\geq 2}_{A,V_5}$ from (2.19).

Proposition 5.7. Let X be a smooth ordinary GM fivefold with associated Lagrangian A. Assume that $Y_A^3 = \emptyset$ and $F_2^{\sigma}(X)$ is smooth. The composition of maps

$$H_1(F^2_{\sigma}(X), \mathbf{Z}) \xrightarrow{\tilde{\sigma}_*} H_1(\widetilde{Y}^{\geq 2}_A, \mathbf{Z}) \xrightarrow{\mathsf{AJ}_Z} H_5(X, \mathbf{Z})$$

is equal to the map $-AJ_{\mathcal{L}^2_{\sigma}(X)}$.

Proof. Analogous to the proof of Proposition 4.15.

The above proposition connects the Abel-Jacobi maps AJ_Z and $AJ_{\mathscr{L}^2_{\sigma}(X)}$. The next lemma uses the Clemens-Tyurin argument (Section 3.3) to show that the latter is surjective.

Lemma 5.8. Let X be a general GM fivefold. The Abel-Jacobi map

$$\mathsf{AJ}_{\mathscr{L}^2_{\sigma}(X)} \colon H_1(F^2_{\sigma}(X), \mathbf{Z}) \longrightarrow H_5(X, \mathbf{Z})$$

is surjective.

Proof. Let Y be a general GM sixfold and let $X \subset Y$ be a general hyperplane section, so that X is a general GM fivefold. Set $F_Y := F_{\sigma}^2(Y)$, the Hilbert scheme of σ -planes contained in Y. We check that the assumptions of Proposition 3.3 (with m = 2) are satisfied.

Assumption (a) holds because $F_{\sigma}^2(Y)$ is a smooth irreducible fourfold by [DK19, Corollary 5.13]. Furthermore, the map $q: \mathscr{L}_{\sigma}^2(Y) \to Y$ is generically finite of degree 12 ([DK19, Lemma 5.15]), hence (b) holds

as well. Next, $F_{\sigma}^2(X)$ is a smooth curve by Lemma 2.10, hence (c) holds. Finally, $H_5(Y, \mathbb{Z}) = H_7(Y, \mathbb{Z}) = 0$ by [DK19, Proposition 3.1], hence (d) holds.

Applying Proposition 3.3, we deduce the surjectivity of $AJ_{\mathscr{L}^2(X)}$.

Combining the above results, we can now prove Theorem 5.3.

Proof of Theorem 5.3. Assume first that the GM fivefold X is general. A combination of Proposition 5.7 and Lemma 5.8 proves that the map AJ_Z is surjective. By Propositions 2.5 and 2.6, its source and target are free abelian groups of rank 20. Therefore, the Abel-Jacobi map is an isomorphism.

Since the Abel–Jacobi map is defined by the cohomology class of an algebraic cycle, it preserves the Hodge structures, hence induces an isomorphism of the corresponding abelian varieties: the Albanese variety of $\widetilde{Y}_{A(X)}^{\geq 2}$ and the intermediate Jacobian of X.

Since the scheme $Z \subset X \times \widetilde{Y}_{A(X)}^{\geq 2}$ was defined in Section 5.1 for all smooth X and all σ -planes $\Pi_0 \subset X$ and since this definition works in families, these two statements follow by continuity for any X such that $Y_{A(X)}^3 = \emptyset$.

It remains to prove that the isomorphism (5.5) respects the principal polarizations. For X very general, the Picard number of Jac(X) is 1 by Corollary 3.6, hence any two principal polarizations on Jac(X) coincide. This proves the claim for very general X and, by continuity, for any smooth X such that $\widetilde{Y}_{A(X)}^{\geq 2}$ is also smooth.

5.5. Simplicity argument

We give an alternative argument for the isomorphism (5.5) for a smooth GM fivefold X, based on a simplicity result of independent interest, analogous to the one proved in Proposition 3.4.

Let S be a smooth connected projective surface and let $j: C \hookrightarrow S$ be a smooth (irreducible) ample curve. By the Lefschetz theorem, the morphism $j_*: H_1(C, \mathbb{Z}) \to H_1(S, \mathbb{Z})$ is surjective, hence the induced morphism

$$\operatorname{Jac}(C) \simeq \operatorname{Alb}(C) \longrightarrow \operatorname{Alb}(S)$$

is surjective with connected kernel. We denote this kernel by K(C, S).

Consider now a connected double étale cover $\pi: \widetilde{S} \to S$ and set $\widetilde{C} := \pi^{-1}(C)$, a smooth ample curve on the surface \widetilde{S} .

Lemma 5.9. There is a commutative diagram

(5.13)

$$K(\widetilde{C},C) \rightarrow P(\widetilde{C},C) \rightarrow P(\widetilde{S},S) \rightarrow 0$$

$$\downarrow \qquad \downarrow \qquad \downarrow$$

$$0 \rightarrow K(\widetilde{C},\widetilde{S}) \rightarrow \operatorname{Jac}(\widetilde{C}) \rightarrow \operatorname{Alb}(\widetilde{S}) \rightarrow 0$$

$$\downarrow \pi_{*} \qquad \downarrow \pi_{*} \qquad \downarrow \pi_{*}$$

$$0 \rightarrow K(C,S) \rightarrow \operatorname{Jac}(C) \rightarrow \operatorname{Alb}(S) \rightarrow 0$$

$$\downarrow \qquad \downarrow \qquad \downarrow$$

$$0 \qquad 0 \qquad 0,$$

where $K(\widetilde{C}, C)$, $P(\widetilde{C}, C)$ (the Prym variety of the double cover $\widetilde{C} \to C$), and $P(\widetilde{S}, S)$ are the neutral components of the respective kernels of the vertical maps π_* induced by π .

Proof. The surjectivity of the maps $Jac(\widetilde{C}) \to Jac(C)$ and $Alb(\widetilde{S}) \to Alb(S)$ is obvious. The only thing we have to prove is the surjectivity of the map $K(\widetilde{C},\widetilde{S}) \to K(C,S)$ or, equivalently, the surjectivity of the map $P(\widetilde{C},C) \to P(\widetilde{S},S)$. On the level of cotangent spaces, the surjectivity of this second map corresponds to the injectivity of the restriction $H^1(S,\eta) \to H^1(C,\eta|_C)$, where η is the line bundle of order 2 on S

corresponding to the double étale covering π . Its kernel is controlled by $H^1(S, \eta(-C))$, which vanishes by Kodaira vanishing and Serre duality because $\eta(C)$ is ample on S. This proves the injectivity of the morphism $H^1(S, \eta) \to H^1(C, \eta|_C)$, hence the lemma.

The next statement is the main result of this section. The definition of "trivial endomorphism ring" can be found in Section 3.4.

Theorem 5.10. Let $S \subset \mathbf{P}^N$ be a smooth connected projective surface and let $\pi : \widetilde{S} \to S$ be a connected double étale cover. Let $H \subset \mathbf{P}^N$ be a very general hyperplane and set $C := S \cap H$. With the notation above, the endomorphism ring of the abelian variety $K(\widetilde{C}, C)$ is trivial.

Proof. We use the notation of Section 3.4. Choose a Lefschetz pencil $f: S \to \mathbf{P}^1$ of hyperplane sections of S. The connected double étale cover $\pi: \widetilde{S} \to S$ induces for each $t \in \mathbf{P}^1$ a connected double étale cover $\pi_t: \widetilde{C}_t \to C_t$ between fibers. Denote by $j_t: C_t \hookrightarrow S$ and $\tilde{j}_t: \widetilde{C}_t \hookrightarrow \widetilde{S}$ the embeddings.

For $t \in \mathbf{P}^1 \setminus \{t_1, \dots, t_r\}$, the involution τ of \widetilde{S} attached to π acts on each summand of the orthogonal direct sum decomposition

$$H^{1}(\widetilde{C}_{t}, \mathbf{Q}) = H^{1}(\widetilde{C}_{t}, \mathbf{Q})_{\mathrm{van}} \oplus \widetilde{j}_{t}^{*} H^{1}(\widetilde{S}, \mathbf{Q})$$

from (3.3) and it preserves the symplectic form given by cup-product. The τ -invariant subspaces are

$$H^1(C_t, \mathbf{Q}) = H^1(C_t, \mathbf{Q})_{\operatorname{van}} \oplus j_t^* H^1(S, \mathbf{Q})_{\operatorname{van}}$$

The isogeny class of the abelian variety $K(C_t, S)$ defined in (5.13) is obtained from the Hodge structure of $H^1(C_t, \mathbf{Q})_{\text{van}}$, hence its endomorphism ring is trivial by Proposition 3.4. Therefore, to study the isogeny class of the neutral component $K(\widetilde{C}_t, C_t)$ of the kernel of the surjection

$$K(\widetilde{C}_t,\widetilde{S}) \longrightarrow K(C_t,S),$$

we need to study the rational Hodge structure on the τ -antiinvariant subspace $H^1(\widetilde{C}_t, \mathbf{Q})^-_{\text{van}}$.

1

For each $i \in \{1, ..., r\}$, the curve C_{t_i} has two nodes over the node of C_{t_i} , hence there are two disjoint vanishing cycles δ'_i and $\delta''_i = \tau^*(\delta'_i)$. Since the vanishing cycles span the vector space $H^1(\widetilde{C}_t, \mathbf{Q})_{\text{van}}$, the cycles $\delta'_1 - \delta''_1, ..., \delta'_r - \delta''_r$ span the antiinvariant subspace $H^1(\widetilde{C}_t, \mathbf{Q})_{\text{van}}^-$. The image of the monodromy representation

$$\tilde{\rho} \colon \pi_1(\mathbf{P}^1 \smallsetminus \{t_1, \dots, t_r\}, t) \longrightarrow \operatorname{Sp}(H^1(\overline{C}_t, \mathbf{Q}))$$

consists of automorphisms that are τ -equivariant and, reasoning as in the proof of [Voi03, Proposition 3.23], we see that, up to changing signs, the classes $\delta'_1 - \delta''_1, \ldots, \delta'_r - \delta''_r$ are all in the same monodromy orbit. Moreover, as in the proof of Proposition 3.4, there is for each $i \in \{1, \ldots, r\}$ an element of $\pi_1(\mathbf{P}^1 \setminus \{t_1, \ldots, t_r\}, 0)$ that acts on $H^1(\widetilde{C}_t, \mathbf{Q})$ by

$$T_i := T_{\delta_i''} \circ T_{\delta_i'} \colon x \longmapsto x - (x \cdot \delta_i')\delta_i' - (x \cdot \delta_i'')\delta_i''.$$

If *x* is τ -antiinvariant, we have

$$(x \cdot \delta_i'') = (x \cdot \tau^*(\delta_i')) = (\tau^*(x) \cdot \delta_i') = -(x \cdot \delta_i'),$$

hence

$$T_i(x) = x - (x \cdot \delta'_i)(\delta'_i - \delta''_i) = x - \frac{1}{2}(x \cdot (\delta'_i - \delta''_i))(\delta'_i - \delta''_i).$$

One then deduces from that and [PS03, Lemma 4] that the monodromy action on $H^1(\widetilde{C}_t, \mathbf{Q})^-_{\text{van}}$ is big; it follows that the Zariski closure of the monodromy group for $K(\widetilde{C}_t, C_t)$ is the full symplectic group $\text{Sp}(H^1(\widetilde{C}_t, \mathbf{Q})^-_{\text{van}})$. As in the proof of [PS03, Theorem 17], for $t \in \mathbf{P}^1$ very general, any endomorphism of $K(\widetilde{C}_t, C_t)$ intertwines every element of the monodromy group, hence every element of the symplectic group. It must therefore be a multiple of the identity.

The endomorphism ring of the abelian variety $K(\tilde{C}_t, C_t)$ is therefore trivial.

We now apply the theorem to GM fivefolds. Let X be a general GM fivefold with Lagrangian data set (V_6, V_5, A) . Our starting point is again the surjectivity, proved in Lemma 5.8, of the Abel-Jacobi map

$$\operatorname{AJ}_{\mathscr{L}^2_{\sigma}(X)} \colon H_1(F^2_{\sigma}(X), \mathbb{Z}) \longrightarrow H_5(X, \mathbb{Z})$$

associated with the Hilbert scheme $F_{\sigma}^{2}(X)$ that parametrizes σ -planes contained in X. This Hilbert scheme is isomorphic to the smooth curve $\widetilde{Y}_{A,V_{5}}^{\geq 2}$ (Lemma 2.10) defined as the inverse image by the double cover

$$\pi_A \colon \widetilde{Y}_A^{\geq 2} \longrightarrow Y_A^{\geq}$$

of the hyperplane section $Y_{A,V_5}^{\geq 2} = Y_A^{\geq 2} \cap \mathbf{P}(V_5)$. The surjectivity of the map $AJ_{\mathscr{L}^2_{\sigma}(X)}$ is therefore equivalent to the connectedness of the kernel of the induced surjective morphism

(5.14)
$$\Phi: \operatorname{Jac}(\widetilde{Y}_{A,V_5}^{\geq 2}) \longrightarrow \operatorname{Jac}(X)$$

between Jacobians. By Lemma 2.10 and Proposition 2.6, the dimension of this kernel is

$$g(\widetilde{Y}_{A,V_5}^{\geq 2}) - \dim(\operatorname{Jac}(X)) = 161 - 10 = 151.$$

Corollary 5.11. Let X be a smooth GM fivefold with Lagrangian data set (V_6, V_5, A) . Assume that the surface $\widetilde{Y}_A^{\geq 2}$ and the curve $\widetilde{Y}_{A,V_5}^{\geq 2}$ are smooth. The morphism Φ from (5.14) then factors as

$$\Phi: \operatorname{Jac}(\widetilde{Y}_{A,V_5}^{\geq 2}) \twoheadrightarrow \operatorname{Alb}(\widetilde{Y}_A^{\geq 2}) \xrightarrow{\sim} \operatorname{Jac}(X),$$

where the left arrow is the Albanese map $Jac(\widetilde{Y}_{A,V_5}^{\geq 2}) = Alb(\widetilde{Y}_{A,V_5}^{\geq 2}) \rightarrow Alb(\widetilde{Y}_A^{\geq 2}).$

Proof. Since Alb $(Y_A^{\geq 2}) = 0$ (Proposition 2.5), the diagram (5.13) reads

(5.15)

$$K_{V_{5}} \longrightarrow P_{A,V_{5}} \longrightarrow \operatorname{Alb}(\widetilde{Y}_{A}^{\geq 2}) \rightarrow 0$$

$$\downarrow \qquad \downarrow \qquad \parallel$$

$$0 \longrightarrow \widetilde{K}_{V_{5}} \longrightarrow \operatorname{Jac}(\widetilde{Y}_{A,V_{5}}^{\geq 2}) \rightarrow \operatorname{Alb}(\widetilde{Y}_{A}^{\geq 2}) \rightarrow 0$$

$$\downarrow \qquad \downarrow$$

$$\operatorname{Jac}(Y_{A,V_{5}}^{\geq 2}) = \operatorname{Jac}(Y_{A,V_{5}}^{\geq 2})$$

$$\downarrow \qquad \downarrow$$

$$0 \qquad 0.$$

where $K_{V_5} := K(\widetilde{Y}_{A,V_5}^{\geq 2}, Y_{A,V_5}^{\geq 2})$, $\widetilde{K}_{V_5} := K(\widetilde{Y}_{A,V_5}^{\geq 2}, \widetilde{Y}_A^{\geq 2})$, and P_{A,V_5} is the Prym variety of the double covering $\widetilde{Y}_{A,V_5}^{\geq 2} \to Y_{A,V_5}^{\geq 2}$. The genus of the curve $Y_{A,V_5}^{\geq 2}$ is 81 by (2.20), hence the dimension of the variety P_{A,V_5} is 80. Also, the dimension of Alb $(\widetilde{Y}_A^{\geq 2})$ is 10 by Proposition 2.5. Therefore, dim $(K_{V_5}) = 70$.

When V_5 is a very general hyperplane in V_6 , the abelian varieties $Jac(Y_{A,V_5}^{\geq 2})$ and K_{V_5} are simple by Proposition 3.4 and Theorem 5.10, hence they are the only two simple factors of the abelian variety \widetilde{K}_{V_5} . Since Jac(X) has dimension 10, the abelian variety \widetilde{K}_{V_5} must therefore be contained in the kernel of Φ .

In other words, the composition $\widetilde{K}_{V_5} \hookrightarrow \operatorname{Jac}(\widetilde{Y}_{A,V_5}^{\geq 2}) \xrightarrow{\Phi} \operatorname{Jac}(X)$ vanishes for V_5 very general. By continuity, it vanishes for all hyperplanes V_5 such that $\widetilde{Y}_{A,V_5}^{\geq 2}$ is smooth. The kernel of Φ , being connected of dimension 151, must then be equal to \widetilde{K}_{V_5} , which implies the corollary.

Note that this argument cannot be applied to GM threefolds, because the corresponding hyperplane sections $Y_{A,V_5}^{\geq 2}$ are very far from being general.

6. Period maps

In this section, we prove Theorems 1.1 and 1.3. We will use the description (2.8) of the coarse moduli space \mathbf{M}_n^{GM} of smooth GM varieties of dimension *n*. Let

(6.1)
$$\mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} = \mathrm{LGr}_{\mathrm{ndv}}(\wedge^{3} V_{6}) / / \mathrm{PGL}(V_{6})$$

be the coarse quasiprojective moduli space of EPW sextics defined by Lagrangian subspaces A with no decomposable vectors (see (2.3) for the definition). This is an open subset of the coarse moduli space \mathbf{M}^{EPW} of EPW sextics defined in (1.6). We denote by \mathbf{r} the involution of $\mathbf{M}_{\text{ndv}}^{\text{EPW}}$ defined by $\mathbf{r}([A]) = ([A^{\perp}])$. The morphism ρ_n was defined in (2.9).

Lemma 6.1. There exists a regular morphism $\bar{\wp}$: $\mathbf{M}_{ndv}^{\text{EPW}}/\mathbf{r} \to \mathbf{A}_{10}$ such that for $n \in \{3, 5\}$, the composition

(6.2)
$$\mathbf{M}_{n}^{\mathrm{GM}} \xrightarrow{\boldsymbol{\rho}_{n}} \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} \longrightarrow \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} / \mathbf{r} \xrightarrow{\boldsymbol{\varrho}} \mathbf{A}_{10}$$

is equal to the period map

$$\wp_n \colon \mathbf{M}_n^{\mathrm{GM}} \longrightarrow \mathbf{A}_{10}$$
$$[X] \longmapsto [\operatorname{Jac}(X)].$$

Proof. Consider the universal EPW variety

$$\mathscr{Y}_{\mathscr{A}^{\perp}}^{5-n} := \{ (A, V_5) \in \mathsf{LGr}_{\mathsf{ndv}}(\bigwedge^3 V_6) \times \mathbf{P}(V_6^{\vee}) \mid \dim(A \cap \bigwedge^3 V_5) = 5 - n \},$$

so that $\mathscr{Y}_{\mathscr{A}^{\perp}}^{5-n}//\operatorname{PGL}(V_6)$ is the coarse moduli space of ordinary GM varieties of dimension n, an open subscheme of $\mathbf{M}_n^{\mathrm{GM}}$. For $n \in \{3, 5\}$, the projection

$$\operatorname{pr}: \mathscr{Y}^{5-n}_{\mathscr{A}^{\perp}} \longrightarrow \operatorname{LGr}_{\operatorname{ndv}}(\bigwedge^{3} V_{6})$$

is a smooth surjective morphism. We show that the map

$$\tilde{\wp}_n \colon \mathscr{Y}_{\mathscr{A}^\perp}^{5-n} \longrightarrow \mathbf{A}_{10}
(A, V_5) \longmapsto [\operatorname{Jac}(X_{A, V_5})]$$

factors through pr. By smooth descent, it is enough for this to show that the two maps

$$\mathscr{Y}^{5-n}_{\mathscr{A}^{\perp}} \times_{\mathsf{LGr}_{\mathsf{ndv}}(\bigwedge^{3} V_{6})} \mathscr{Y}^{5-n}_{\mathscr{A}^{\perp}} \Longrightarrow \mathbf{A}_{10}$$

defined as compositions of the projections to the factors with the map $\tilde{\wp}_n$ are equal. Since the fiber product is a smooth variety, it is enough to verify the equality pointwise on an open subset. In other words, we need to check that for general GM varieties X, X' of the same dimension $n \in \{3, 5\}$ with A(X) = A(X'), there is an isomorphism $Jac(X) \simeq Jac(X')$ of principally polarized abelian varieties. For n = 3, this holds by Theorem 4.4, and for n = 5, by Theorem 5.3; in both cases, the intermediate Jacobians are isomorphic to $Alb(\widetilde{Y}_A^{\geq 2})$ as soon as A = A(X) = A(X') is such that $Y_A^{\geq 3} = \emptyset$. This proves that in both cases, the morphism $\tilde{\wp}_n$ factors as

$$\mathscr{Y}^{5-n}_{\mathscr{A}^{\perp}} \xrightarrow{\operatorname{pr}} \operatorname{LGr}_{\operatorname{ndv}}(\bigwedge^{3} V_{6}) \longrightarrow \mathbf{A}_{10}.$$

The maps $\tilde{\wp}_n$ are PGL(V_6)-invariant, hence so is the map

$$\operatorname{LGr}_{\operatorname{ndv}}(\bigwedge^{3} V_{6}) \longrightarrow \mathbf{A}_{10}$$

constructed via factorization. Therefore, it factors through a regular map

$$\mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} = \mathrm{LGr}_{\mathrm{ndv}}(\wedge^{3} V_{6}) / / \mathrm{PGL}(V_{6}) \longrightarrow \mathbf{A}_{10}$$

Similarly, this map is r-invariant by Theorem 4.4, hence we obtain a regular morphism

$$\bar{\wp}: \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}}/\mathbf{r} \longrightarrow \mathbf{A}_{10}$$

The composition (6.2) agrees with the period map by construction.

Recall that we denoted by $Alb(\widetilde{Y}_A^{\geq 2})$ the Albanese variety of (any desingularization of) the double EPW surface $\widetilde{Y}_A^{\geq 2}$.

Proposition 6.2. For any Lagrangian $[A] \in LGr_{ndv}(\bigwedge^3 V_6)$, we have $\bar{\wp}([A]) = [Alb(\widetilde{Y}_A^{\geq 2})]$.

Proof. If $Y_A^3 = \emptyset$, the equality holds by Theorems 4.4 and 5.3, and Lemma 6.1.

Assume $Y_A^3 \neq \emptyset$. Consider a neighborhood $U \subset \mathsf{LGr}_{ndv}(\bigwedge^3 V_6)$ of [A] such that the determinant of the tautological bundle \mathscr{A} is trivial over U. Consider a universal family $\mathscr{Y}_{\mathscr{A}}^{\geq 2} \to U$ of EPW surfaces over it and the composition

$$p: \widetilde{\mathscr{Y}}_{\mathscr{A}}^{\geq 2} \xrightarrow{\pi} \mathscr{Y}_{\mathscr{A}}^{\geq 2} \longrightarrow U,$$

where π is the double covering constructed as in the proof of [DK20a, Theorem 5.2(2)]. This proof implies that $\widetilde{\mathcal{Y}}_{\mathscr{A}}^{\geq 2}$ is a smooth variety. The fiber of p over [A] is the surface $\widetilde{Y}_{A}^{\geq 2}$, which is smooth away from a finite number of ordinary double points ([DK20a, Theorem 5.2(2)]).

Let $C \subset U$ be a general smooth affine curve passing through the point [A], so that its tangent space at [A] lies outside the finitely many hyperplanes which are the images of the differential of p at the various singular points of the fiber $p^{-1}([A])$. Upon shrinking U, the base change

$$\widetilde{\mathscr{Y}}_C^{\geq 2} := \widetilde{\mathscr{Y}}_{\mathscr{A}}^{\geq 2} \times_U C$$

is then a smooth threefold and the morphism $p_C \colon \widetilde{\mathscr{Y}}_C^{\geq 2} \to C$ is smooth over the complement of the point $[A] \in C$.

Consider a double covering $\widetilde{C} \to C$ branched at [A]. The base change

$$\widetilde{\mathscr{Y}}_{\widetilde{C}}^{\geq 2} := \widetilde{\mathscr{Y}}_{C}^{\geq 2} \times_{C} \widetilde{C}$$

is a threefold with finitely many ordinary double points in the central fiber. By [Ati58], there is an analytic simultaneous resolution $(\widetilde{\mathcal{Y}}_{\widetilde{C}}^{\geq 2})' \to \widetilde{\mathcal{Y}}_{\widetilde{C}}^{\geq 2}$ such that the composition

$$p' \colon (\widetilde{\mathscr{Y}}_{\widetilde{C}}^{\geq 2})' \longrightarrow \widetilde{\mathscr{Y}}_{\widetilde{C}}^{\geq 2} \longrightarrow \widetilde{C}$$

is a smooth morphism with central fiber isomorphic to the (smooth) blow up $(\widetilde{Y}_{\widetilde{C}}^{\geq 2})' \to \widetilde{Y}_{A}^{\geq 2}$ of the singular points of $\widetilde{Y}_{A}^{\geq 2}$. The sheaf $R^1p'_*(\underline{Z})$ is then locally constant and its stalk at [A] is isomorphic to $H^1((\widetilde{Y}_{\widetilde{C}}^{\geq 2})', \mathbf{Z})$.

By Theorems 4.4 and 5.3, this sheaf carries, away from the point [A], a variation of Hodge structure that comes from the middle cohomology of a family of smooth projective varieties of odd dimension, hence it has a canonical principal polarization. Since the sheaf $R^1 p'_*(\underline{Z})$ is locally constant on the whole \widetilde{C} , this polarization extends across this point. In particular, the natural Hodge structure on the stalk $H^1((\widetilde{Y}_{\widetilde{C}}^{\geq 2})', \mathbf{Z})$ at [A] has a principal polarization, hence provides a principal polarization on the Albanese variety $Alb((\widetilde{Y}_{\widetilde{C}}^{\geq 2})')$ and the map $\widetilde{C} \to \mathbf{A}_{10}$ defined by the above variation takes the point [A] to $[Alb((\widetilde{Y}_{\widetilde{C}}^{\geq 2})')]$. Since this map agrees on $\widetilde{C} \setminus \{[A]\}$ with the composition

$$\widetilde{C} \longrightarrow C \longrightarrow U \longrightarrow \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}} \longrightarrow \mathbf{M}_{\mathrm{ndv}}^{\mathrm{EPW}}/\mathbf{r} \xrightarrow{\widetilde{\wp}} \mathbf{A}_{10},$$

it agrees everywhere, hence $\bar{\wp}([A]) = [\operatorname{Alb}((\widetilde{Y}_{\widetilde{C}}^{\geq 2})')].$

We now use these results to prove Theorem 1.3.

Proof of Theorem 1.3. The factorization of the period map \mathscr{P}_n is proved in Lemma 6.1 and the equality $\mathscr{P}_n([X]) = [\operatorname{Alb}(\widetilde{Y}_{A(X)}^{\geq 2})]$ follows from this factorization and Proposition 6.2.

Remark 6.3. Consider the natural action of $PGL(V_6)$ on $LGr_{ndv}(\bigwedge^3 V_6) \times \mathbf{P}(V_6^{\vee})$, linearized as in [DK20b, Section 5.4]. For each $n \in \{3, 4, 5, 6\}$, there is by [DK20b, Theorem 5.15] a canonical embedding

(6.3)
$$\mathbf{M}_n^{\mathrm{GM}} \subset (\mathsf{LGr}_{\mathrm{ndv}}(\bigwedge^3 V_6) \times \mathbf{P}(V_6^{\vee})) /\!\!/ \operatorname{PGL}(V_6)$$

and $(LGr_{ndv}(\wedge^{3}V_{6})\times \mathbb{P}(V_{6}^{\vee}))//PGL(V_{6}) \to \mathbb{M}_{ndv}^{EPW}$ is generically a \mathbb{P}^{5} -fibration (the fiber over any point [A] is isomorphic to $\mathbb{P}(V_{6}^{\vee})//PGL(V_{6})_{A}$). The inclusion (6.3) is an open embedding for n = 5, a closed embedding for n = 3, and

$$(\operatorname{LGr}_{\operatorname{ndv}}(\bigwedge^{3} V_{6}) \times \mathbf{P}(V_{6}^{\vee})) / / \operatorname{PGL}(V_{6}) = \mathbf{M}_{5}^{\operatorname{GM}} \sqcup \mathbf{M}_{3}^{\operatorname{GM}}$$

by (2.10). This property is reminiscent of the Satake compactification.

We can also complete the proof of Theorem 1.1.

Proof of Theorem 1.1. By Proposition 6.2, the morphism $\bar{\wp}$ defines for each Lagrangian A with no decomposable vectors a principal polarization on Alb $(\tilde{Y}_A^{\geq 2})$ such that (1.2) holds; this proves the first part of the theorem.

By Lemma 6.1, the isomorphism (1.4) of principally polarized abelian varieties holds for all smooth GM varieties X of dimension $n \in \{3, 5\}$; this proves the last part of the theorem.

Finally, the isomorphism (1.3) for A with $Y_A^3 = \emptyset$ was established in Theorems 4.4 and 5.3. For A with $Y_A^3 \neq \emptyset$, the proof of Proposition 6.2 gives an isomorphism

$$H_n(X, \mathbb{Z}) \simeq H_1((\widetilde{Y}_{A(X)}^{\geq 2})', \mathbb{Z}),$$

where $(\widetilde{Y}_{A(X)}^{\geq 2})'$ is a desingularization of $\widetilde{Y}_{A(X)}^{\geq 2}$. Since the only singularities of $\widetilde{Y}_{A}^{\geq 2}$ are ordinary double points, there is a canonical isomorphism $H_1((\widetilde{Y}_A^{\geq 2})', \mathbb{Z}) \xrightarrow{\sim} H_1(\widetilde{Y}_A^{\geq 2}, \mathbb{Z})$. This proves the second part of the theorem.

References

[Ati58] M. F. Atiyah, On analytic surfaces with double points, Proc. Roy. Soc. London 247 (1958), 237-244.

- [BL99] C. Birkenhake, H. Lange, Complex tori, Progress in Mathematics 177, Birkhäuser Boston, Inc., Boston, MA, 1999.
- [BFM19] M. Bernardara, E. Fatighenti, L. Manivel, Nested varieties of K3 type, preprint arXiv:1912.03144 (2019).
- [BM79] S. Bloch, J.P. Murre, On the Chow group of certain types of Fano threefolds, Compos. Math. 39 (1979), 47-105.
- [CvG92] C. Ciliberto, G. van der Geer, On the Jacobian of a hyperplane section of a surface, in: Classification of irregular varieties (Trento, 1990), 33-40, Lecture Notes in Math. 1515, Springer, Berlin, 1992.
- [Deb20] O. Debarre, Gushel-Mukai varieties, preprint arXiv:2001.03485 (2020).
- [DIM12] O. Debarre, A. Iliev, L. Manivel, On the period map for prime Fano threefolds of degree 10, J. Algebraic Geom. 21 (2012), 21-59.
- [DK18] O. Debarre, A. Kuznetsov, Gushel-Mukai varieties: classification and birationalities, Algebr. Geom. 5 (2018), 15-76.
- [DK19] _____, Gushel-Mukai varieties: linear spaces and periods, Kyoto J. Math. 59 (2019), 897-953.
- [DK20a] _____, Double covers of quadratic degeneracy and Lagrangian intersection loci, Math. Ann. 378 (2020), 1435–1469.
- [DK20b] _____, Gushel-Mukai varieties: moduli, Internat. J. Math. 31 (2020) 2050013.
- [DK20+] _____, Gushel-Mukai varieties: quadrics, in preparation.
- [Del71] P. Deligne, Théorie de Hodge. II, Inst. Hautes Études Sci. Publ. Math. 40 (1971), 5-57.
- [Fer12] A. Ferretti, Special subvarieties of EPW sextics, Math. Z. 272 (2012), 1137-1164.

- [FW08] B. Fu, C.-L. Wang, Motivic and quantum invariance under stratified Mukai flops, J. Differential Geom. 80 (2008), 261–280.
- [IM11] A. Iliev, L. Manivel, Fano manifolds of degree 10 and EPW sextics, Ann. Sci. École Norm. Sup. 44 (2011), 393-426.
- [IM07] _____, Prime Fano threefolds and integrable systems, Math. Ann. 339 (2007), 937–955.
- [IP99] V. A. Iskovskikh, Yu. Prokhorov, Fano varieties, in: Algebraic geometry, V, 1-247, Encyclopaedia Math. Sci. 47, Springer-Verlag, Berlin, 1999.
- [Kuz16] A. Kuznetsov, Küchle fivefolds of type c5, Math. Z. 284 (2016), 1245-1278.
- [KP18] A. Kuznetsov, A. Perry, Derived categories of Gushel-Mukai varieties, Compos. Math. 154 (2018), 1362-1406.
- [KP19] _____, Categorical cones and quadratic homological projective duality, preprint arXiv:1902.09824 (2019).
- [KPS18] A. Kuznetsov, Yu. Prokhorov, C. Shramov, Hilbert schemes of lines and conics and automorphism groups of Fano threefolds, Jpn. J. Math. 13 (2018), 109–185.
- [KS18] A. Kuznetsov, E. Shinder, Grothendieck ring of varieties, D- and L-equivalence, and families of quadrics, Selecta Math 24 (2018), 3475-3500.
- [Log12] D. Logachev, Fano threefolds of genus 6, Asian J. Math. 16 (2012), 515-560.
- [Mar81] D. G. Markushevic, Numerical invariants of families of lines on some Fano varieties, (in Russian) Mat. Sb. 116(158) (1981), no. 2(10), 265–288. English translation: Math. USSR Sbornik 44 (1983), no. 2, 239–260.
- [Nag98] J. Nagel, The generalized Hodge conjecture for the quadratic complex of lines in projective four-space, Math. Ann. 312 (1998), 387-401.
- [O'Gr08a] K. O'Grady, Irreducible symplectic 4-folds numerically equivalent to (K3)^[2], Commun. Contemp. Math. 10 (2008), 553-608.
- [O'Gr08b] _____, Dual double EPW-sextics and their periods, Pure Appl. Math. Q, 4 (2008), 427-468.
- [O'Gr13] _____, Double covers of EPW-sextics, Michigan Math. J. 62 (2013), 143-184.
- [PS83] C. A. M. Peters, J. H. M. Steenbrink, Infinitesimal variations of Hodge structure and the generic Torelli problem for projective hypersurfaces (after Carlson, Donagi, Green, Griffiths, Harris), in: Classification of algebraic and analytic manifolds (Katata, 1982), 399-463, Progr. Math. 39, Birkhäuser Boston, Boston, MA, 1983.
- [PS03] _____, Monodromy of variations of Hodge structure, Monodromy and differential equations (Moscow, 2001), Acta Appl. Math. 75 (2003), 183-194.
- [Tju72] A. N. Tjurin, Five lectures on three-dimensional varieties (in Russian), Uspehi Mat. Nauk 27 (1972), no. 5, (167), 3-50. English translation: Russian Math. Surveys 27 (1972), 1-53.
- [Voi88] C. Voisin, Sur la jacobienne intermédiaire du double solide d'indice deux, Duke Math. J. 57 (1988), 629-646.
- [Voi03] _____, Hodge theory and complex algebraic geometry. II, Translated from the French by Leila Schneps. Cambridge Studies in Advanced Mathematics 77, Cambridge University Press, Cambridge, 2003.
- [Wel81] G. E. Welters, Abel-Jacobi isogenies for certain types of Fano threefolds, Mathematical Centre Tracts 141, Mathematisch Centrum, Amsterdam, 1981.